

FILTERED FROBENIUS ALGEBRAS IN MONOIDAL CATEGORIES

talk by Harshit Yadav

(arXiv: 2106.01999
joint work Chelsea Walton)

① Motivation

(i) From noncommutative ring theory

Filtered algebras

$$A_0 \subset A_1 \subset A_2 \dots$$

$$A = \bigcup_{i \in \mathbb{N}_0} A_i$$

$$A_i \cdot A_j \subset A_{i+j}$$

Graded algebras

$$B = \bigoplus B_i$$

$$B_i \cdot B_j \subset B_{i+j}$$

$$\bullet \quad A \xrightarrow{\text{gr}} \text{gr}(A) = \bigoplus_{i \in \mathbb{N}} \frac{A_i}{A_{i-1}} \quad \left(\begin{array}{l} \text{associated} \\ \text{graded} \\ \text{algebra} \\ \text{of } A \end{array} \right)$$

• A is called a filtered deformation of $\text{gr}(A)$.

→ $\text{gr}(A)$ is a graded algebra

Example:

A	$\text{gr}(A)$
$U(\mathfrak{g})$	$S(\mathfrak{g})$
$U(V, B)$	$\wedge(V)$

In fact, many nice properties of $\text{gr}(A)$ transfer to A.

If $\text{gr}(A)$ is integral domain
Noetherian
prime then so is $\text{gr}(A)$.

The talk is about the property of being Frobenius.

But, what are Frobenius algebras and why should we care about them?

This brings us to the second motivation

(ii) From Quantum algebra

Frobenius algebras in monoidal category

$$= (\mathcal{C}, \otimes, \mathbb{1})$$

(like vector spaces) • $\mathcal{C} =$ a category

(like tensor of vector spaces) • $\otimes: \mathcal{C} \times \mathcal{C} \rightarrow \mathcal{C}$ bifunctor

(like ground field \mathbb{k}) • $\mathbb{1}$: unit object of \mathcal{C}
unit for \otimes product




(allows us to do algebra in \mathcal{C})

WHAT?

A Frobenius algebra in $(\mathcal{C}, \otimes, \mathbb{1})$ is a 5-tuple

- algebra in \mathcal{C}
- $A =$ object in \mathcal{C}
 - $m = A \otimes A \rightarrow A$
 - $\eta: \mathbb{1} \rightarrow A$
 - $\Delta: A \rightarrow A \otimes A$
 - $\epsilon: A \rightarrow \mathbb{1}$

coalgebra in \mathcal{C} +



Example: $\mathbb{k}G$ with $\Delta(g) = \sum_{h \in G} gh^{-1} \otimes h$
 $\epsilon(g) = \delta_{g, e_G}$
 is a Frobenius algebra in $\mathcal{C} = \text{Vec}_{\mathbb{k}}$.

WHY?

Frobenius algebras in monoidal categories, show up in work on

- (i) TQFTs and CFTs
- (ii) Morita theory
- (iii) Classification of subfactors
- (iv) Computer Science

With these motivations in mind
the following NC ring theory result provides a context
for our work.

Theorem (Bongale, 1967)

($A_0 = k$)
Let A be a finite dimensional connected, filtered algebra
over k . If $\text{gr}(A)$ is Frobenius, then so is A .

we generalize this to get our main result

MAIN THEOREM (Walton-Y., 21)

Let \mathcal{C} be an abelian, rigid monoidal category. Let
 A be a connected, filtered algebra in \mathcal{C} with
finite filtration. If $\text{gr}(A)$ is a Frobenius algebra
in \mathcal{C} , then so is A .

As an application of this, we are able to
prove that.

Theorem (Walton-Y.)

Every exact module category \mathcal{M} over a symmetric
finite tensor category \mathcal{C} is represented by a
Frobenius algebra A in \mathcal{C} , i.e., $\mathcal{M} = \mathcal{C}_A$.

(more details at the end)

Let's come back to proving the main theorem.
We need to develop two tools to prove it.

- (i) Associated graded algebra construction
- (ii) New characterization of Frobenius algebras.

(i) Associated graded functor

For \mathcal{C} an abelian, monoidal category with \otimes biexact.
We construct a monoidal associated graded functor

$$\text{gr}: \text{Fil}(\mathcal{C}) \longrightarrow \text{Gr}(\mathcal{C})$$

$$(A, F_A) \longmapsto \coprod_{i \in \mathbb{N}_0} \frac{F_A(i)}{F_A(i-1)}$$

Objects: (X, F_X) with $X \in \mathcal{C}$,
 $F_X: \mathbb{N}_0 \rightarrow \mathcal{C}$ s.t.
 $\text{colim}_i F_X(i) = X$

Morphisms: $X \xrightarrow{f} Y$ in \mathcal{C} that
 preserve filtration

$(X, F_X) \otimes (Y, F_Y) := (X \otimes Y, \text{Day convolution of } F_X, F_Y)$

Objects: $X = \coprod_{i \in \mathbb{N}_0} X_i$

Morphisms: $X \rightarrow Y$ compatible
 with \coprod
 decomposition

$(X \otimes Y)_k = \coprod_{i+j=k} (X_i \otimes Y_j)$

UPSHOT If A is a filtered algebra in \mathcal{C} , then
 $\text{gr}(A)$ is a graded algebra in \mathcal{C} .

(ii) New characterization of Frobenius algebras

Theorem

Let \mathcal{C} be a rigid monoidal category. An algebra
 (A, m, u) in \mathcal{C} is Frobenius
 if and only if

$\exists v: A \rightarrow \mathbb{1}$ such that any left/right ideal
 of A that factors through $\ker(v)$ is zero.

Proof sketch of the main theorem:

- (A, F_A) has finite filtration $\Rightarrow A \cong F_A(n)$ for some $n \in \mathbb{N}$

- Take $\nu: A \xrightarrow{\sim} F_A(n) \rightarrow \frac{F_A(n)}{F_A(n)} \cong \mathbb{1}$ ↙ because A is connected

- Take any ideal I of A so that

$$\begin{array}{ccc} \ker(\nu) & \longrightarrow & A \xrightarrow{\nu} \mathbb{1} \\ \uparrow & \nearrow \phi & \\ I & & \end{array}$$

(by our characterization of Frobenius algs, it suffices to) show that $I=0$

- Consider $\text{gr}(\phi): \text{gr}(I) \rightarrow \text{gr}(A)$

- Show $\text{gr}(\phi)$ factors through the kernel of the Frobenius form on $\text{gr}(A)$. Hence,
 $\text{gr}(I) = 0$

- Hence, $I=0$.

x — x

Directions for future work

- ① Generalize and study other ring theoretical properties that lift under filtered deformations
- ② Construct braided Clifford algebras and show that they are Frobenius by showing that its associated graded is the exterior algebra.

Details of the application

Take $\mathcal{C} =$ symmetric finite tensor category
 $\mathcal{M} =$ exact module category over \mathcal{C}

Etingof - Ostrik showed that $\mathcal{M} = \mathcal{C}_A$ where

$$A = \text{Ind}_{\mathbb{H}}^{\mathbb{H}} \left(\text{Ind}_{\mathbb{H}}^{\mathbb{H}} (\text{End}(V)) \otimes \text{Cl}_W \right)$$

Key idea : $\text{gr}(\text{Cl}_W) = \Lambda_W$ is the exterior algebra

Λ_W is Frobenius

$\Rightarrow \text{Cl}_W$ is Frobenius

- Clearly $\text{End}(V)$ is also Frobenius
- $\text{Ind}_{\mathbb{H}}^{\mathbb{H}}$ are Frobenius monoidal
- \otimes of Frob. algs is Frobenius

$\Rightarrow A$ is Frobenius.