

POINCARÉ DUALITY FOR TAUTOLOGICAL CHERN SUBRINGS OF ORTHOGONAL GRASSMANNIANS

NIKITA A. KARPENKO AND ALEXANDER S. MERKURJEV

ABSTRACT. Let X be an orthogonal grassmannian of a nondegenerate quadratic form q over a field. Let C be the subring in the Chow ring $\mathrm{CH}(X)$ generated by the Chern classes of the tautological vector bundle on X . We prove Poincaré duality for C . For q of odd dimension, the result was already known due to an identification between C and the Chow ring of certain symplectic grassmannian. For q of even dimension, such an identification is not available.

1. INTRODUCTION

Let q be an arbitrary nondegenerate quadratic form over an arbitrary field F of an arbitrary dimension $d = \dim q$ (see [2, §7.A] for the definition of a nondegenerate quadratic form in any characteristic including 2). We take any integer m with $1 \leq m \leq d/2$ and consider the Chow ring $\mathrm{CH}(X_m)$ of the orthogonal grassmannian variety X_m of totally isotropic m -planes of q . Here we use the affine numbering of grassmannians so that X_1 is the projective quadric given by q .

Assume that the form q is split (it is automatically split if the field F is algebraically closed). Then the group $\mathrm{CH}(X_m)$ is a free abelian group of finite rank. It does not depend on the field F and has a canonical basis given by Schubert classes. The bilinear form $(a, b) \mapsto \deg(ab) \in \mathbb{Z}$ on $\mathrm{CH}(X_m)$, where $\deg: \mathrm{CH}(X_m) \rightarrow \mathbb{Z}$ is the degree map (vanishing on the homogeneous components of $\mathrm{CH}(X_m)$ other than the component $\mathrm{CH}_0(X_m)$ given by the 0-cycles), turns out to be unimodular in this split case. In other words, $\mathrm{CH}(X_m)$ satisfies Poincaré duality with respect to the bilinear form. Moreover, the Schubert basis is *weekly self-dual*, i.e., dual to a permutation of itself. The unimodularity can also be explained by the fact that the variety X_m is (absolutely) cellular (see [2, §66]). Therefore its Chow motive is split (see [2, Corollary 66.4]). The latter fact implies the Poincaré duality, c.f. [7, Remark 5.6] or [9, Proposition 1.5].

In general (for arbitrary F and nonsplit q), the change of field homomorphism

$$(1.1) \quad \mathrm{CH}(X_m) \rightarrow \mathrm{CH}((X_m)_K),$$

for a field extension K/F is neither injective nor surjective. Nevertheless, the ring $\mathrm{CH}(X_m)$ contains a “core” subring C , called the *tautological Chern subring*, generated by the

Date: 20 Nov 2021.

Key words and phrases. Algebraic groups; projective homogeneous varieties; Chow groups. *Mathematical Subject Classification (2010):* 20G15; 14C25.

The first author was supported by a Discovery Grant from the National Science and Engineering Research Council of Canada and a Centennial Grant from the University of Alberta. The second author was supported by the NSF grant DMS #1801530.

Chern classes of the tautological (rank m) vector bundle on X_m . By [5, Theorem 2.1], the homomorphism (1.1) maps C isomorphically onto “itself” – the tautological Chern subring of $\mathrm{CH}((X_m)_K)$. In fact, C is determined by the integers d and m alone and does not depend on F and q . Note that the additive group of C is free of finite rank and that $\deg(C) = 2^m \mathbb{Z}$ (see, e.g., [5, Theorem 2.1] and [12, Statement 2.15]). By [5, Theorem 6.1], if the quadratic form q is *generic*, the subring C coincides with the entire Chow ring of X_m .

Not only the ring C itself, but also the restriction of the degree map $\deg: \mathrm{CH}(X_m) \rightarrow \mathbb{Z}$ to $C \subset \mathrm{CH}(X_m)$ is determined by d and m . We consider the bilinear form on C given by the modified degree map $2^{-m} \deg: C \rightarrow \mathbb{Z}$ and show that C satisfies Poincaré duality with respect to it:

Theorem 1.2. *The bilinear form*

$$C \times C \rightarrow \mathbb{Z}, \quad (a, b) \mapsto 2^{-m} \deg(a \cdot b)$$

is unimodular. In other terms, every \mathbb{Z} -basis for C admits a dual basis.

The idea of considering Poincaré duality for C originated in [11, §4], where it was proven for $m = \lfloor d/2 \rfloor$ and then successfully applied in order to determine the torsion index of spin groups.

The unimodularity means invertibility of the determinant of the Gramm matrix. Therefore, replacing the coefficient ring \mathbb{Z} by an arbitrary field k we get an equivalent version of Theorem 1.2:

Theorem 1.3. *For any field k , the bilinear form on the finite dimensional k -vector space $C \otimes k$, given by $2^{-m} \deg$, is nondegenerate.*

Nondegeneracy in Theorem 1.3 is the precise analogue of unimodularity in Theorem 1.2. Both mean that the induced map of the space to its dual is an isomorphism. But since the coefficient ring in Theorem 1.3 is a field, it is enough to only have injectivity of the map. In other terms, it is enough to check triviality of the radical.

For odd d , Theorems 1.2 and 1.3 have already been shown in [4, Proof of Theorem 4.1] by the argument of [11, §4] – using an identification of C with the entire Chow ring of the m -th symplectic grassmannian Y_m of a nondegenerate alternating bilinear form of dimension $\dim q - 1$. It also identifies the modified degree map on C with the usual degree map on $\mathrm{CH}(Y_m)$. The Schubert classes form a basis for the additive group of $\mathrm{CH}(Y_m)$ and this basis is weakly self-dual.

Possession of a self-dual basis is a stronger property than just the unimodularity. To prove the latter, one can argue (as we already did for $\mathrm{CH}(X_m)$ with split q) without involving the Schubert basis: since the variety Y_m is cellular, its Chow motive is split implying the integral Poincaré duality for $\mathrm{CH}(Y_m)$.

For even d however, an identification like above is not available (aside from the extreme values 1 and $d/2$ of m – see Examples 1.4 and 1.5) and therefore a different approach is needed. A proof of Theorem 1.3 for even d is given in §2.

Example 1.4. For even $d = 2n + 2$ and split q , the quadric X_1 has dimension $2n$ and the Schubert basis for $\mathrm{CH}(X_1)$ consists of the powers h^i for $i = 0, 1, \dots, n - 1$ of the class h of a hyperplane section, the classes l_i of linear subspaces of dimensions $i = 0, 1, \dots, n - 1$,

and two distinct classes l'_n and l''_n of linear subspaces of dimension n (see [2, §68]). The subring $C \subset \text{CH}(X_1)$ has a weakly self-dual basis consisting of h^i for $i = 0, 1, \dots, 2n$. Note that $h^n = l'_n + l''_n$ and $h^{2n-i} = 2l_i$ for $i = 0, 1, \dots, n-1$. The ring C endowed with the modified degree map $2^{-1} \text{deg}: C \rightarrow \mathbb{Z}$, is isomorphic to the entire Chow ring of the projective space \mathbb{P}^{2n} , endowed with its usual degree map.

Example 1.5. For $m = d/2$ (with even d) and split q , the variety X_m has two connected components isomorphic (to each other as well as) to the $(m-1)$ -st grassmannian X'_{m-1} of any nondegenerate $(d-1)$ -dimensional subform q' of q . Via this isomorphism, the tautological Chern subring of $\text{CH}(X_m)$ together with its modified (by 2^m) degree map is identified with the tautological Chern subring of $\text{CH}(X'_{m-1})$ endowed with its modified (by 2^{m-1}) degree map. As already discussed, the latter can be identified with the entire Chow ring of the corresponding symplectic grassmannian.

Theorems 1.2 and 1.3 for odd d are crucial for [4, Theorem 4.1] providing an algorithm for computation of indexes of generic $\text{Spin}(d)$ -grassmannians. A similar algorithm for even d is not available yet; we expect that our results will help to obtain it.

2. PROOF OF THEOREM 1.3 FOR EVEN d

Below we provide a proof of the Poincaré duality with coefficients in a field k for even $d = 2n$ by reduction to the duality for $d = 2n + 1$.

We choose a nondegenerate $(2n+1)$ -dimensional quadratic form q , containing a given nondegenerate $(2n)$ -dimensional quadratic form q' as a subform. Writing X'_m for the m -th orthogonal grassmannian of q' , we have a closed imbedding $in: X'_m \hookrightarrow X_m$. The tautological vector bundle \mathcal{T}' on X'_m is the pull-back of the tautological vector bundle \mathcal{T} on X_m . Therefore the pull-back homomorphism of Chow rings $in^*: \text{CH}(X_m) \rightarrow \text{CH}(X'_m)$ maps C surjectively onto C' . Note that by [5, Theorem 2.1], $in^*(c_{2n-m}(-\mathcal{T})) = 0$. Here and below we write $c_i(\mathcal{E})$ for the i -th Chern class of a vector bundle \mathcal{E} and we write $c_i(-\mathcal{E})$ for the i -th Chern class of $-\mathcal{E}$ (or, in other terms, for the i -th Segre class of \mathcal{E} , see [3, Chapter 3]).

Lemma 2.1. *The class $[X'_m] \in \text{CH}(X_m)$ of $X'_m \subset X_m$ coincides with $c_m(\mathcal{T})$.*

Proof. Let V be the vector space of definition of q and let $V' \subset V$ be the hyperplane on which q' is defined. The grassmannian $\Gamma_m(V')$ of m -planes in V' is a closed subvariety in the grassmannian $\Gamma_m(V)$ and $[\Gamma_m(V')] \in \text{CH}(\Gamma_m(V))$ is the m -th Chern class of the tautological vector bundle on $\Gamma_m(V)$, [3, §14.7]. We have a pull-back square of closed embeddings:

$$\begin{array}{ccc} \Gamma_m(V') & \longrightarrow & \Gamma_m(V) \\ \uparrow & & \uparrow \\ X'_m & \longrightarrow & X_m \end{array}$$

Since $\dim \Gamma_m(V) - \dim \Gamma_m(V') = m = \dim X_m - \dim X'_m$, the statement follows from [2, Corollary 57.20]. \square

In particular, it follows that $in_*(C') \subset C$: any element of C' has the form $in^*(x)$ with $x \in C$ and $in_*(in^*(x)) = c_m(\mathcal{T}) \cdot x \in C$ by projection formula.

The push-forward map $in_*: \text{CH}(X'_m) \rightarrow \text{CH}(X_m)$ satisfies $\text{deg} \circ in_* = \text{deg}'$ with

$$\text{deg}': \text{CH}(X'_m) \rightarrow \mathbb{Z}$$

the degree map of X'_m .

We now switch to coefficients k considerations. As in Theorem 1.3, we assume that k is a field.

Let $in_* \otimes k$ be the additive homomorphism $C' \otimes k \rightarrow C \otimes k$ given by in_* . By projection formula, Poincaré duality with coefficients k for C' will follow from that for C once we know that $in_* \otimes k$ is injective. Indeed, given any nonzero $a' \in C' \otimes k$, we find $a \in C \otimes k$ non-orthogonal to $in_*(a')$ and get that a' is not orthogonal to $in^*(a)$.

Since $C' = in^*(C)$, the injectivity we need is implied by the following property of C alone (making no reference to C' anymore):

Proposition 2.2. *Any element $f \in C \otimes k$, vanishing under multiplication by $c_m(\mathcal{T})$, is divisible by $c_{2n-m}(-\mathcal{T})$.*

Proof. In the beginning of this proof we allow k to be an arbitrary commutative ring. As already mentioned in §1, C is identified with the Chow ring $\text{CH}(Y_m)$ of the m -th grassmannian of a $2n$ -dimensional nondegenerate alternating bilinear form b . Under this identification, the Chern classes in C of the tautological vector bundle on X_m correspond to the respective Chern classes in $\text{CH}(Y_m)$ of the tautological vector bundle $'\mathcal{T}$ on Y_m .

We have $Y_m = G/P$ for certain parabolic subgroup $P \subset G := \text{Sp}(2n)$. Namely, P is the standard maximal parabolic subgroup given by the m -th vertex of the Dynkin diagram of $\text{Sp}(2n)$. Therefore $C \otimes k$ is identified with the coefficient k version

$$\text{Ch}(G/P) := \text{CH}(G/P) \otimes k$$

of the Chow ring.

We are going to operate with the *Chow rings of the classifying spaces* of algebraic groups introduced in [10]. The Weyl group W of P acts on the Chow ring $\text{Ch}(BT)$ of the classifying space BT of the standard split maximal torus $T \subset P$. The ring of W -invariants $\text{Ch}(BT)^W$ maps surjectively onto $\text{Ch}(G/P)$:

$$(2.3) \quad \text{Ch}(BT)^W \twoheadrightarrow \text{Ch}(G/P).$$

Since G has no torsion primes, it follows by [1, Corollaire 2] that the kernel is generated by the non-constant part of the image of the (injective) ring homomorphism $\text{Ch}(BG) \rightarrow \text{Ch}(BT)^W$ from the Chow ring of the classifying space of G .

The ring $\text{Ch}(BT)$ is identified with the polynomial ring

$$\tilde{R} := k[x_1, \dots, x_m, y_1, \dots, y_l]$$

in the n variables $x_1, \dots, x_m, y_1, \dots, y_l$, where $l := n - m$. We view \tilde{R} as a graded ring with respect to the standard grading $\text{deg } x_i = \text{deg } y_j = 1$. Write u_i for the i -th elementary symmetric polynomial in x_1, \dots, x_m ($i = 1, \dots, m$) and v_j for the j -th elementary symmetric polynomial in y_1^2, \dots, y_l^2 ($j = 1, \dots, l$). Let R be the subring $k[u_1, \dots, u_m, v_1, \dots, v_l]$ of \tilde{R} . The ring R is also a polynomial ring over k in $n = m + l$ variables. Since the Weyl group W acts on $\tilde{R} = \text{Ch}(BT)$ by interchanging the variables x_1, \dots, x_m , interchanging the variables y_1, \dots, y_l , and changing signs of the latter variables, we have $R = \text{Ch}(BT)^W$.

Write w_i for the i -th elementary symmetric polynomial in the squares

$$x_1^2, \dots, x_m^2, y_1^2, \dots, y_l^2$$

of our n variables ($i = 1, \dots, n$). Let $I \subset R$ be the ideal generated by w_1, w_2, \dots, w_n . The computation of $\text{CH}(BG)$ given in [10, §15] (see also [6, Example 5.2]) tells us that I is the kernel of the surjective ring homomorphism (2.3)

$$(2.4) \quad R = \text{Ch}(BT)^W \twoheadrightarrow \text{Ch}(G/P) = C \otimes k.$$

To have a more complete understanding of the objects involved, let us mention that $\text{Ch}(BG) = \text{Ch}(BT)^{W_G} = \tilde{R}^{W_G}$, where W_G is the Weyl group of G , and w_1, \dots, w_n generate this ring of invariants. The quotient of \tilde{R} by the ideal $I\tilde{R}$, generated by w_1, \dots, w_n , is therefore the Chow ring $\text{Ch}(G/B)$ for the standard Borel subgroup $B \subset G$.

Consider the monomials $u := u_m = x_1 \cdots x_m \in R$ and $w := u_m v_l = u y_1^2 \cdots y_l^2 \in R$. Note that $w_n = uw$. Besides, the image of u under (2.4) is $c_m(\mathcal{T})$ whereas the image of w is $(-1)^n c_{2n-m}(-'\mathcal{T})$. In order to justify the statement on the image of w , let us mention that whereas the images of x_1, \dots, x_m are the roots of the vector bundle $'\mathcal{T}$, the roots of the containing $'\mathcal{T}$ trivial vector bundle, given by the $2n$ -dimensional vector space V_b of definition of our alternating bilinear form b , are $\pm x_1, \dots, \pm x_m, \pm y_1, \dots, \pm y_l$. Therefore $c_{2n-m}(-'\mathcal{T}) = c_{2n-m}(V_b/'\mathcal{T})$ is computed as claimed.

Proposition 2.2 translates as follows:

$$(2.5) \quad \text{Let } f \in R \text{ be a polynomial such that } uf \in I. \text{ Then } f \in wR + I.$$

To prove (2.5), since I is a homogeneous ideal, we may assume that the polynomial f is also homogeneous. By assumption,

$$uf = w_1 f_1 + w_2 f_2 + \dots + w_n f_n$$

for some $f_1, f_2, \dots, f_n \in R$, or equivalently,

$$(2.6) \quad u f_0 + w_1 f_1 + w_2 f_2 + \dots + w_{n-1} f_{n-1} = 0,$$

where $f_0 = w f_n - f$.

From now on k is assumed to be a field. Let $M \subset R$ be the ideal generated by $u_1, \dots, u_m, v_1, \dots, v_l$. Clearly, M is a maximal ideal of R containing I .

We claim that M is the radical \sqrt{I} of I . For every polynomial $p \in R$ with trivial constant term, the coefficients of the polynomial $\prod (t - \sigma p)$, where σ runs over the Weyl group W_G of G , are W_G -symmetric polynomials with trivial constant terms, and hence they belong to I . Since p is a root of the polynomial, we have $p^{|W_G|} \in I$, hence $p \in \sqrt{I}$.

Let $S = R_M$ be the localization of R at the maximal ideal M . Then S is a regular local ring of dimension n . Consider the sequence of polynomials in R :

$$(2.7) \quad w_0, w_1, w_2, \dots, w_{n-1}$$

with $w_0 := u$. Since u divides w_n , the ideal in R generated by these polynomials contains I and hence its radical is equal to $M = \sqrt{I}$. This means that the sequence (2.7) is a system of parameters in S . Since S is a regular local ring, by [8, Theorem 31], the sequence

(2.7) is regular. It follows from [8, Theorem 43], that the sequence (which is a part of the Koszul complex)

$$(2.8) \quad \Lambda^2(S^n) \xrightarrow{d'} S^n \xrightarrow{d} S,$$

where $d(e_i) = w_i$ and $d'(e_i \wedge e_j) = w_j e_i - w_i e_j$ (here e_0, e_1, \dots, e_{n-1} is the standard basis for S^n) is exact.

By (2.6), the n -tuple $(f_0, f_1, \dots, f_{n-1})$ is in the kernel of d . The exactness of (2.8) yields elements $g_i \in S$ such that

$$f_0 = w_1 g_1 + w_2 g_2 + \dots + w_{n-1} g_{n-1} \in IS.$$

It follows that $f = w f_n - f_0 \in wS + IS$. As $S = R_M$, there is a polynomial $h \in R$ with a nonzero constant term, satisfying $fh \in wR + I$. Since the polynomial f is homogeneous, it is also contained in $wR + I$. \square

REFERENCES

- [1] DEMAZURE, M. Invariants symétriques entiers des groupes de Weyl et torsion. *Invent. Math.* 21 (1973), 287–301.
- [2] ELMAN, R., KARPENKO, N., AND MERKURJEV, A. *The algebraic and geometric theory of quadratic forms*, vol. 56 of *American Mathematical Society Colloquium Publications*. American Mathematical Society, Providence, RI, 2008.
- [3] FULTON, W. *Intersection theory*, second ed., vol. 2 of *Ergebnisse der Mathematik und ihrer Grenzgebiete. 3. Folge. A Series of Modern Surveys in Mathematics [Results in Mathematics and Related Areas. 3rd Series. A Series of Modern Surveys in Mathematics]*. Springer-Verlag, Berlin, 1998.
- [4] KARPENKO, N. A. On generic flag varieties for odd spin groups. Preprint (4 May 2021, 11 pages). Available on authors' webpage.
- [5] KARPENKO, N. A. On generic quadratic forms. *Pacific J. Math.* 297, 2 (2018), 367–380.
- [6] KARPENKO, N. A. On generically split generic flag varieties. *Bull. Lond. Math. Soc.* 50 (2018), 496–508.
- [7] KARPENKO, N. A., AND MERKURJEV, A. S. Canonical p -dimension of algebraic groups. *Adv. Math.* 205, 2 (2006), 410–433.
- [8] MATSUMURA, H. *Commutative algebra*, second ed., vol. 56 of *Mathematics Lecture Note Series*. Benjamin/Cummings Publishing Co., Inc., Reading, Mass., 1980.
- [9] MERKURJEV, A. R -equivalence on three-dimensional tori and zero-cycles. *Algebra Number Theory* 2, 1 (2008), 69–89.
- [10] TOTARO, B. The Chow ring of a classifying space. In *Algebraic K-theory (Seattle, WA, 1997)*, vol. 67 of *Proc. Sympos. Pure Math.* Amer. Math. Soc., Providence, RI, 1999, pp. 249–281.
- [11] TOTARO, B. The torsion index of the spin groups. *Duke Math. J.* 129, 2 (2005), 249–290.
- [12] VISHIK, A. Fields of u -invariant $2^r + 1$. In *Algebra, arithmetic, and geometry: in honor of Yu. I. Manin. Vol. II*, vol. 270 of *Progr. Math.* Birkhäuser Boston Inc., Boston, MA, 2009, pp. 661–685.

(N. Karpenko) MATHEMATICAL & STATISTICAL SCIENCES, UNIVERSITY OF ALBERTA, EDMONTON, CANADA

Email address: karpenko@ualberta.ca

URL: www.ualberta.ca/~karpenko

(A. Merkurjev) DEPARTMENT OF MATHEMATICS, UNIVERSITY OF CALIFORNIA, LOS ANGELES, CA, USA

Email address: merkurev@math.ucla.edu

URL: www.math.ucla.edu/~merkurev