

# ON GENERICALLY SPLIT GENERIC FLAG VARIETIES

NIKITA A. KARPENKO

ABSTRACT. Let  $G$  be a split semisimple algebraic group over an arbitrary field  $F$ , let  $E$  be a  $G$ -torsor over  $F$ , and let  $P$  be a parabolic subgroup of  $G$ . The quotient variety  $X := E/P$ , known as a *flag variety*, is *generically split*, if the parabolic subgroup  $P$  is special. It is *generic*, provided that the  $G$ -torsor  $E$  over  $F$  is a standard generic  $G_k$ -torsor for a subfield  $k \subset F$  and a split semisimple algebraic group  $G_k$  over  $k$  with  $(G_k)_F = G$ .

For any generically split generic flag variety  $X$ , we show that the Chow ring  $\mathrm{CH} X$  is generated by Chern classes (of vector bundles over  $X$ ). This implies that the topological filtration on the Grothendieck ring of  $X$  coincides with the computable gamma filtration. The results were already known in some cases including the case where  $P$  is a Borel subgroup.

We also provide a complete classification of generically split generic flag varieties and, equivalently, of special parabolic subgroups for split simple groups.

## CONTENTS

0. Introduction	1
1. Special groups over algebraically closed fields	2
2. Split special reductive groups over arbitrary fields	3
3. Special parabolic subgroups of split semisimple groups	4
4. Chow rings of classifying spaces	4
5. Chern subrings for classifying spaces	5
6. Chern subrings for special parabolic subgroups	7
7. Chern subring for generic generically split flag varieties	8
8. Classification of special parabolic subgroups	10
References	14

## 0. INTRODUCTION

Let  $G$  be a split semisimple algebraic group over an arbitrary field  $k$  and let  $E$  be a standard generic  $G$ -torsor. By definition,  $E$  is the generic fiber of a standard versal  $G$ -torsor  $U \rightarrow U/G$ , whose total space  $U$  is a nonempty open  $G$ -equivariant subvariety in a finite-dimensional linear  $G$ -representation. The base of  $E$  is therefore the function field  $F := k(U/G)$ .

---

*Date:* 16 April 2017. Revised 10 April 2018.

*Key words and phrases.* Algebraic groups; projective homogeneous varieties; Chow groups. *Mathematical Subject Classification (2010):* 20G15; 14C25.

This work has been supported by a Discovery Grant from the National Science and Engineering Board of Canada.

Let  $B$  be a Borel subgroup of  $G$ . There is a conjecture (see [12, Conjecture 1.1]) that for the variety  $X := E/B$  – a generic variety of complete flags – the canonical epimorphism of the Chow ring  $\mathrm{CH} X$  onto the associated graded ring  $GK(X)$  of the topological filtration on the Grothendieck ring  $K(X)$  is an isomorphism. Since the ring  $K(X)$  is computed ([18]) and the topological filtration on it is known to coincide with the computable gamma filtration ([11, Example 2.4]), this conjecture, proved in several cases already, provides a way to compute the Chow ring.

There is an equivalent version of the above conjecture, where the Borel subgroup  $B$  is replaced by any fixed *special* parabolic subgroup  $P$ , see [13, Lemma 4.2]. By definition,  $P$  is special if any  $P$ -torsor over any extension field of the base field is trivial. This is equivalent to the fact that the generic flag variety  $X := E/P$  is *generically split*, see Lemma 7.1.

In the present paper we show that a generically split generic flag variety  $X = E/P$  has the same property as in the above particular case with  $P = B$ : the topological filtration on the Grothendieck ring  $K(X)$  coincides with the gamma filtration. This confirms that the introduced by A. Grothendieck gamma filtration is a good approximation of the topological filtration. And not only in the well-known sense that they both coincide after tensoring by  $\mathbb{Q}$ .

The reason for this coincidence (with  $\mathbb{Z}$ -coefficients) is our main result (Theorem 7.3) saying that the ring  $\mathrm{CH} X$  coincides with its Chern subring. In its turn, Theorem 7.3 is a consequence of the similar statement for the Chow ring of the classifying space of  $P$ , see Proposition 5.5.

The assumption that  $P$  is special is essential. For instance, taking for  $G$  a spinor group  $\mathrm{Spin}(n)$ , we may take a parabolic subgroup  $P \subset G$  such that the variety  $X = E/P$  is a projective quadric. Then the gamma filtration on  $K(X)$  is different from the topological filtration provided that  $n \geq 9$ . Indeed, as shown in [9], the component of codimension 2 of the graded ring associated with the gamma filtration contains an element of order 2; on the other hand, by [8, Theorem 6.1 and §3.1], the component of codimension 2 of the graded ring associated with the topological filtration is torsion-free for such  $n$ .

For an overview of sections of the paper we refer to the table of contents: the titles of sections are self-explaining. Note that the paper ends with a classification of special parabolic subgroups (or, equivalently, of generically split generic flag varieties).

ACKNOWLEDGEMENTS. I am grateful to Vladimir Chernousov and Alexander Merkurjev for useful comments. Many thanks to Alexander Merkurjev for his proofs of Lemmas 5.4 and 7.1.

## 1. SPECIAL GROUPS OVER ALGEBRAICALLY CLOSED FIELDS

An affine algebraic group  $G$  over a field  $F$  is *special*, if any  $G$ -torsor over any extension field of  $F$  is trivial. Since for any field extension  $L/F$ , the set of isomorphism classes of  $G$ -torsors over  $L$  is identified with the Galois cohomology set  $H^1(L, G)$ ,  $G$  is special if and only if  $H^1(L, G) = 1$  for any  $L$ .

Below in this section, we are assuming that the field  $F$  is algebraically closed.

**Theorem 1.1** ([1, Exposé 5, Théorème 3]). *Any special affine algebraic group over an algebraically closed field  $F$  is connected. A semisimple group  $G$  over  $F$  is special if and only if it is isomorphic to a finite direct product, where each factor is the special linear group  $\mathrm{SL}(n)$  (for some  $n \geq 2$ ) or the symplectic group  $\mathrm{Sp}(2n)$  (for some  $n \geq 2$ ).*

Classification of arbitrary special groups over  $F$  easily reduces to the semisimple case:

**Lemma 1.2** (see also [7, Theorem 1]). *Let  $G$  be an affine connected algebraic group over an algebraically closed field  $F$ . The following three conditions are equivalent:*

- (1)  $G$  is special;
- (2) the reductive group  $G/R_u$  is special, where  $R_u$  is the unipotent radical of  $G$ ;
- (3) the semisimple group  $(G/R_u)'$  is special, where  $(G/R_u)'$  is the commutator subgroup of  $G/R_u$ .

*Proof.* Let us prove equivalence of (1) and (2). By [2, Theorem 10.6], a connected unipotent group  $U$  over  $F$  has a finite filtration by closed normal subgroups with all successive quotients isomorphic to  $\mathbb{G}_a$ . Since for any field  $L/F$ ,  $H^i(L, \mathbb{G}_a) = 0$  for  $i > 0$ , we get that  $H^i(L, U) = 0$  for  $i > 0$  provided that  $U$  is abelian. The derived series of the unipotent radical  $R_u$  of  $G$  is a finite filtration of  $R_u$  by closed connected characteristic subgroups with abelian successive quotients. It follows that the map of the set  $H^1(L, G)$  of isomorphism classes of  $G$ -torsor over  $L$  to the set  $H^1(L, G/R_u)$ , induced by the projection  $G \rightarrow G/R_u$ , is a bijection.

Now we prove the equivalence of (2) and (3). We may assume that  $G$  is reductive, i.e.  $G = G/R_u$ . The quotient  $S := G/G'$  is then a (split) torus (called the *coradical* of  $G$ ). By [2, Proposition 11.20], the image in  $S$  of a maximal torus  $T$  of  $G$  is a maximal torus of  $S$  and therefore equals  $S$ . The intersection  $T' := T \cap G'$  is a (split) maximal torus of  $G'$ . For any field extension  $L/F$ , the exact sequence of tori

$$1 \rightarrow T' \rightarrow T \rightarrow S \rightarrow 1$$

produces an exact sequence

$$T(L) \rightarrow S(L) \rightarrow H^1(L, T').$$

Since  $H^1(L, T') = 1$ , the map  $T(L) \rightarrow S(L)$  is surjective. It follows that the map  $G(L) \rightarrow S(L)$  is also surjective. From the exact sequence

$$G(L) \rightarrow S(L) \rightarrow H^1(L, G') \rightarrow H^1(L, G) \rightarrow H^1(L, S)$$

with  $H^1(L, S) = 1$  we conclude that the map  $H^1(L, G') \rightarrow H^1(L, G)$  is a bijection.  $\square$

## 2. SPLIT SPECIAL REDUCTIVE GROUPS OVER ARBITRARY FIELDS

Let  $G$  be a split reductive group over an arbitrary field  $F$  and let  $G'$  be the split semisimple group given by the commutator subgroup of  $G$ . If  $G$  is special, then  $G$  remains special over any extension field of  $F$ . In particular, it remains special over an algebraic closure of  $F$ . By the results of the preceding section and classification of split semisimple groups, it follows that  $G'$  (over  $F$ ) is isomorphic to a direct product of special linear and symplectic groups. The converse also holds:

**Theorem 2.1** (see also [7, Proposition 15]). *For a split reductive group  $G$  over an arbitrary field the following statements are equivalent:*

- (1)  $G$  is special;
- (2)  $G'$  is special;
- (3)  $G'$  is isomorphic to a direct product of special linear and symplectic groups.

*Proof.* The equivalence of (1) and (2) can be shown precisely the same way as the equivalence of (2) and (3) in Lemma 1.2. It has been shown that (2) implies (3) in the discussion before Theorem 2.1. Since special linear groups, symplectic groups, and direct products of special groups are special, (3) implies (2).

Let us also note that for a split reductive  $G$ , the image of a split maximal torus in  $G$  under the projection of  $G$  onto the coradical  $G/G'$  of  $G$  is a split maximal torus. Hence the torus  $G/G'$  is split and, in particular, special. This way Theorem 2.1 becomes a particular case of [7, Proposition 15].  $\square$

### 3. SPECIAL PARABOLIC SUBGROUPS OF SPLIT SEMISIMPLE GROUPS

Let  $P$  be a parabolic subgroup of a split semisimple group  $G$  over an arbitrary field  $F$ . By [2, Proposition 20.5], the unipotent radical  $R_u$  of  $P$  is defined over  $F$ . Moreover,  $P$  contains a Levi subgroup  $L$  (over  $F$ );  $L$  is a split reductive group, the composition  $L \hookrightarrow P \rightarrow P/R_u$  is an isomorphism.

The same proof as in Lemma 1.2 (equivalence of (1) and (2)) shows that  $P$  is special if and only if  $L$  is special. We should mention here that an arbitrary connected unipotent group  $U$  over  $F$  is known to possess a filtration by closed normal subgroups with quotients isomorphic to  $\mathbb{G}_a$  only in the case of perfect  $F$ , [4, IV, §2, Corollaire 3.9]. This is fine for  $U = R_u$ , because  $P$  and therefore its unipotent radical  $R_u$  are defined over the prime subfield of  $F$ , which is perfect.

By Theorem 2.1 we know that  $L$  is special if and only if the split semisimple group  $L'$  is special which means that  $L'$  is isomorphic to a direct product of special linear and symplectic groups.

### 4. CHOW RINGS OF CLASSIFYING SPACES

Let  $G$  be a split connected reductive algebraic group over an arbitrary field  $F$ . Let  $T$  be a split maximal torus of  $G$  and let  $W := N_G(T)/T$  be the Weyl group of  $G$  with respect to  $T$ . We consider the  $G$ -equivariant Chow ring of the point  $\mathrm{CH}_G(\mathrm{Spec} F)$  also called the Chow ring of the classifying space of  $G$  and introduced in [27].

Assume that  $G$  is special. According to [5, Proposition 6], the homomorphism

$$\mathrm{CH}_G(\mathrm{Spec} F) \rightarrow \mathrm{CH}_T(\mathrm{Spec} F),$$

induced by the imbedding  $T \hookrightarrow G$  (see [5, §3.2]), identifies the ring  $\mathrm{CH}_G(\mathrm{Spec} F)$  with the subring  $\mathrm{CH}_T(\mathrm{Spec} F)^W$  of  $W$ -invariant elements in  $\mathrm{CH}_T(\mathrm{Spec} F)$ .

Viewing a character of  $T$  as a  $T$ -equivariant vector bundle over  $\mathrm{Spec} F$ , we can take its first equivariant Chern class in  $\mathrm{CH}_T^1(\mathrm{Spec} F)$ , [5, §3.2]. This assignment gives rise to a ring homomorphism

$$S(\hat{T}) \rightarrow \mathrm{CH}_T(\mathrm{Spec} F),$$

where  $S(\hat{T})$  is the symmetric ring of the character group  $\hat{T}$  of  $T$ . Since  $T$  is split, this homomorphism is a ring isomorphism so that we get an identification

$$\mathrm{CH}_G(\mathrm{Spec} F) = S(\hat{T})^W.$$

Summarizing we get

**Theorem 4.1** ([5, Proposition 6]). *Let  $G$  be a special reductive group over a field  $F$  containing a split maximal torus  $T$ . Then  $\mathrm{CH}_G(\mathrm{Spec} F) = S(\hat{T})^W$ .*

## 5. CHERN SUBRINGS FOR CLASSIFYING SPACES

Let  $G$  be an affine algebraic group over an arbitrary field  $F$ . The Chern subring of  $\mathrm{CH}_G(\mathrm{Spec} F)$  is the subring generated by the equivariant Chern classes of finite-dimensional linear  $G$ -representations over  $F$ . Note that the representation ring  $R(G)$  can be viewed as the  $G$ -equivariant Grothendieck ring of  $\mathrm{Spec} F$ , [17, Example 2]. The total equivariant Chern class  $c : R(G) \rightarrow \mathrm{CH}_G(\mathrm{Spec} F)$  is a homomorphism of the additive group  $R(G)$  into the multiplicative group of the elements in  $\mathrm{CH}_G(\mathrm{Spec} F)$  with the homogeneous component of codimension 0 equal to 1.

For any reductive group  $G$  containing a split maximal torus  $T$ , the restriction homomorphism  $R(G) \rightarrow R(T) = \mathbb{Z}[\hat{T}]$  identifies  $R(G)$  with the ring  $\mathbb{Z}[\hat{T}]^W$ , [22, Théorème 4]. Since the addition in  $\hat{T}$  becomes multiplication in  $\mathbb{Z}[\hat{T}]$ , we use the traditional exponential notation: we write  $e^\chi$  for a character  $\chi \in \hat{T}$  considered as an element of  $\mathbb{Z}[\hat{T}]$ . The above total Chern class  $c$  is a restriction of the total Chern class

$$c : \mathbb{Z}[\hat{T}] = R(T) \rightarrow \mathrm{CH}_T(\mathrm{Spec} F) = S(\hat{T}),$$

which is the homomorphism of the additive group of  $\mathbb{Z}[\hat{T}]$  to the multiplicative group of the elements in  $S(\hat{T})$  with the constant term 1, associating to  $e^\chi \in \mathbb{Z}[\hat{T}]$  the element  $1 + \chi \in S(\hat{T})$ . In particular, the first Chern class of  $e^\chi$  is  $\chi \in S(\hat{T})$ . Note that when we identify  $\hat{T}$  with a subset in  $S(\hat{T})$ , the operation on  $\hat{T}$  is the addition of the ring  $S(\hat{T})$ .

**Example 5.1.** For any integer  $n \geq 2$ , let us consider the special linear group  $G := \mathrm{SL}(n)$ . The group of characters  $\hat{T}$  of a split maximal torus  $T \subset G$  can be identified with the quotient of a free abelian group of rank  $n$  on generators  $x_1, \dots, x_n$  modulo the subgroup generated by  $x_1 + \dots + x_n$ . Moreover, the Weyl group  $W$  is the symmetric group on  $x_1, \dots, x_n$ . Identifying  $\mathrm{CH}_G(\mathrm{Spec} F)$  with  $S(\hat{T})^W$  and using [14, Lemma 8.1] to compute the  $W$ -invariants, we get that  $\mathrm{CH}_G(\mathrm{Spec} F)$  is the polynomial ring (over  $\mathbb{Z}$ ) in the elementary symmetric polynomials  $\sigma_2, \dots, \sigma_n$  in  $x_1, \dots, x_n$ . Since  $\sigma_i$  as an element of  $\mathrm{CH}_G(\mathrm{Spec} F)$  is the  $i$ th Chern class of the tautological representation of  $G$  (whose class in  $R(G) = \mathbb{Z}[\hat{T}]^W$  is equal to  $e^{x_1} + \dots + e^{x_n}$ ), we see that the ring  $\mathrm{CH}_G(\mathrm{Spec} F)$  coincides with its Chern subring.

**Example 5.2.** For any integer  $n \geq 1$ , let us consider the symplectic group  $G := \mathrm{Sp}(2n)$ . The group of characters  $\hat{T}$  of a split maximal torus  $T \subset G$  can be identified with a free abelian group of rank  $n$  on generators  $x_1, \dots, x_n$ . Moreover, the Weyl group  $W$  is a semidirect product of the symmetric group on  $x_1, \dots, x_n$  by a direct product of  $n$  copies of  $\mathbb{Z}/2\mathbb{Z}$ , where the  $i$ th copy of  $\mathbb{Z}/2\mathbb{Z}$  changes the sign of  $x_i$  and acts trivially on the

remaining generators. Identifying  $\mathrm{CH}_G(\mathrm{Spec} F)$  with  $S(\hat{T})^W$ , we see that  $\mathrm{CH}_G(\mathrm{Spec} F)$  is the polynomial ring  $\mathbb{Z}[\sigma_1, \dots, \sigma_n]$  in the elementary symmetric polynomials  $\sigma_1, \dots, \sigma_n$  in squares of  $x_1, \dots, x_n$ . Since  $\sigma_i$  is the  $2i$ th Chern class of the tautological representation of  $G$  (whose class in  $R(G) = \mathbb{Z}[\hat{T}]^W$  is equal to  $e^{x_1} + e^{x_1^{-1}} + \dots + e^{x_n} + e^{x_n^{-1}}$  so that the Chern classes are the  $2n$  elementary symmetric polynomials in  $\pm x_1, \dots, \pm x_n$ ), we see that the ring  $\mathrm{CH}_G(\mathrm{Spec} F)$  is generated by Chern classes. Note that the odd Chern classes of the tautological representation vanish.

**Example 5.3.** Let  $G = G_1 \times \dots \times G_n$  with  $n \geq 1$  and with each  $G_i$  isomorphic to a symplectic or a special linear group. Let  $T_i$  be a split maximal torus in  $G_i$ . Then  $T := T_1 \times \dots \times T_n$  is a split maximal torus in  $G$  and the Weyl group  $W$  of  $G$  relative to  $T$  is the product  $W_1 \times \dots \times W_n$  with  $W_i$  being the Weyl group of  $G_i$  relative to  $T_i$ . It follows that

$$\begin{aligned} \mathrm{CH}_G(\mathrm{Spec} F) &= S(\hat{T})^W = S(\hat{T}_1)^{W_1} \otimes \dots \otimes S(\hat{T}_n)^{W_n} = \\ &= \mathrm{CH}_{G_1}(\mathrm{Spec} F) \otimes \dots \otimes \mathrm{CH}_{G_n}(\mathrm{Spec} F). \end{aligned}$$

Since for any  $i$  the ring  $\mathrm{CH}_{G_i}(\mathrm{Spec} F)$  coincides with its Chern subring, the same holds for  $\mathrm{CH}_G(\mathrm{Spec} F)$ .

Let  $G$  be a split reductive group over  $F$  and let  $G'$  be its commutator subgroup. The following lemma shows that the homomorphism of the representation rings  $R(G) \rightarrow R(G')$ , given by restriction of representations, is surjective (cf. the beginning of proof of Proposition 5.5).

**Lemma 5.4** (A. S. Merkurjev, c.f. [16, Lemma 5.2]). *Let  $M$  be an abelian group endowed with an action of a finite (abstract) group  $W$  and let  $M'' \subset M^W$  be a torsion-free subgroup of  $W$ -invariant elements. For  $M' := M/M''$ , the homomorphism  $\mathbb{Z}[M]^W \rightarrow \mathbb{Z}[M']^W$  is surjective.*

*Proof.* For any subgroup  $W' \subset W$ , since the action of  $W'$  on  $M''$  is trivial, we have  $H^1(W', M'') = \mathrm{Hom}(W', M'')$ . Since  $W'$  is finite and the group  $M''$  is torsion-free, we have  $\mathrm{Hom}(W', M'') = 0$ . The exact cohomology sequence

$$M^{W'} \longrightarrow (M')^{W'} \longrightarrow H^1(W', M') = 0$$

therefore tells us that the homomorphism  $M^{W'} \rightarrow (M')^{W'}$  is surjective. For any  $m' \in M'$ , the sum  $s_{m'} \in \mathbb{Z}[M']$  of the exponents of the elements of the  $W'$ -orbit of  $m'$  in  $M'$  is  $W'$ -equivariant. The additive group of  $\mathbb{Z}[M']^W$  is generated by all such  $s_{m'}$ .

So, we only need to show that  $s_{m'}$  is in the image of  $\mathbb{Z}[M]^W$  for any given  $m' \in M'$ . For this, we consider the stabilizer  $W' \subset W$  of  $m'$ . By surjectivity of  $M^{W'} \rightarrow (M')^{W'}$ , we can find a  $W'$ -invariant element  $m \in M$  mapped to  $m'$ . The  $W'$ -orbit of  $e^m \in \mathbb{Z}[M]$  produces an element  $s_m \in \mathbb{Z}[M]^W$  mapped to  $s_{m'}$ .  $\square$

**Proposition 5.5.** *For any split special reductive group  $G$  over an arbitrary field, the ring  $\mathrm{CH}_G(\mathrm{Spec} F)$  coincides with its Chern subring.*

*Proof.* Let  $T$  be a split maximal torus in  $G$  and let  $T'$  be the split maximal torus  $T \cap G'$  of  $G'$ . The Weyl group  $W$  of  $G$  relative to  $T$  is identified with the Weyl group of  $G'$  relative to  $T'$ , and this identification respects the action on  $T'$ . Moreover, the action of  $W$  on

$T/T'$  is trivial. Therefore, setting  $M := \hat{T}$  and  $M'' := \widehat{T/T'}$ , we reduce the proof to the following elementary statement (see Lemma 5.6 below).

Let  $M$  be a lattice endowed with an action of a finite group  $W$ . We define the (abstract) total Chern class map  $c : \mathbb{Z}[M] \rightarrow S(M)$  as the homomorphism of the additive group  $\mathbb{Z}[M]$  into the multiplicative group of the elements in  $S(M)$  with the homogeneous component of degree 0 equal to 1, assigning to  $e^m \in \mathbb{Z}[M]$  for each  $m \in M$  the element  $1+m \in S(M)$ . Note that  $c(\mathbb{Z}[M]^W) \subset S(M)^W$ . We also write  $c$  for the map  $\mathbb{Z}[M]^W \rightarrow S(M)^W$  thus obtained, and also call it the (abstract) total Chern class map. We say that the ring  $S(M)^W$  is generated by Chern classes, if it is generated by the homogeneous component of the elements from the image of  $c$ .

**Lemma 5.6.** *Let  $M$  be a lattice endowed with an action of a finite group  $W$  and let  $M'' \subset M^W$  be a subgroup of  $W$ -invariant elements such that the quotient  $M' := M/M''$  is free. Assume that the ring  $S(M')^W$  is generated by Chern classes. Then the ring  $S(M)^W$  is also generated by Chern classes.*

*Proof.* Choosing a basis of  $M''$ , taking the corresponding flag of subgroups in  $M''$ , and decomposing the projection  $M \rightarrow M'$  in a finite chain of epimorphisms with cyclic kernels, we reduce the proof to the case of cyclic  $M''$ .

In the commutative diagram

$$\begin{array}{ccc} R(G) = \mathbb{Z}[M]^W & \longrightarrow & R(G') = \mathbb{Z}[M']^W \\ \downarrow c & & \downarrow c \\ \mathrm{CH}_G(\mathrm{Spec} F) = S(M)^W & \longrightarrow & \mathrm{CH}_{G'}(\mathrm{Spec} F) = S(M')^W \end{array}$$

the upper map is an epimorphism by Lemma 5.4. We fix a generator  $y$  of  $M''$ . We show that a homogeneous element  $f \in S(M)^W$  is a polynomial in Chern classes using induction on  $\deg f$ . For  $f$  of degree 0 there is nothing to prove. Assume that the degree of  $f$  is positive and write its image  $f' \in S(M')^W$  as a polynomial in Chern classes of some elements in  $\mathbb{Z}[M']^W$ . Lifting these elements to  $\mathbb{Z}[M]^W$  in an arbitrary way and taking the same polynomial in the Chern classes of the liftings, we get an element  $g \in S(M)^W$  such that the difference  $f - g$  vanishes in  $S(M')^W$ . Since the kernel of the ring homomorphism  $S(M) \rightarrow S(M')$  is the ideal in  $S(M)$  generated by  $y$ , we can write  $f - g = yh$  with some homogeneous  $h \in S(M)$  of degree  $\deg h = \deg f - 1$ . Since  $y$  and  $yh$  are  $W$ -invariant while the ring  $S(W)$  is a polynomial ring, it follows that  $h \in S(M)^W$ . So, by induction hypothesis,  $h$  can be written as a polynomial in Chern classes.  $\square$

$\square$

## 6. CHERN SUBRINGS FOR SPECIAL PARABOLIC SUBGROUPS

**Proposition 6.1.** *Let  $P$  be a special parabolic subgroup of a split semisimple algebraic group over an arbitrary field  $F$ . Then the ring  $\mathrm{CH}_P(\mathrm{Spec} F)$  coincides with its Chern subring.*

*Proof.* Let  $L$  be a Levi subgroup of  $P$ . Then  $L$  is a split special reductive group so that the ring  $\mathrm{CH}_L(\mathrm{Spec} F)$  coincides with its Chern subring. Since the composition

$R(L) \rightarrow R(P) \rightarrow R(L)$  is the identity, the homomorphism  $R(P) \rightarrow R(L)$  is surjective. Therefore it suffices to show that the homomorphism  $\mathrm{CH}_P(\mathrm{Spec} F) \rightarrow \mathrm{CH}_L(\mathrm{Spec} F)$  is an isomorphism.

In order to show that  $\mathrm{CH}_P^i(\mathrm{Spec} F) \rightarrow \mathrm{CH}_L^i(\mathrm{Spec} F)$  is an isomorphism for any  $i \geq 0$ , we recall that  $\mathrm{CH}_P^i(\mathrm{Spec} F) = \mathrm{CH}^i(U/P)$  for an open subset  $U \subset V$  of a finite-dimensional linear  $P$ -representation  $V$  admitting a  $P$ -torsor  $U \rightarrow U/P$  and such that  $\mathrm{codim}_V(V \setminus U) > i$ . For any such  $U$  we also have  $\mathrm{CH}_L^i(\mathrm{Spec} F) = \mathrm{CH}^i(U/L)$  and the homomorphism  $\mathrm{CH}^i(U/P) \rightarrow \mathrm{CH}^i(U/L)$  we are talking about is the pull-back with respect to the projection  $U/L \rightarrow U/P$ .

Let  $R_u$  be the unipotent radical of  $P$ . Then  $P$  is the semidirect product  $R_u \rtimes L$ . In particular, the quotient variety  $P/L$  is isomorphic to the underlying variety of  $R_u$ . Since the algebraic group  $R_u$  is connected, unipotent, and defined over the prime subfield of  $F$ , the variety  $R_u$  is isomorphic to an affine space, [4, IV, §2, Corollaire 3.9 and IV, §4, Théorème (de Lazard) 4.1].

Since  $P$  is special, for any point  $x \in U/P$  the fiber of the morphism  $U \rightarrow U/P$  over  $x$  is isomorphic to  $P_{F(x)}$ . Therefore the fiber over  $x$  of the morphism  $U/L \rightarrow U/P$  is isomorphic to the quotient variety  $(P/L)_{F(x)}$ . As explained above,  $P/L$  is an affine space so that the pull-back of Chow groups  $\mathrm{CH}(U/P) \rightarrow \mathrm{CH}(U/L)$  is an isomorphism by the homotopy invariance of Chow groups, [6, Theorem 57.13].

In fact, proving that  $\mathrm{CH}_P(\mathrm{Spec} F) \rightarrow \mathrm{CH}_L(\mathrm{Spec} F)$  is an isomorphism, we don't really need the assumption that the parabolic subgroup  $P$  is special. Indeed, in general, every fiber of  $U/L \rightarrow U/P$  is the twist of  $R_u$  by a  $P$ -torsor. It follows by [24, Theorem (iii) of §4.4.1 in Chapter 2], that all fibers are isomorphic to affine spaces.  $\square$

## 7. CHERN SUBRING FOR GENERIC GENERICALLY SPLIT FLAG VARIETIES

Let  $F$  be a field,  $G$  a split semisimple algebraic group over  $F$ ,  $P$  a parabolic subgroup of  $G$ , and  $E$  a  $G$ -torsor over  $F$ . The flag variety  $X := E/P$  is *generically split* (cf. [19] and [20]) if the  $G$ -torsor  $E$  becomes trivial over the function field  $F(X)$ .

The flag variety  $X$  is *generic*, provided that the  $G$ -torsor  $E$  over  $F$  is a standard generic  $G_k$ -torsor for a subfield  $k \subset F$  and a split semisimple algebraic group  $G_k$  over  $k$  with  $(G_k)_F = G$ .

The following lemma can be used to detect the special parabolic subgroups (cf. §8):

**Lemma 7.1** (cf. [13, Lemma 4.1]). *A generic flag variety  $X = E/P$  is generically split if and only if the parabolic subgroup  $P$  is special.*

*Proof.* If  $P$  is special, the variety  $E'/P$  is generically split for any  $G$ -torsor  $E'$  over  $F$  by [14, Lemma 6.5]. In particular, this holds for  $E' = E$ .

Conversely, if a generic flag variety  $X = E/P$  is generically split, then for any field  $L/F$  and any  $G$ -torsor  $E'$  over  $L$ , the flag variety  $E'/P$  is also generically split. It follows by [21, §5] that  $P$  is special.

Below is a direct proof of both implications due to A. Merkurjev.

Assume that  $P$  is special. The pull-back of a  $G$ -torsor  $E$  over the field  $F$  to the field  $F(E/P)$  coincides with the push-forward of the generic fiber of the  $P$ -torsor  $E \rightarrow E/P$  and hence is trivial since  $P$  is special.

Conversely, assume that a generic flag variety  $E/P$  is generically split. We may assume that  $G$  is defined over a subfield  $k \subset F$  such that the  $G$ -torsor  $E$  over  $F$  is the generic fiber of the  $G$ -torsor  $U \rightarrow U/G$  for a nonempty open subset  $U$  of a finite-dimensional linear  $G$ -representation over  $k$ . In particular,  $F$  is the function field  $k(U/G)$ . By assumption, the pull-back  $E'$  of  $E$  to the field  $F(E/P)$  is trivial. On the other hand,  $E'$  is the push-forward of the generic fiber  $E''$  of the  $P$ -torsor  $U \rightarrow U/P$ . Since for any field  $L/F$  the map  $G(L) \rightarrow (G/P)(L)$  is surjective by [2, Proposition 20.5], the map  $H^1(L, P) \rightarrow H^1(L, G)$  has trivial kernel. It follows that  $E''$  is trivial, i.e., the morphism  $U \rightarrow U/P$  has a section over a nonempty open subset  $W \subset U/P$ . Since the  $P$ -torsor  $U \rightarrow U/P$  is versal, every  $P$ -torsor over an infinite field  $K/k$  is isomorphic to the fiber of  $U \rightarrow U/P$  over a point in  $W(K)$  ([23, §5 of Chapter I]) and hence is trivial. Therefore,  $P$  is special.  $\square$

Here is an extended version of Lemma 7.1:

**Corollary 7.2.** *Let  $G$  be a split semisimple algebraic group over an arbitrary field  $k$ , let  $P$  be a parabolic subgroup of  $G$ , and let  $\tilde{E}$  be a standard generic  $G$ -torsor. The following statements are equivalent:*

- (1) *The parabolic subgroup  $P$  is special.*
- (2) *The generic flag variety  $\tilde{E}/P$  is generically split.*
- (3) *For any field  $F/k$  and any  $G$ -torsor  $E$  over  $F$ , the flag variety  $E/P$  is generically split.*
- (4) *For any field  $F/k$  and any non-trivial  $G$ -torsor  $E$  over  $F$ , the flag variety  $E/P$  has no rational points.*

*Proof.* Equivalence of (1), (2) and (3) is already contained in the statement or in the proof of Lemma 7.1.

If  $E/P$  is generically split and has a rational point, then the  $G$ -torsor  $E$  becomes trivial over the function field  $F(E/P)$  and this function field is purely transcendental. It follows that  $E$  is trivial already over  $F$ . Therefore (3) implies (4).

Conversely, let us assume (4). Since over  $F(E/P)$  the variety  $E/P$  gets a rational point, the  $G$ -torsor  $E$  becomes trivial over  $F(E/P)$ . This means that the flag variety  $E/P$  is generically split.  $\square$

Here is the main result of the paper:

**Theorem 7.3.** *For any generically split generic flag variety  $X$  over a field  $F$ , the Chow ring  $\mathrm{CH} X$  coincides with its Chern subring.*

*Proof.* The Chow ring  $\mathrm{CH} X = \mathrm{CH}(E/P)$  can be viewed as the  $P$ -equivariant Chow ring  $\mathrm{CH}_P(E)$ , [5]. The characteristic map  $\mathrm{CH}_P(\mathrm{Spec} F) \rightarrow \mathrm{CH}_P(E)$  is then the pull-back with respect to the structure morphism  $E \rightarrow \mathrm{Spec} F$ . It follows that the characteristic map maps the Chern subring of  $\mathrm{CH}_P(\mathrm{Spec} F)$  into the Chern subring of  $\mathrm{CH} X$ . Since the characteristic map is surjective (see, e.g., [11, §2]) and the ring  $\mathrm{CH}_P(\mathrm{Spec} F)$  coincides with its Chern subring (Proposition 6.1), the statement follows.  $\square$

**Corollary 7.4.** *For any generic generically split flag variety  $X$ , the topological filtration on the Grothendieck ring  $K(X)$  coincides with the gamma filtration.*

*Proof.* For any regular algebraic variety  $X$  over a field with the property that the ring  $\mathrm{CH} X$  coincides with its Chern subring, the topological filtration on  $K(X)$  coincides with the gamma filtration, cf. [10, Proof of Theorem 3.7]. More precisely, gamma filtration coincides with the topological one if and only if the graded ring  $GK(X)$  associated with the topological filtration coincides with its Chern subring.  $\square$

## 8. CLASSIFICATION OF SPECIAL PARABOLIC SUBGROUPS

In this section we classify special parabolic subgroups of split simple algebraic groups. In view of Lemma 7.1, this is equivalent to a classification of generically split generic flag varieties.

As is already clear from the description in §3, the classification does not depend on the base field.

Since parabolic subgroups of products of semisimple groups are products of parabolic subgroups of the factors and are special if and only if each factor is special, we also get a classification of special parabolic subgroups for split semisimple groups which are products of simple ones.

Let  $G$  be a split semisimple algebraic group over an arbitrary field  $F$ . To a parabolic subgroup  $P$  of  $G$ , a subset of the set of vertices of the Dynkin diagram of  $G$  is associated, [25, §1.6]. For instance, the empty subset is associated to a Borel subgroup while the set of all vertices is associated to  $P = G$ . The subset associated to  $P$  is called the *type of  $P$* . Most of the time, the classification is easier to formulate in terms of the complement of  $P$  which we call the *co-type of  $P$* .

Recall that the Dynkin diagram of the semisimple part of (a Levi subgroup of)  $P$  is obtained by removing from the Dynkin diagram of  $G$  the vertices of the co-type of  $P$  (as well as the edges attached to at least one vertex removed). By §3, a necessary condition for  $P$  to be special is that the Dynkin diagram of its semisimple part is a disjoint union of diagrams of types A and/or C. As explained below, this necessary condition turns out to be also sufficient for simply connected  $G$ .

Corollary 7.2 relates detection of special parabolic subgroups in  $G$  with determination of possible Tits indexes of the inner twisted forms of  $G$ . Indeed, if  $E$  is a  $G$ -torsor over  $F$ , the flag variety  $E/P$  is a projective homogeneous variety under the twisted form  $G^E$  of  $G$  given by  $E$ . Note that the group  $G^E$  is split (i.e.,  $G^E \simeq G$ ) if and only if the torsor  $E$  is trivial: if  $G^E$  is split, the variety  $E/B$  for a Borel subgroup  $B \subset G$  has a rational point; therefore  $E$  is trivial since  $B$  is special. Also note that for adjoint  $G$ , the groups  $G^E$  are precisely all (up to an isomorphism) inner forms of  $G$ . And for simply connected  $G$ , the groups  $G^E$  are precisely all strongly inner forms of  $G$ .

Recall that the Tits index of  $G^E$  is, like the type of  $P$ , also a subset of vertices of the Dynkin diagram of  $G$ . In terms of the Tits indexes, the equivalence (1)  $\Leftrightarrow$  (4) in Corollary 7.2 translates as follows:  $P$  is special if and only if every Tits index that occurs for  $G^E$  when  $E$  varies over all  $G$ -torsors over all field extensions of  $F$ , coincides with the total Dynkin diagram of  $G$  provided that it contains the co-type of  $P$ .

From now on  $G$  is additionally supposed to be simple. The part of the classification of Tits indexes given in [25], which deals with the groups of inner type, provides the list of all Tits indexes of  $G^E$  in the case of adjoint  $G$ . The classification given below in the

adjoint case is therefore justified by this information. For non-adjoint  $G$ , however, the list of possible Tits indexes of  $G^E$  is a subset of the adjoint list, enlarging the list of co-types of special parabolic subgroups and requiring additional arguments.

For simply connected  $G$  such an additional argument is [26, Corollary 1] (see also [26, Theorem 2(i)], thanks to the referee for indicating these references). It shows that (as already claimed)  $P$  is special if and only if removing the co-type of  $P$  from the Dynkin diagram of  $G$  we get a disjoint union of diagrams of types A and/or C. (We recall that the only if part has been already shown several lines above.)

It remains to discuss only every such  $G$  which is neither adjoint nor simply connected. Then  $G$  is of classical type and for  $G$  of classical types, the most natural argument, at least for the author, relies on realizations of the group  $G$  and the varieties  $E/P$ . This argument is provided below after the complete classification list is stated.

We use the numbering of the vertices of the Dynkin diagram of  $G$  as in [3]. (For classical types, this numbering is also described in [15, §24.A].) This way the (co-)type of  $P$  becomes a set of integers. We consider all split simple algebraic groups  $G$  – type by type – and in each case we provide a necessary and sufficient condition for a parabolic subgroup  $P \subset G$  to be special (in terms of the (co-)type of  $P$ ):

- $G$  is of type  $A_n$  ( $n \geq 1$ ),  $G \simeq \mathrm{SL}(n+1)/\mu_m$ : *the greatest common divisor of the co-type of  $P$  and  $m$  is 1.*
- $G$  is of type  $B_n$  ( $n \geq 1$ ).
  - $G$  is adjoint:  *$n$  does not belong to the type of  $P$ .*
  - $G$  is simply connected: *at least one of the numbers  $n, n-1, n-2$  does not belong to the type of  $P$ .* (For  $n \geq 3$ , the condition means that at least one of  $n, n-1, n-2$  belongs to the co-type.) In particular, any parabolic subgroup (including  $G$ ) is special if  $n \leq 2$ ; for  $n = 3$  any parabolic subgroup different from  $G$  is special.
- $G$  is of type  $C_n$  ( $n \geq 1$ ).
  - $G$  is adjoint: *the co-type of  $P$  contains at least one odd number.*
  - $G$  is simply connected: *no condition.* In other words, all parabolic subgroups of  $G$  (including  $G$  itself) are special.
- $G$  is of type  $D_n$  ( $n \geq 3$ ).
  - $G$  is adjoint: *the co-type of  $P$  satisfies at least one of the following three conditions: (1) contains at least one of the numbers  $n, n-1$  and at least one odd number  $\leq n-2$ ; (2) contains both of the numbers  $n, n-1$ ; (3) for odd  $n$ , contains at least one of the numbers  $n, n-1$ .*
  - $G = O^+(2n)$  (special orthogonal group): *the co-type of  $P$  contains at least one of the numbers  $n$  and  $n-1$ .*
  - $G = \mathrm{Spin}^\sim(2n)$  (semispinor group,  $n$  is even): *the co-type of  $P$  satisfies at least one of the following two conditions: (1) contains at least one of the numbers  $n, n-1, n-2, n-3$  and at least one odd number  $\leq n-3$ ; (2) contains both of the numbers  $n, n-1$ .*
  - $G$  is simply connected: *at least one of the numbers  $n, n-1, n-2, n-3$  is not in the type of  $P$ .* (For  $n \geq 4$ , the condition means that at least one of  $n, n-1, n-2, n-3$  is in the co-type of  $P$ .) In particular, for  $n = 3$  all

parabolic subgroups are special (including  $G = \text{Spin}(6) = \text{SL}(4)$ ); for  $n = 4$ , all parabolic subgroups different from  $G$  are special.

- $G$  is of type  $G_2$ : *the co-type of  $P$  is non-empty*. In other words, any parabolic subgroup  $P \subsetneq G$  is special (while  $G$  itself is not).
- $G$  is of type  $F_4$ : *the co-type of  $P$  is not contained in  $\{4\}$* .
- $G$  is of type  $E_6$ .
  - $G$  is adjoint: *the co-type of  $P$  is contained neither in  $\{1, 6\}$  nor in  $\{2, 4\}$* .
  - $G$  is simply connected: *the co-type of  $P$  is not contained in  $\{1, 6\}$* .
- $G$  is of type  $E_7$ .
  - $G$  is adjoint: *the co-type of  $P$  is contained neither in  $\{1, 6, 7\}$  nor in  $\{1, 3, 4, 6\}$* .
  - $G$  is simply connected: *the co-type of  $P$  is not contained in  $\{1, 6, 7\}$* .
- $G$  is of type  $E_8$ : *the co-type of  $P$  is not contained in  $\{1, 6, 7, 8\}$* .

To finish, here are the proofs for all  $G$  which are neither adjoint nor simply connected.

**Type A.** Let  $G$  be of type  $A_n$  ( $n \geq 1$ ). Then  $G$  is isomorphic to  $\text{SL}(n+1)/\mu_m$ , where  $m \geq 1$  is a divisor of  $n+1$ . (The simply connected and adjoint cases correspond to  $m = 1$  and  $m = n+1$  respectively, we don't need to exclude them from the proof.) Let  $r_1 < \dots < r_k$  be the co-type of  $P$ . We are going to prove that

- *$P$  is special if and only if the greatest common divisor of the integers  $r_1, \dots, r_k$  and  $m$  is 1.*

In particular, for  $m = 1$  all parabolic subgroups (including  $G = \text{SL}(n+1)$  itself) are special.

Note that a  $G$ -torsor  $E$  over  $F$  gives rise to a central simple  $F$ -algebra  $A$  of degree  $n+1$  and of exponent dividing  $m$ . The isomorphism class of any central simple  $F$ -algebra  $A$  of degree  $n+1$  and of exponent dividing  $m$  can be obtained this way. Moreover, the algebra given by  $E$  is split if and only if  $E$  is trivial. The variety  $E/P$  can be identified with the variety of flags of right ideals in  $A$  of reduced dimensions  $r_1, \dots, r_k$ .

To prove the above classification of special parabolic subgroups in  $G$ , we use Corollary 7.2. If the numbers  $r_1, \dots, r_k, m$  have a common divisor  $d > 1$ , we find – over an appropriate field  $L/F$  – a central simple algebra  $A$  of degree  $n+1$  and of index and exponent  $d$ . Let  $E$  be a  $G$ -torsor over  $L$  producing the algebra  $A$ . Since  $\text{ind } A = d \neq 1$ ,  $E$  is nontrivial. However the variety  $E/P$  of flags of right ideals in  $A$  of reduced dimensions  $r_1, \dots, r_k$  has a rational point. Consequently,  $P$  is not special.

On the other hand, if the greatest common divisor of  $r_1, \dots, r_k, m$  is 1, then for any field  $L/F$  and any  $G$ -torsor  $E$  over  $L$ , the index of the central simple  $L$ -algebra  $A$  associated to  $E$  is 1 provided that the flag variety  $E/P$  (over  $L$ ) possesses a rational point. This means that  $E$  is trivial. Therefore  $P$  is special.

**Type D.** We start with the case  $G = O^+(2n)$ ,  $n \geq 3$ . Recall the characterization we are going to prove:

- *A parabolic subgroup  $P$  of  $G$  is special if and only if its co-type contains at least one of the numbers  $n$  and  $n-1$ .*

Isomorphism classes of  $G$ -torsors over  $F$  are in natural bijection with the isomorphism classes of the  $2n$ -dimensional non-degenerate quadratic forms over  $F$  of trivial discriminant. Under this bijection, the trivial  $G$ -torsor corresponds to the hyperbolic quadratic

form. If a  $G$ -torsor  $E$  corresponds to a quadratic form  $q$  and the co-type of  $P$  consists of the numbers  $r_1 < \cdots < r_k$ , the flag variety  $E/P$  is interpreted as follows. If  $r_k \leq n - 2$ ,  $E/P$  is the variety of flags of totally isotropic subspaces in  $q$  of dimensions  $r_1, \dots, r_k$ . If  $r_k = n$  and  $r_{k-1} = n - 1$ , then  $E/P$  is the variety of flags of totally isotropic subspaces in  $q$  of dimensions  $r_1, \dots, r_{k-1}$ . Finally, if  $r_{k-1} \leq n - 2$  and  $r_k \geq n - 1$ ,  $E/P$  is one of two (isomorphic) connected components of the variety of flags of totally isotropic subspaces in  $q$  of dimensions  $r_1, \dots, r_{k-1}, n$ .

Now we prove the above classification. If none of  $n$  and  $n - 1$  is in the co-type of  $P$  (meaning that  $r_k \leq n - 2$ ), we show that  $P$  is not special, considering a field  $L/F$  admitting a  $2n$ -dimensional quadratic form  $q$  of trivial discriminant and of Witt index  $n - 2$ . For the corresponding  $E$ , the variety  $E/P$  has then a rational point, but  $E$  is not trivial.

If at least one of  $n, n - 1$  is in the co-type of  $P$  and  $E/P$  has a rational point for some  $E$  over some  $L/F$ , the Witt index of the quadratic form  $q$  corresponding to  $E$  is at least  $n - 1$ . Since the discriminant of  $q$  is trivial, it follows that the Witt index is  $n$ , i.e.,  $q$  is hyperbolic. Therefore  $E$  is trivial and we proved  $P$  is special.

Our second and last case is that of a semispinor group  $G = \text{Spin}^\sim(2n)$  (here  $n$  is at least 4 and even). Let us recall the statement under proof:

- *A parabolic subgroup  $P$  of  $G$  is special if and only if its co-type satisfies at least one of the following two conditions:*
  - (1) *contains at least one of the numbers  $n, n - 1, n - 2, n - 3$  and at least one odd number  $\leq n - 3$ ;*
  - (2) *contains both of the numbers  $n, n - 1$ .*

We have a canonical surjection with trivial kernel of the set of isomorphism classes of  $G$ -torsors over  $F$  onto the set of isomorphism classes of central simple  $F$ -algebras of degree  $2n$  endowed with a quadratic pair with trivial discriminant and a split component of the Clifford algebra, [15, §29.F]. (Note that the index of any central simple algebra, carrying a quadratic pair, is a power of 2.) If a  $G$ -torsor  $E$  gives rise to an algebra  $A$  and the co-type of a parabolic subgroup  $P \subset G$  consists of the numbers  $r_1 < \cdots < r_k$ , the variety  $E/P$  is interpreted as the flag variety of isotropic right ideals in  $A$  of the reduced dimensions described as follows. If  $r_k \leq n - 2$ , the reduced dimensions are  $r_1, \dots, r_k$ . If  $r_k = n$  and  $r_{k-1} = n - 1$ , the reduced dimensions are  $r_1, \dots, r_{k-1}$ . Finally, if  $r_{k-1} \leq n - 2$  and  $r_k \geq n - 1$ , the variety  $E/P$  is one of two (not necessarily isomorphic) connected components of the variety of flags of isotropic ideals in  $A$  of reduced dimensions  $r_1, \dots, r_{k-1}, n$ .

Assume that  $P$  satisfies the condition of the classification. Then at least one of the reduced dimensions in the description of the variety  $E/P$  is odd. Therefore, the central simple algebra  $A$  is split as soon as the variety  $E/P$  has a rational point. In this case the quadratic pair on  $A$  is adjoint to a  $2n$ -dimensional non-degenerate quadratic form of trivial discriminant and Clifford invariant and of Witt index  $\geq n - 3$ . Such a quadratic form is necessarily hyperbolic, implying triviality of  $E$ .

We proved that the condition of the classification is sufficient (for  $P$  to be special). To prove that it is necessary, we first note – by looking at the simply connected case – that the part “the co-type of  $P$  contains at least one of  $n, n - 1, n - 2, n - 3$ ” is indeed

necessary. So, assume that  $P$  satisfies this part of the condition but does not satisfy the entire condition meaning that the co-type does not contain any odd number  $\leq n - 3$  and does not contain at least one of  $n, n - 1$ . In such a situation, we show that  $P$  is not special, considering a field  $L/F$  admitting a central simple  $L$ -algebra  $A$  of degree  $2n$  and index 2. We endow such  $A$  with a hyperbolic quadratic pair. By [15, Proposition 8.31], one of two components of the corresponding Clifford algebra is split. Therefore  $A$  is given by some  $G$ -torsor  $E$  over  $L$ . For such  $E$ , the variety  $E/P$  has a rational point. However the torsor  $E$  is not trivial, showing that  $P$  is not special.

### REFERENCES

- [1] *Séminaire C. Chevalley; 2e année: 1958. Anneaux de Chow et applications*. Secrétariat mathématique, 11 rue Pierre Curie, Paris, 1958.
- [2] BOREL, A. *Linear algebraic groups*, second ed., vol. 126 of *Graduate Texts in Mathematics*. Springer-Verlag, New York, 1991.
- [3] BOURBAKI, N. *Lie groups and Lie algebras. Chapters 4–6*. Elements of Mathematics (Berlin). Springer-Verlag, Berlin, 2002. Translated from the 1968 French original by Andrew Pressley.
- [4] DEMAZURE, M., AND GABRIEL, P. *Groupes algébriques. Tome I: Géométrie algébrique, généralités, groupes commutatifs*. Masson & Cie, Éditeur, Paris; North-Holland Publishing Co., Amsterdam, 1970. Avec un appendice *Corps de classes local* par Michiel Hazewinkel.
- [5] EDIDIN, D., AND GRAHAM, W. Equivariant intersection theory. *Invent. Math.* 131, 3 (1998), 595–634.
- [6] ELMAN, R., KARPENKO, N., AND MERKURJEV, A. *The algebraic and geometric theory of quadratic forms*, vol. 56 of *American Mathematical Society Colloquium Publications*. American Mathematical Society, Providence, RI, 2008.
- [7] HURUGUEN, M. Special reductive groups over an arbitrary field. *Transform. Groups* 21, 4 (2016), 1079–1104.
- [8] KARPENKO, N. A. Algebro-geometric invariants of quadratic forms. *Algebra i Analiz* 2, 1 (1990), 141–162.
- [9] KARPENKO, N. A. The Grothendieck ring of quadrics, and gamma filtration. In *Rings and modules. Limit theorems of probability theory, No. 3 (Russian)*. Izd. St.-Peterbg. Univ., St. Petersburg, 1993, pp. 39–61, 256.
- [10] KARPENKO, N. A. Codimension 2 cycles on Severi-Brauer varieties. *K-Theory* 13, 4 (1998), 305–330.
- [11] KARPENKO, N. A. Chow groups of some generically twisted flag varieties. *Annals of K-Theory* 2, 2 (2017).
- [12] KARPENKO, N. A. Chow ring of generic flag varieties. *Math. Nachr.* 290, 16 (2017), 2641–2647.
- [13] KARPENKO, N. A. Chow ring of generically twisted varieties of complete flags. *Adv. Math.* 306 (2017), 789–806.
- [14] KARPENKO, N. A., AND MERKURJEV, A. S. Canonical  $p$ -dimension of algebraic groups. *Adv. Math.* 205, 2 (2006), 410–433.
- [15] KNUS, M.-A., MERKURJEV, A., ROST, M., AND TIGNOL, J.-P. *The book of involutions*, vol. 44 of *American Mathematical Society Colloquium Publications*. American Mathematical Society, Providence, RI, 1998. With a preface in French by J. Tits.
- [16] MERKURJEV, A. Unramified degree three invariants of reductive groups. *Adv. Math.* 293 (2016), 697–719.
- [17] MERKURJEV, A. S. Equivariant  $K$ -theory. In *Handbook of K-theory. Vol. 1, 2*. Springer, Berlin, 2005, pp. 925–954.
- [18] PANIN, I. A. On the algebraic  $K$ -theory of twisted flag varieties. *K-Theory* 8, 6 (1994), 541–585.
- [19] PETROV, V., AND SEMENOV, N. Generically split projective homogeneous varieties. *Duke Math. J.* 152, 1 (2010), 155–173.

- [20] PETROV, V., AND SEMENOV, N. Generically split projective homogeneous varieties. II. *J. K-Theory* 10, 1 (2012), 1–8.
- [21] PETROV, V., AND SEMENOV, N. Rost motives, affine varieties, and classifying spaces. *J. Lond. Math. Soc. (2)* 95, 3 (2017), 895–918.
- [22] SERRE, J.-P. Groupes de Grothendieck des schémas en groupes réductifs déployés. *Inst. Hautes Études Sci. Publ. Math.*, 34 (1968), 37–52.
- [23] SERRE, J.-P. Cohomological invariants, Witt invariants, and trace forms. In *Cohomological invariants in Galois cohomology*, vol. 28 of *Univ. Lecture Ser.* Amer. Math. Soc., Providence, RI, 2003, pp. 1–100. Notes by Skip Garibaldi.
- [24] SPRINGER, T. A. Linear algebraic groups. In *Algebraic geometry, 4 (Russian)*, Itogi Nauki i Tekhniki. Akad. Nauk SSSR, Vsesoyuz. Inst. Nauchn. i Tekhn. Inform., Moscow, 1989, pp. 5–136, 310–314, 315. Translated from the English.
- [25] TITS, J. Classification of algebraic semisimple groups. In *Algebraic Groups and Discontinuous Subgroups (Proc. Sympos. Pure Math., Boulder, Colo., 1965)*. Amer. Math. Soc., Providence, R.I., 1966, pp. 33–62.
- [26] TITS, J. Strongly inner anisotropic forms of simple algebraic groups. *J. Algebra* 131, 2 (1990), 648–677.
- [27] TOTARO, B. The Chow ring of a classifying space. In *Algebraic K-theory (Seattle, WA, 1997)*, vol. 67 of *Proc. Sympos. Pure Math.* Amer. Math. Soc., Providence, RI, 1999, pp. 249–281.

MATHEMATICAL & STATISTICAL SCIENCES, UNIVERSITY OF ALBERTA, EDMONTON, CANADA  
Email address: [karpenko at ualberta.ca](mailto:karpenko@ualberta.ca), web page: [www.ualberta.ca/~karpenko](http://www.ualberta.ca/~karpenko)