

QUADRATIC FORMS IN I^n OF DIMENSION $2^n + 2^{n-1}$

CURTIS R. HARVEY AND NIKITA A. KARPENKO

ABSTRACT. For $n \geq 3$, confirming a weak version of a conjecture of Hoffmann, we show that every anisotropic quadratic form in I^n of dimension $2^n + 2^{n-1}$ splits over a finite extension of the base field of degree not divisible by 4. The first new case is $n = 4$, where we obtain a classification of the corresponding quadratic forms up to odd degree base field extensions and get this way a strong upper bound on their essential 2-dimension. As well, we compute the reduced Chow group of the maximal orthogonal grassmannian of the quadratic form and conclude that its canonical 2-dimension is $2^n + 2^{n-2} - 2$.

Let F be a field (of any characteristic) and let $I = I(F)$ be the Witt group of classes of even-dimensional non-degenerate quadratic forms over F defined as in [4, §8] (and denoted $I_q(F)$ there). For $n \geq 2$, we write $I^n = I^n(F)$ for the subgroup in $I(F)$ generated by the n -fold Pfister forms. We refer to [4, 9.B] for other equivalent definitions of $I^n(F)$ (denoted $I_q^n(F)$ there).

Any element of I is represented by an anisotropic quadratic form. By the Arason-Pfister Hauptsatz, the smallest possible dimension of a nonzero anisotropic quadratic form in I^n is 2^n (see [4, Theorem 23.7(1)] for the characteristic-free version). The quadratic forms in I^n of dimension 2^n are classified: as a consequence of the Arason-Pfister Hauptsatz and [4, Corollary 23.4)], they are exactly the forms similar to n -fold Pfister forms. In particular, any 2^n -dimensional quadratic form in I^n splits over a finite base field extension of degree dividing 2.

The smallest possible dimension exceeding 2^n of an anisotropic quadratic form in I^n is $2^n + 2^{n-1}$. For $n = 3$ this has been shown in [12] (characteristic $\neq 2$) and in [1] (arbitrary characteristic); for $n = 4$ in [6] (characteristic $\neq 2$) and in [5, Théorème 4.2.11] (characteristic 2); for arbitrary n and characteristic $\neq 2$ a proof has been given in [15, Theorem 5.4] and then extended to characteristic 2 in [13, Proposition 11.5].

For $n = 2$, quadratic forms in I^n of dimension $2^n + 2^{n-1}$ are the well-understood Albert forms. For $n \geq 3$, by a conjecture of Hoffmann ([6, Conjecture 2] for characteristic $\neq 2$ and [5, Conjecture 4.3.1] for characteristic 2), quadratic forms in I^n of dimension $2^n + 2^{n-1}$ should be classified as products of an Albert bilinear form (i.e., a 6-dimensional symmetric bilinear form of determinant -1) by a Pfister form (of foldness $n - 2$). In particular, such forms should split over a finite base field extension of degree dividing 2 as well.

Date: 10 Mar 2024. Revised: 1 Jul 2024.

Key words and phrases. Quadratic forms over fields; projective homogeneous varieties; Chow rings. *Mathematical Subject Classification (2020):* 11E04; 14C25.

This work has been supported by a Discovery Grant from the National Science and Engineering Research Council of Canada and completed when the second author was a visitor at the Université de Lorraine.

However, the two above conjectures are so far proved for $n = 3$ only: the proof for characteristic $\neq 2$ of [12] is extended to characteristic 2 in [5, Proposition 4.1.2]. The main result of the present note is

Theorem 1. *For any $n \geq 3$ and in any characteristic, every quadratic form in I^n of dimension $2^n + 2^{n-1}$ splits over a finite base field extension of degree not divisible by 4.*

Proof. Let X be a connected component of the highest orthogonal grassmannian of a quadratic form q in $I^n(F)$ of dimension $2^n + 2^{n-1}$. Theorem 1 means that the *index* $i(X)$ of the variety X , defined as the g.c.d. of the degrees of closed points on X , divides 2. In other terms, taking into account Springer's Theorem [4, Corollary 18.5], $i(X) = 2$ provided that q is not split.

We write \bar{X} for X over an algebraic closure of F and we write $\bar{\text{C}}\text{H}(X)$ for the ring given by the image of the change of field homomorphism $\text{CH}(X) \rightarrow \text{CH}(\bar{X})$ of the Chow rings. Note that the kernel of the change of field homomorphism is the ideal of the elements of finite order. For this reason, $\bar{\text{C}}\text{H}(X)$ is sometimes called the *reduced Chow group* of X .

By [4, Theorem 86.12], the ring $\text{CH}(\bar{X})$ is generated by certain homogeneous elements e_1, \dots, e_l of codimensions $1, \dots, l := 2^{n-1} + 2^{n-2} - 1$. It is convenient to define $e_i := 0$ for $i > l$. For any $i \geq 1$, the element e_i is characterized by the property that $(-1)^i 2e_i$ is the i th Chern class of the tautological vector bundle on \bar{X} (see [4, Proposition 87.13]); in particular, $2e_i \in \bar{\text{C}}\text{H}(X)$.

Since for any field extension K/F , the anisotropic part of the quadratic form q_K over the field K is either 0, or 2^n , or $2^n + 2^{n-1}$, it follows by [4, Corollary 88.6] (see also [4, Corollary 88.7]) that $e_i \in \bar{\text{C}}\text{H}(X)$ for all i different from $k := 2^{n-1} - 1$ and l . By [4, (86.15)], for any $i \geq 1$, we have

$$e_i^2 - 2e_{i-1}e_{i+1} + 2e_{i-2}e_{i+2} - \dots + (-1)^{i-1}2e_1e_{2i-1} + (-1)^i e_{2i} = 0 \in \text{CH}(\bar{X}).$$

In particular,

$$2e_k e_l = 2e_{k+1}e_{l-1} - 2e_{k+2}e_{l-2} + \dots \pm 2e_{m-1}e_{m+1} \pm e_m^2 \in \bar{\text{C}}\text{H}(X),$$

where $m := (k + l)/2$. Therefore $2e \in \bar{\text{C}}\text{H}(X)$, where $e \in \text{CH}(\bar{X})$ is the product $e_1 \dots e_l$ of all the generators. Since e is the class of a 0-cycle of degree 1 (see [4, Corollary 86.10]), the variety X (over F) possesses a 0-cycle of degree 2. \square

For $n = 4$, in view of [6, Proposition 4.1] and [5, Proposition 4.3.2], Theorem 1 provides a classification of the corresponding quadratic forms “up to odd degree extensions” which yields a strong upper bound on their *essential 2-dimension*. We provide details right below, starting with the classification result:

Theorem 2. *For a field F (of any characteristic), let q be a quadratic form in $I^4(F)$ of dimension $24 = 2^4 + 2^3$. Then there exists a finite field extension K/F of odd degree such that q_K is isomorphic to the tensor product of an Albert bilinear form by a Pfister form.*

Proof. By Theorem 1, we can find a finite field extension K/F of odd degree and a field extension L/K of degree dividing 2 such that q_L is split. The description of q_K then follows from [6, Proposition 4.1] (for characteristic $\neq 2$) and [5, Proposition 4.3.2] (for characteristic 2). \square

To formulate the result on the essential 2-dimension, let us consider the functor I_{24}^4 , associating to every extension field K of a fixed field F the set of isomorphism classes of quadratic forms in $I^4(K)$ of dimension 24. The essential 2-dimension of an element in $I_{24}^4(K)$ as well as the essential 2-dimension $\text{ed}_2 I_{24}^4$ of the functor I_{24}^4 are defined as in [3, §1].

Corollary 3. *One has $\text{ed}_2 I_{24}^4 \leq 7$.*

Proof. Writing F for the base field and taking $q \in I_{24}^4(K)$ for a field extension K/F , we find by Theorem 2 an odd degree field extension L/K such that q_L is isomorphic to the tensor product of the diagonal Albert bilinear form $\langle a_1, a_2, a_3, a_4, a_5, -a_1 a_2 a_3 a_4 a_5 \rangle$ by the Pfister form $\langle\langle b_1, b_2 \rangle\rangle$ for some nonzero $a_1, \dots, a_5, b_1 \in L$ and some $b_2 \in L$, where in characteristic $\neq 2$ the element b_2 is also nonzero. The subfield $F(a_1, \dots, a_5, b_1, b_2) \subset L$, whose transcendence degree over F is at most 7, is then a *field of definition* of q_L . It follows that $\text{ed}_2 q = \text{ed}_2 q_L \leq 7$ and so $\text{ed}_2 I_{24}^4 \leq 7$. \square

Recall that the essential 2-dimension is a 2-local version of and constitutes a lower bound for the *essential dimension*, measuring, informally speaking, how many independent parameters are required to describe an isomorphism class of the corresponding type of objects; in particular, $\text{ed}_2 I_{24}^4 \leq \text{ed} I_{24}^4$. For $n = 3$, since the description of the corresponding quadratic forms does not involve odd degree extensions, similar to the proof of Corollary 3 arguments show that $\text{ed}_2 I_{12}^3 \leq \text{ed} I_{12}^3 \leq 6$. In fact, in characteristic $\neq 2$, $\text{ed}_2 I_{12}^3 = \text{ed} I_{12}^3 = 6$ by [3, Theorem 7.1]: the lower bound $6 \leq \text{ed}_2 I_{12}^3$ is obtained by constructing a nontrivial degree 6 cohomological invariant with coefficients in $\mathbb{Z}/2\mathbb{Z}$ for I_{12}^3 .

For $n \geq 4$, assuming [6, Conjecture 2], one gets

$$\text{ed}_2 I_{2^n+2^{n-1}}^n \leq \text{ed} I_{2^n+2^{n-1}}^n \leq n + 3.$$

Finally, one has $\text{ed}_2 I_{2^n}^n = \text{ed} I_{2^n}^n = n + 1$ for any n . Indeed, as already mentioned, any $q \in I_{2^n}^n$ is isomorphic to $b \cdot \langle\langle b_1, b_2, \dots, b_n \rangle\rangle$ for some $n + 1$ parameters b, b_1, \dots, b_n , ensuring that $n + 1$ is an upper bound for $\text{ed} I_{2^n}^n$. On the other hand, associating in characteristic $\neq 2$ to q the symbol (b, b_1, \dots, b_n) in the $(n + 1)$ st Galois cohomology group with coefficients in $\mathbb{Z}/2\mathbb{Z}$, one gets a nontrivial degree $n + 1$ cohomological invariant showing that $n + 1$ is a lower bound for $\text{ed}_2 I_{2^n}^n$ (see [11, Theorem 3.4]). The non-triviality of the cohomological invariant is shown in [2, §3]. The characteristic 2 case is treated similarly using cohomological invariants with values in étale motivic cohomology groups (cf. [14, §3] and especially [14, Proof of Lemma 3.1]); the non-triviality of the cohomological invariant follows from [7].

To conclude, let us return to the case of arbitrary $n \geq 2$. Let X be the highest orthogonal grassmannian of an anisotropic quadratic form $q \in I^n$. If $\dim q = 2^n$, then $i(X) = 2$ and therefore the ring $\overline{\text{CH}}(X)$ contains $2\text{CH}(\overline{X})$. By [4, Corollary 88.6], $\overline{\text{CH}}(X)$ also contains the elements e_1, \dots, e_{2^n-1-2} – the generators of the ring $\text{CH}(\overline{X})$ with exception of the very last one e_{2^n-1-1} . Since $i(X) \neq 1$, we conclude that $\overline{\text{CH}}(X)$ is exactly the subring in $\text{CH}(\overline{X})$ generated by $2\text{CH}(\overline{X})$ and e_1, \dots, e_{2^n-1-2} (cf. [4, Example 88.10]).

Now let us assume that $\dim q = 2^n + 2^{n-1}$, where $n \geq 3$. Since $i(X) = 2$ by Theorem 1, we still have the inclusion $\overline{\text{CH}}(X) \supset 2\text{CH}(\overline{X})$. Besides, it has been shown in the proof of Theorem 1 that $\overline{\text{CH}}(X) \ni e_i$ for all i except $i = k := 2^{n-1} - 1$ and $i = l := 2^{n-1} + 2^{n-2} - 1$.

Theorem 4. *For any $n \geq 3$ and any anisotropic quadratic form q in I^n of dimension $2^n + 2^{n-1}$, the ring $\overline{\text{CH}}(X)$ of its highest grassmannian X is generated by $2\text{CH}(\overline{X})$ and all e_i with $i \notin \{k, l\}$.*

Proof. Since $\overline{\text{CH}}(X) \supset 2\text{CH}(\overline{X})$, it suffices to show that the ring $\overline{\text{Ch}}(X)$ is generated by e_i with $i \notin \{k, l\}$, where $\text{Ch}(X) := \text{CH}(X)/2\text{CH}(X)$ and $\overline{\text{Ch}}(X) := \text{Im}(\text{Ch}(X) \rightarrow \text{Ch}(\overline{X}))$. By [4, Theorem 87.7] (originally proved in [16]), it suffices to show that neither e_k nor e_l is in $\overline{\text{Ch}}(X)$.

By [4, Corollary 82.3] once again, the anisotropic part of q over the function field of its quadric Y has dimension 2^n . By [4, Corollary 88.7], we conclude that $e_k \notin \overline{\text{Ch}}(X)$.

Finally, let us assume that $e_l \in \overline{\text{Ch}}(X)$ and seek for a contradiction. By [4, Theorem 90.3] (originally proved in [16]), the *canonical 2-dimension* of the variety X equals k and does not change when the base field is extended to the function field of Y . It follows by [10, Theorem 3.2] that a shift of the *upper Chow motive* $U(X)$ with coefficients $\mathbb{Z}/2\mathbb{Z}$ is a direct summand of the motive of Y . On the other hand, by [4, Lemma 82.4], the complete motivic decomposition of the quadric Y consists only of shifts of the upper motive $U(Y)$. Moreover, since the variety $X_{F(Y)}$ has no 0-cycle of odd degree, the motives $U(Y)$ and $U(X)$ are not isomorphic, see [9, Corollary 2.15]. The contradiction obtained proves Theorem 4. \square

Regarding the motives of the varieties X and Y from the above proof, each of them decomposes in a finite direct sum of indecomposable motives; moreover, by [9, Corollary 2.6], such a decomposition is unique in the usual sense. The upper motive $U(X)$ (resp., $U(Y)$) is defined as the summand with nontrivial Ch^0 (unique in any decomposition given). By [9, Corollary 2.15], the motives $U(X)$ and $U(Y)$ are isomorphic if and only if each of the two varieties $X_{F(Y)}$ and $Y_{F(X)}$ possesses a 0-cycle of odd degree.

Let us also recall that *canonical dimension* $\text{cd}(X)$ of a smooth projective variety X is the minimum of dimension of the image of a rational self-map $X \dashrightarrow X$, c.f. [8]. See also [8, Definition 1.3] for a definition using the essential dimension of a certain functor related to X . Canonical 2-dimension, which appeared in the above proof, is its 2-local version also providing a lower bound for it.

Corollary 5. *For any anisotropic q as in Theorem 4, the canonical 2-dimension $\text{cd}_2(X)$ of its highest grassmannian X is equal to $k + l = 2^n + 2^{n-2} - 2$.*

Proof. As in the proof of Theorem 4, $e_i \in \overline{\text{Ch}}(X)$ for $i \notin \{k, l\}$. Moreover, it follows from Theorem 4 (and has been shown in its proof explicitly) that neither e_k nor e_l is in $\overline{\text{Ch}}(X)$. Thus in terms of the J -invariant in [4, Chapter 88], we have $J(q) = \{k, l\}$ and so [4, Theorem 90.3] tells us that $\text{cd}_2(X) = k + l$. \square

ACKNOWLEDGEMENTS. We thank Detlev Hoffmann for providing reference [5]. We also thank Alexander Merkurjev and Alexander Vishik for useful comments.

REFERENCES

- [1] BAEZA, R. *Quadratic forms over semilocal rings*. Lecture Notes in Mathematics, Vol. 655. Springer-Verlag, Berlin-New York, 1978.
- [2] BERHUY, G., AND FAVI, G. Essential dimension: a functorial point of view (after A. Merkurjev). *Doc. Math.* 8 (2003), 279–330 (electronic).
- [3] CHERNOUSOV, V., AND MERKURJEV, A. Essential dimension of spinor and Clifford groups. *Algebra Number Theory* 8, 2 (2014), 457–472.
- [4] ELMAN, R., KARPENKO, N., AND MERKURJEV, A. *The algebraic and geometric theory of quadratic forms*, vol. 56 of *American Mathematical Society Colloquium Publications*. American Mathematical Society, Providence, RI, 2008.
- [5] FAIVRE, F. Liaison des formes de Pfister et corps de fonctions de quadriques en caractéristique 2. Ph.D. thesis, Université de Franche-Compte, Besançon (2006).
- [6] HOFFMANN, D. W. On the dimensions of anisotropic quadratic forms in I^4 . *Invent. Math.* 131, 1 (1998), 185–198.
- [7] IZHBOLDIN, O. T. On the cohomology groups of the field of rational functions. In *Mathematics in St. Petersburg*, vol. 174 of *Amer. Math. Soc. Transl. Ser. 2*. Amer. Math. Soc., Providence, RI, 1996, pp. 21–44.
- [8] KARPENKO, N. A. Canonical dimension. In *Proceedings of the International Congress of Mathematicians. Volume II* (New Delhi, 2010), Hindustan Book Agency, pp. 146–161.
- [9] KARPENKO, N. A. Upper motives of algebraic groups and incompressibility of Severi-Brauer varieties. *J. Reine Angew. Math.* 677 (2013), 179–198.
- [10] KARPENKO, N. A. Minimal canonical dimensions of quadratic forms. *Doc. Math.* (2015), 367–385.
- [11] MERKURJEV, A. S. Essential dimension: a survey. *Transform. Groups* 18, 2 (2013), 415–481.
- [12] PFISTER, A. Quadratische Formen in beliebigen Körpern. *Invent. Math.* 1 (1966), 116–132.
- [13] PRIMOZIC, E. Motivic Steenrod operations in characteristic p . *Forum Math. Sigma* 8 (2020), Paper No. e52, 25.
- [14] TOTARO, B. Essential dimension of the spin groups in characteristic 2. *Comment. Math. Helv.* 94, 1 (2019), 1–20.
- [15] VISHIK, A. Motives of quadrics with applications to the theory of quadratic forms. In *Geometric methods in the algebraic theory of quadratic forms*, vol. 1835 of *Lecture Notes in Math*. Springer, Berlin, 2004, pp. 25–101.
- [16] VISHIK, A. On the Chow groups of quadratic Grassmannians. *Doc. Math.* 10 (2005), 111–130 (electronic).

MATHEMATICAL & STATISTICAL SCIENCES, UNIVERSITY OF ALBERTA, EDMONTON, CANADA
 Email address: rcharvey@ualberta.ca

MATHEMATICAL & STATISTICAL SCIENCES, UNIVERSITY OF ALBERTA, EDMONTON, CANADA
 Email address: karpenko@ualberta.ca
 URL: www.ualberta.ca/~karpenko