

**CODIMENSION 2 CYCLES
ON QUADRATIC WEIL TRANSFER
OF BIQUATERNIONIC SEVERI-BRAUER VARIETY**

NIKITA A. KARPENKO

ABSTRACT. Let F be a field, B a biquaternion F -algebra, L/F an étale quadratic extension, X the Weil transfer with respect to L/F of the Severi-Brauer variety of B_L . We show that the Chow group of codimension 2 cycle classes on X is torsion-free.

Our Chow groups are those with integral coefficients. The motives used in the proof are the Grothendieck Chow motives (still with the integral coefficients) as in [1, §64].

Theorem 1. *Let F be a field, B a biquaternion F -algebra, L/F an étale quadratic extension, X the Weil transfer with respect to L/F of the Severi-Brauer variety of B_L . Then the Chow group $\mathrm{CH}^2(X)$ is torsion-free.*

Proof. If L is not a field, then $L \simeq F \times F$ and $X \simeq \mathrm{SB}(B) \times \mathrm{SB}(B)$. Since X is a rank 3 projective bundle over $\mathrm{SB}(B)$, the Chow motive of X is then isomorphic to the sum $M \oplus M(1) \oplus M(2) \oplus M(3)$, where M is the motive of the variety $\mathrm{SB}(B)$. Therefore the total Chow group of X is a direct sum of four (shifted) copies of the total Chow group of $\mathrm{SB}(B)$. Since the total Chow group of $\mathrm{SB}(B)$ is torsion-free [5, Corollary 7], the total Chow group of X is also torsion-free; in particular, so is the group $\mathrm{CH}^2(X)$.

Now let us assume that L is a field and B_L is not a division algebra. In this case, by [8, Proposition 16.2], there exists a quaternion F -algebra Q such that the algebra B_L is isomorphic to the algebra of (2×2) -matrices over Q_L . Let now M be the motive (over L) of the Severi-Brauer variety of Q_L . Then the motive of $\mathrm{SB}(B_L)$ is isomorphic to $M \oplus M(2)$ [4, Theorem 1.3.1]. Therefore, by [3, Lemma 2.1], the motive (over F) of X is isomorphic to the sum $R_{L/F}M \oplus (R_{L/F}M)(4) \oplus \mathrm{cor}_{L/F}(M \otimes \sigma M)(2)$, where σ is the non-trivial automorphism of L/F , $R_{L/F}$ is the motivic Weil transfer functor of [6], and $\mathrm{cor}_{L/F}$ is the motivic corestriction functor introduced in [7]. The motive $R_{L/F}M$ is the motive of the variety $R_{L/F} \mathrm{SB}(Q_L)$ (abusing notation, we write $R_{L/F}$ also for the Weil transfer on varieties). This variety is a smooth projective quadric surface; its total Chow group is torsion-free. Concerning the remaining summand $\mathrm{cor}_{L/F}(M \otimes \sigma M)(2)$, we first note that the corestriction functor preserves the Chow group. The motive $M \otimes \sigma M$ is the motive of the variety $\mathrm{SB}(Q_L) \times \mathrm{SB}(Q_L)$ which is a rank 1 projective bundle over $\mathrm{SB}(Q_L)$. Since the total Chow group of $\mathrm{SB}(Q_L)$ is torsion-free, the total Chow group of $\mathrm{cor}_{L/F}(M \otimes \sigma M)$ is torsion-free and it follows that the total Chow group of X is torsion-free in this case also.

To finish the proof of Theorem 1, we consider the remaining case where B_L is a skew field (and L is a field). We consider the Grothendieck group $K(X)$ of classes of coherent \mathcal{O}_X -modules together with its topological filtration, and we are going to show that the

total associated graded group is torsion-free. Since its codimension 2 graded piece is isomorphic to $\text{CH}^2(X)$, this is sufficient for our purpose.

We start by describing the group $K(X)$. Note that the Grothendieck group (but not the topological filtration on it) is computed for any projective homogeneous variety in [9]. More precisely, for any projective homogeneous variety Y over a field l , an isomorphism $K_0(Y) \simeq K_0(A)$ with certain separable l -algebra A is constructed (and $K(Y) = K_0(Y)$). Moreover, if l/k is a finite separable field extension, then $K_0(R_{l/k}Y) \simeq K_0(N_{l/k}A)$, where $N_{l/k}A$ is the norm algebra.

We turn back to our particular variety X . According to [9] and since X is projective homogeneous, the restriction homomorphism $K(X) \rightarrow K(X_E)$ is injective for any field extension E/F . We will describe $K(X)$ as a subgroup of $K(X_E)$ with E/F such that X_E is “simple enough”.

Let E/F be a maximal subfield of the F -algebra B . Since B_L is a skew field, $EL := E \otimes_F L$ is a field. The EL -variety X_{EL} is isomorphic to $\mathbb{P}^3 \times \mathbb{P}^3$. Let x be the class of $\mathcal{O}(-1)$ in $K(\mathbb{P}^3)$. The ring $K(\mathbb{P}^3)$ is generated by x with the only relation $(1-x)^4 = 0$. In particular, the group $K(\mathbb{P}^3)$ is freely generated by x^i , $i \in \{0, 1, 2, 3\}$. The ring $K(X_{EL}) = K(\mathbb{P}^3 \times \mathbb{P}^3)$ is isomorphic to the tensor product of 2 copies of the ring $K(\mathbb{P}^3) = \mathbb{Z}[x]/(1-x)^4$, the isomorphism is given by the exterior product. It follows that the group $K(X_{EL})$ is freely generated by $x^i \times x^j$ with $i, j \in \{0, 1, 2, 3\}$.

The non-trivial element of the Galois group of EL/E acts on $K(X_{EL})$ by exchanging the factors. The subgroup (subring) $K(X_E) \subset K(X_{EL})$ is the set of invariant elements; it is freely generated by $x^i \times x^i$, $i \in \{0, 1, 2, 3\}$ and $x^i \times x^j + x^j \times x^i$ with $i \neq j \in \{0, 1, 2, 3\}$.

The subgroup (subring) $K(X) \subset K(X_E)$ is generated by certain multiples of the generators of $K(X_E)$. Namely, by $x^i \times x^i$, $i \in \{0, 1, 2, 3\}$ (each of which comes with the coefficient $1 = \text{ind } N_{L/F} B_L^{\otimes i}$), by $x^i \times x^j + x^j \times x^i$ with even $i+j$ (still with the coefficient $1 = \text{ind } B_L^{\otimes(i+j)}$), and, finally, by $4(x^i \times x^j + x^j \times x^i)$ with odd $i+j$ (the coefficient 4 being the index of $B_L^{\otimes(i+j)}$). This is our desired description of $K(X)$ and now we come to the consideration of its topological filtration.

First of all, we know the topological filtration on $K(X_{EL})$. It is easily described in terms of $h^i \times h^j$, a different system of additive generators, where $h := 1 - x$ is the hyperplane class. In terms of these generators, the topological filtration is simply the degree filtration: the term of codimension d is generated by all $h^i \times h^j$ with $i + j \geq d$.

We claim that the topological filtration on $K(X_E)$ is induced by that on $K(X_{EL})$. In general, we only have the inclusion $K(X_E)^{(d)} \subset K(X_{EL})^{(d)}$ for any d . This gives a homomorphism of the associated graded groups. By a transfer argument, the kernel of this homomorphism consists of torsion. On the other hand, as we already know, the total Chow group of X_E has no torsion. Therefore, the graded group of the topological filtration on $K(X_E)$ is also torsion-free. It follows that the homomorphism of the graded groups is injective and this means that the inclusion $K(X_E)^{(d)} \subset K(X_{EL})^{(d)}$ is in fact the equality $K(X_E)^{(d)} = K(X_{EL})^{(d)} \cap K(X_E)$, that is, the filtration on $K(X_E)$ is the induced one.

We come to the crucial point of the proof. We claim that the graded group of $K(X)$ is also torsion-free. In other words, we claim that the filtration on $K(X)$ is induced by that on $K(X_E)$. We prove this by the method of [2, Lemma 2.1]. Since the index of $K(X)$ in

$K(X_E)$ is $4^4 = 2^8$, it suffices to show that the order of the cokernel of the homomorphism of the graded groups is at most 2^8 .

The graded group of $K(X_E)$ is generated by the classes of the following elements:

$$\begin{aligned}
\text{codim} = 0 : & & & 1 \times 1, \\
\text{codim} = 1 : & & & 1 \times h + h \times 1, \\
\text{codim} = 2 : & & & 1 \times h^2 + h^2 \times 1 \quad \text{and} \quad h \times h, \\
\text{codim} = 3 : & & & 1 \times h^3 + h^3 \times 1 \quad \text{and} \quad h \times h^2 + h^2 \times h, \\
\text{codim} = 4 : & & & h \times h^3 + h^3 \times h \quad \text{and} \quad h^2 \times h^2, \\
\text{codim} = 5 : & & & h^2 \times h^3 + h^3 \times h^2, \\
\text{codim} = 6 : & & & h^3 \times h^3.
\end{aligned}$$

The cokernel order in codimension 0 is 1.

The image in codimension 1 contains the class of the element $1 \times h + h \times 1$. Indeed,

$$\begin{aligned}
K(X) \ni \alpha := 1 \times 1 - x \times x &= 1 \times 1 - (1 - h) \times (1 - h) = \\
& 1 \times h + h \times 1 - h \times h \equiv 1 \times h + h \times 1 \pmod{K(X_E)^{(2)}}.
\end{aligned}$$

Since $\alpha \in K(X_E)^{(1)}$ and the quotient $K(X)^{(0)}/K(X)^{(1)}$ is torsion-free, $\alpha \in K(X)^{(1)}$. It follows that the cokernel order in codimension 1 is also 1.

The image in codimension 2 contains the class of $1 \times h^2 + h^2 \times 1 + 2(h \times h)$, as this is the square of the class of $1 \times h + h \times 1$. It also contains the class of $4(h \times h)$: by a transfer argument applied to the degree 4 field extension E/F , every element of the graded group of $K(X_E)$ multiplied by 4 is in the image. It follows that the cokernel order in codimension 2 is at most 4.

The crucial thing happens in codimension 3. First of all we have the class of

$$(1 \times h + h \times 1)^3 = (1 \times h^3 + h^3 \times 1) + 3(h \times h^2 + h^2 \times h)$$

in the image. On the other hand, we have a morphism $\text{SB}(B) \rightarrow R_{L/F} \text{SB}(B_L) = X$ becoming over L the diagonal morphism $\text{SB}(B_L) \rightarrow \text{SB}(B_L) \times \text{SB}(B_L)$. This is the morphism corresponding to the identity under the canonical bijection

$$\text{Mor}(\text{SB}(B), R_{L/F} \text{SB}(B_L)) = \text{Mor}(\text{SB}(B_L), \text{SB}(B_L))$$

which we have because the Weil restriction functor is right-adjoint to the scalar extension functor. Therefore, $\text{SB}(B)$ is a closed subvariety of X whose class in $\text{CH}^3(X_E) = K(X_E)^{(3)}/K(X_E)^{(4)}$ is equal to

$$(1 \times h^3 + h^3 \times 1) + (h \times h^2 + h^2 \times h).$$

It follows that the cokernel order in codimension 3 is at most $2 = \left| \det \begin{pmatrix} 1 & 1 \\ 3 & 1 \end{pmatrix} \right|$. We also see that the class of $2(h \times h^2 + h^2 \times h)$ is in the image.

In codimension 4 the image contains $2(h^2 \times h^2)$ as this is $(1 \times h + h \times 1)^4$ modulo 4. We also have $2(h \times h^3 + h^3 \times h)$ as the product of $2(h \times h^2 + h^2 \times h)$ by $1 \times h + h \times 1$ modulo 4. Therefore the cokernel order in codimension 4 is at most 4.

The doubled generator in codimension 5 is in the image as the product of $2(h \times h^3 + h^3 \times h)$ by $1 \times h + h \times 1$. So, the cokernel order in codimension 5 is at most 2.

Finally, the cokernel order in codimension 6 is at most 4.

Summarizing (multiplying) over all codimension, we see that the total cokernel order is at most $1 \cdot 1 \cdot 4 \cdot 2 \cdot 4 \cdot 2 \cdot 4 = 2^8$, as desired. \square

REFERENCES

- [1] ELMAN, R., KARPENKO, N., AND MERKURJEV, A. *The algebraic and geometric theory of quadratic forms*, vol. 56 of *American Mathematical Society Colloquium Publications*. American Mathematical Society, Providence, RI, 2008.
- [2] IZHBOLDIN, O. T., AND KARPENKO, N. A. On the group $H^3(F(\psi, D)/F)$. *Doc. Math.* 2 (1997), 297–311 (electronic).
- [3] KARPENKO, N. A. Incompressibility of quadratic Weil transfer of generalized Severi-Brauer varieties. *Linear Algebraic Groups and Related Structures* (preprint server) 362 (2009, Oct 28), 12 pages.
- [4] KARPENKO, N. A. Grothendieck Chow motives of Severi-Brauer varieties. *Algebra i Analiz* 7, 4 (1995), 196–213.
- [5] KARPENKO, N. A. On topological filtration for Severi-Brauer varieties. II. In *Mathematics in St. Petersburg*, vol. 174 of *Amer. Math. Soc. Transl. Ser. 2*. Amer. Math. Soc., Providence, RI, 1996, pp. 45–48.
- [6] KARPENKO, N. A. Weil transfer of algebraic cycles. *Indag. Math. (N.S.)* 11, 1 (2000), 73–86.
- [7] KARPENKO, N. A. Upper motives of outer algebraic groups. In *Quadratic forms, linear algebraic groups, and cohomology*, vol. (to appear) of *Dev. Math.* Springer, New York, 2010.
- [8] KNUS, M.-A., MERKURJEV, A., ROST, M., AND TIGNOL, J.-P. *The book of involutions*, vol. 44 of *American Mathematical Society Colloquium Publications*. American Mathematical Society, Providence, RI, 1998. With a preface in French by J. Tits.
- [9] PANIN, I. A. On the algebraic K -theory of twisted flag varieties. *K-Theory* 8, 6 (1994), 541–585.

UPMC UNIV PARIS 06, INSTITUT DE MATHÉMATIQUES DE JUSSIEU, F-75252 PARIS, FRANCE

Web page: www.math.jussieu.fr/~karpenko

E-mail address: [karpenko at math.jussieu.fr](mailto:karpenko@math.jussieu.fr)