

A Research Story: Compound Equations and Dynamics. Part 3

James Muldowney, University of Alberta

July 25, 2019

Curves and Surfaces

A smooth *curve* γ in \mathbb{R}^n is a C^1 function $s \rightarrow x(s)$, $s \in I \subset \mathbb{R}$, $x(s) \in \mathbb{R}^n$.

A measure of the *length* of γ is

$$l(\gamma) = \int_{\gamma} dl \stackrel{\text{def}}{=} \int_I \left\| \frac{dx}{ds}(s) \right\| ds$$

where $\|\cdot\|$ is a norm on \mathbb{R}^n . For example, the euclidean norm $\|x\| = \sqrt{(x_1)^2 + \cdots + (x_n)^2}$ gives the usual measure of length

$$l(\gamma) = \int_I \sqrt{\frac{dx_1}{ds}^2 + \cdots + \frac{dx_n}{ds}^2} ds$$

A smooth 2-surface σ in \mathbb{R}^n is a C^1 function $(s_1, s_2) \rightarrow x(s_1, s_2)$, $(s_1, s_2) \in U \subset \mathbb{R}^2$, $x(s_1, s_2) \in \mathbb{R}^n$.

A measure of the *area* of σ is

$$a_2(\sigma) = \int_{\sigma} da \stackrel{\text{def}}{=} \int_U \|x_{s_1} \wedge x_{s_2}\| ds_1 ds_2$$

where $x_{s_i} = \frac{\partial}{\partial s_i} x(s_1, s_2)$ and $\|\cdot\|$ is a norm on $\mathbb{R}^{\binom{n}{2}}$. If $\|\cdot\|$ is the Euclidean norm we have

$$a_2(\sigma) = \int_U \sqrt{\sum_{1 \leq i < j \leq n} \left(\frac{\partial(x_i, x_j)}{\partial(s_1, s_2)} \right)^2} ds_1 ds_2$$

where

$$\frac{\partial(x_i, x_j)}{\partial(s_1, s_2)} = \det \begin{bmatrix} \frac{\partial x_i}{\partial s_1} & \frac{\partial x_i}{\partial s_2} \\ \frac{\partial x_j}{\partial s_1} & \frac{\partial x_j}{\partial s_2} \end{bmatrix}.$$

A smooth k -surface σ in \mathbb{R}^n is a C^1 function

$$(s_1, \dots, s_k) \rightarrow x(s_1, \dots, s_k), \quad s_1, \dots, s_k \in U \subset \mathbb{R}^k, \quad x(s_1, \dots, s_k) \in \mathbb{R}^n.$$

A measure of the k -area of σ is

$$a_k(\sigma) = \int_{\sigma} da_k \stackrel{\text{def}}{=} \int_U \|x_{s_1} \wedge \dots \wedge x_{s_k}\| ds_1 \cdots ds_k$$

where $x_{s_i} = \frac{\partial}{\partial s_i} x(s_1, \dots, s_k)$ and $\|\cdot\|$ is a norm on $\mathbb{R}^{\binom{n}{k}}$. If $\|\cdot\|$ is the Euclidean norm we have

$$a_k(\sigma) = \int_U \sqrt{\sum_{1 \leq i_1 < \dots < i_k \leq n} \frac{\partial(x_{i_1}, \dots, x_{i_k})^2}{\partial(s_1, \dots, s_k)}} ds_1 ds_2$$

where

$$\frac{\partial(x_{i_1}, \dots, x_{i_k})}{\partial(s_1, \dots, s_k)} = \det \begin{bmatrix} \frac{\partial x_{i_1}}{\partial s_1} & \frac{\partial x_{i_1}}{\partial s_2} & \cdots & \frac{\partial x_{i_1}}{\partial s_k} \\ \frac{\partial x_{i_2}}{\partial s_1} & \frac{\partial x_{i_2}}{\partial s_2} & \cdots & \frac{\partial x_{i_2}}{\partial s_k} \\ \vdots & \vdots & \ddots & \vdots \\ \frac{\partial x_{i_k}}{\partial s_1} & \frac{\partial x_{i_k}}{\partial s_2} & \cdots & \frac{\partial x_{i_k}}{\partial s_k} \end{bmatrix}.$$

Nonlinear Differential Equations

$$f \in C^1(\mathbb{R}^n \rightarrow \mathbb{R}^n)$$

$$\dot{x} = f(x) \tag{N}$$

Solution : $x(t) = \phi(t) = \phi(t, x_0)$, is uniquely determined by $x(0) = x_0$ and, for simplicity, we will only consider equations for which solutions exist for all $t > 0$

If $\phi(t, x_0) = x_0$ for all t , then x_0 is called an *equilibrium*.

If $\phi(t + \omega) = \phi(t)$, $\omega > 0$, the solution is *periodic* of period ω .

An *orbit* (positive semi-orbit) is a set $\{\phi(t) : 0 \leq t < \infty\}$.

The orbit of an equilibrium is a single point.

The orbit of a periodic solution is a simple closed curve (Jordan curve)

Linearization about a solution $\phi(t)$:

$$\dot{y} = \frac{\partial f}{\partial x}(\phi(t)) y \quad (\text{L})$$

Solution is:

$$y(t) = \frac{\partial \phi}{\partial x_0}(t, x_0) y(0), \quad x_0 = \phi(0)$$

$$Y(t) = \frac{\partial \phi}{\partial x_0}(t, x_0) \text{ fundamental matrix, } Y(0) = I$$

"Proof": $y = \phi(t, x_0)$ solves $\dot{y} = f(y)$

$$\Rightarrow \frac{\partial \phi}{\partial t}(t, x_0) = f(\phi(t, x_0))$$

Differentiate with respect to x_0

$$\Rightarrow \frac{\partial^2 \phi}{\partial t \partial x_0}(t, x_0) = \frac{\partial^2 \phi}{\partial x_0 \partial t}(t, x_0) = \frac{\partial f}{\partial x}(\phi(t, x_0)) \frac{\partial \phi}{\partial x_0}(t, x_0)$$

$$\Rightarrow \dot{Y} = \frac{\partial f}{\partial x}(\phi(t)) Y$$

The k -th compound equation of (L) is:

$$\dot{z} = \frac{\partial f^{[k]}}{\partial x} (\phi(t)) z \quad (L_k)$$

Solution: $z(t) = \frac{\partial \phi^{(k)}}{\partial x_0} (t, x_0) z(0)$, $x_0 = \phi(0)$

The case $k = n$ of (L_k) is the Liouville equation:

$$\dot{z} = \operatorname{div} f (\phi(t)) z \quad (L_n)$$

Solution: $z(t) = \det \frac{\partial \phi}{\partial x_0} (t, x_0) z(0)$, $x_0 = \phi(0)$

Suppose that $D \subset \mathbb{R}^n$ has finite n -dimensional measure $a_n(D)$, then the measure of $\phi(t, D)$ is

$$a_n(\phi(t, D)) = \int_{x \in \phi(t, D)} dx = \int_{x_0 \in D} \left| \det \frac{\partial \phi}{\partial x_0}(t, x_0) \right| dx_0$$

$(L_n) \Rightarrow \det \frac{\partial \phi}{\partial x_0}(t, x_0) = \exp \left[\int_0^t \operatorname{div} f(\phi(s, x_0)) ds \right]$. So, for example, if $\operatorname{div} f < 0$ in \mathbb{R}^n , then the measure of the set $\phi(t, D)$ decreases with time.

When $n = 2$ this observation implies that no simply connected region where $\operatorname{div} f < 0$ can contain a non-trivial periodic orbit of (L) . This is known as *Bendixson's Condition*. Most textbooks prove this as a very nice application of Green's Theorem.

Stability of the linearized equations (L) and its compounds (L_k) have many implications for the dynamics of (N)

If $\gamma_0: x = x_0(s)$, $0 \leq s \leq 1$ is a curve in \mathbb{R}^n , then $\gamma_t: x = \phi(t, x_0(s))$, $0 \leq s \leq 1$ is also a curve in \mathbb{R}^n for each $t \geq 0$.

$$\begin{aligned} l\gamma_0 &= \int_0^1 \left\| \frac{d}{ds} x_0(s) \right\| ds \\ l\gamma_t &= \int_0^1 \left\| \frac{d}{ds} \phi(t, x_0(s)) \right\| ds = \int_0^1 \left\| \frac{\partial \phi}{\partial x_0}(t, x_0(s)) \frac{d}{ds} (x_0(s)) \right\| ds \\ &\leq \int_0^1 \left\| \frac{\partial \phi}{\partial x_0}(t, x_0(s)) \right\| \left\| \frac{d}{ds} (x_0(s)) \right\| ds \end{aligned}$$

We can conclude for example that, if $\left\| \frac{\partial \phi}{\partial x_0}(t, x_0) \right\| \xrightarrow{t \rightarrow \infty} 0$ uniformly with respect to $x_0 \in \mathbb{R}^n$, then

- there is at most one equilibrium of (N) and,
- any equilibrium attracts all other orbits

If $\sigma_0 : (s_1, s_2) \rightarrow x(s_1, s_2)$ is a 2-surface in \mathbb{R}^n then so also is $\sigma_t : (s_1, s_2) \rightarrow \phi(t, x(s_1, s_2))$.

We can use similar ideas to get higher dimensional *Bendixson Conditions* to rule out the existence of periodic orbits. These are conditions on (L_2) that typically imply that some measure of surface area decreases in the dynamics. Another related type of condition would imply that $a_2 \sigma_t \xrightarrow{t \rightarrow \infty} 0$.

The central idea is to observe that a periodic orbit γ is invariant in the dynamics, $\phi(t, \gamma) = \gamma$. So, if Σ_0 is any surface which has γ as its boundary, then $\Sigma_t = \phi(t, \Sigma_0)$ is also a surface with γ as boundary. But if, among all surfaces with boundary γ , Σ_0 is a surface with minimum area and (N) diminishes area we would contradict the minimality of Σ_0 . So no such invariant closed curve can exist.

The following are Bendixson conditions for various measures of 2-surface area. Each reduces to the classical result when $n = 2$:

$$\lambda_1 + \lambda_2 < 0 \text{ (RA Smith)}$$

$$\max_{r \neq s} \left\{ \frac{\partial f_r}{\partial x_r} + \frac{\partial f_s}{\partial x_s} + \sum_{q \neq r,s} \left(\left| \frac{\partial f_r}{\partial x_q} \right| + \left| \frac{\partial f_s}{\partial x_q} \right| \right) \right\} < 0$$

$$\max_{r \neq s} \left\{ \frac{\partial f_r}{\partial x_r} + \frac{\partial f_s}{\partial x_s} + \sum_{q \neq r,s} \left(\left| \frac{\partial f_q}{\partial x_r} \right| + \left| \frac{\partial f_q}{\partial x_s} \right| \right) \right\} < 0$$

$$\lambda_{n-1} + \lambda_n > 0$$

$$\min_{r \neq s} \left\{ \frac{\partial f_r}{\partial x_r} + \frac{\partial f_s}{\partial x_s} - \sum_{q \neq r,s} \left(\left| \frac{\partial f_r}{\partial x_q} \right| + \left| \frac{\partial f_s}{\partial x_q} \right| \right) \right\} > 0$$

$$\min_{r \neq s} \left\{ \frac{\partial f_r}{\partial x_r} + \frac{\partial f_s}{\partial x_s} - \sum_{q \neq r,s} \left(\left| \frac{\partial f_q}{\partial x_r} \right| + \left| \frac{\partial f_q}{\partial x_s} \right| \right) \right\} > 0$$

$\lambda_1 \geq \lambda_2 \geq \dots \geq \lambda_n$ are the eigenvalues of $\frac{1}{2} \left(\frac{\partial f}{\partial x}^* + \frac{\partial f}{\partial x} \right)$

General Compounds

M Fiedler, Czech Math J 24(1974), pp 392 - 402

$\mathbb{X} \subset \mathbb{Y}$: General compound $A^{[k]} \in \mathcal{L} \left(\wedge^k \mathbb{X} \rightarrow \wedge^k \mathbb{Y} \right)$. $0 \leq m \leq k$

$$A^{[k,m]} \left(v^1 \wedge \dots \wedge v^k \right) \stackrel{\text{def}}{=} \sum_{(\varepsilon_1, \dots, \varepsilon_k)} A^{\varepsilon_1} v^1 \wedge A^{\varepsilon_2} v^2 \wedge \dots \wedge A^{\varepsilon_k} v^k$$

$$\varepsilon_i \in \{0, 1\}, \quad \varepsilon_1 + \dots + \varepsilon_k = m, \quad A^0 = I$$

$$A^{[k,0]} = I^{(k)}, \quad A^{[k,1]} = A^{[k]}, \quad A^{[k,k]} = A^{(k)}$$

$$D_h^m (I + hA)^{(k)} \Big|_{t=0} = m! A^{[k,m]}$$

$$D_h^m (I + hA)^{(k)} \Big|_{t=0} = m! A^{[k,m]}$$

$$\begin{aligned} (I + hA)^{(k)} &= \sum_{m=0}^k h^m A^{[k,m]} \\ &= hA^{[k,1]} + h^2 A^{[k,2]} + \dots + h^k A^{[k,k]} \end{aligned}$$

If $\lambda_1, \dots, \lambda_n$ are the eigenvalues of A with eigenvectors v^1, \dots, v^n , then the eigenvalues of $(I + hA)^{(k)}$ are

$$h(\lambda_{i_1} + \dots + \lambda_{i_k}) + h^2(\lambda_{i_1}\lambda_{i_2} + \dots + \lambda_{i_{k-1}}\lambda_{i_k}) + \dots + h^k(\lambda_{i_1}\lambda_{i_2} \dots \lambda_{i_k})$$

with eigenvectors $v^{i_1} \wedge v^{i_2} \wedge \dots \wedge v^{i_k}$.

Bibliography

1. P Constantin, C Foias, *Global Lyapunov exponents, P Constantin, C Foias, Global Lyapunov exponents, Kaplan-Yorke formulas and the dimension of the attractors for 2D Navier-Stokes equations, Comm Pur Appl Math* **38** (1985), 1-27.
2. P Constantin, C Foias & R Temam, *Attractors representing turbulent flows, AMS Memoirs*, Vol **53** (1985), No314
3. WB Demidowitsch, *Eine verallgemeinerung des kriterium von Bendixson, ZAngewMathMech*, **46**(2)(1996), 145-146.
4. M Fiedler, *Additive compound matrices and inequality for eigenvalues of stochastic matrices, CzechMathJ*, **99**(1974), 392-402.
5. MY Li, *Geometrical Studies on the Global Asymptotic Behaviour of Dissipative Dynamical Systems*, University of Alberta PhD thesis, 1993.
6. MY Li, *Bendixson's criterion for autonomous systems with an invariant linear subspace, RockyMountainJMath* **25**(1995), 351-363.
7. MY Li, *Dulac criteria for autonomous systems having an invariant affine manifold, JMathAnalApplications* **199**(1996), 374-190.

8. MY Li, *Bendixson's criterion for autonomous systems with invariant linear subspaces*, RockyMountainJMath **25**(1995), 351-363.
9. MY Li, *Dulac criteria for autonomous systems having an invariant affine manifold*, JMathAnalAppl **199**(1996), 374-390.
10. MY Li and JS Muldowney, *On Bendixson's criterion*, JDiffEquations **106**(1994), 27-39.
11. MY Li and JS Muldowney, *Lower bounds for the Hausdorff dimension of attractors*, JDynamics&DifferentialEqns **7**(1995), 457-469.
12. MY Li and JS Muldowney, *On RA Smith's autonomous convergence theorem*, RockyMountainJMath **25**(1995), 365-379.
13. MY Li and JS Muldowney, *Poincaré's stability condition for quasi-periodic orbits*, CanadianAppliedMathQuarterly **6**(1998), 367-381.
14. MY Li and JS Muldowney, *Dynamics of differential equations on invariant manifolds*, JDifferentialEquations **168**(2000), 295-320.

15. MY Li and L Wang, *A criterion for stability of matrices*, JMathAnalApplications **225**(1998), 249-264.
16. D London, *On derivations arising in differential equations*, Linear and Multilinear Algebra **4** (1976), 179-189.
17. CC McCluskey, *Bendixson Criteria for Difference Equations*. University of Alberta MSc thesis, 1996
18. CC McCluskey, *Global stability in epidemiological models* University of Alberta PhD thesis, 2002.
19. CC McCluskey and JS Muldowney, *Stability implications of Bendixson's criterion*, SIAM Review **40**(1998), 931-934.
20. CC McCluskey and JS. Muldowney, *Bendixson-Dulac Criteria for Difference Equations*. Journal of Dynamics and Differential Equations. **10** (1998), 567-575.
21. C. C. McCluskey and J. S. Muldowney. *Stability implications of Bendixson's conditions for difference equations*. In B. Aulbach, S. Elaydi, and G. Ladas, editors, *New Progress in Difference Equations*, pages 181-188, 2004.

22. JS Muldowney, *On the dimension of the zero or infinity tending sets for linear differential equations*, Proc AMS **83** (1981), 705-709.
23. JS Muldowney, *Dichotomies and asymptotic behaviour for linear differential systems*, Trans AMS **283** (1984), 465-484.
24. JS Muldowney, *Compound matrices and ordinary differential equations*, RockyMountainJMath **20**(1990), 857-872.
25. B Schwarz, *Totally positive differential systems*, Pacific J Math **32**(1970), 203-229
26. RA Smith, *Some applications of Hausdorff dimension inequalities for ordinary differential equations*, ProcRoySocEdinburgh **104A** (1986), 235-259.
27. R Temam, *Infinite-Dimensional Dynamical Systems in Mechanics and Physics*, Applied Mathematical Sciences, vol 68, Springer-Verlag, New York, 1988.
28. H Wielandt, *Topics in the analytic theory of matrices*, Lecture Notes prepared by RR Meyer, University of Wisconsin, Madison, 1967
29. B Wards, *Dynamics of differential equations on invariant manifolds*, University of Alberta MSc thesis, 2005.

30. Q Wang, *Compound operators and infinite dimensional dynamical systems*, University of Alberta PhD thesis, 2008.
31. E Samuylova, *On the dimension of stable solution subspaces of differential equations*, University of Alberta MSc thesis, 2009.