

HOFFMANN'S CONJECTURE FOR TOTALLY SINGULAR FORMS OF PRIME DEGREE

STEPHEN SCULLY

ABSTRACT. One of the most significant discrete invariants of a quadratic form ϕ over a field k is its *(full) splitting pattern*, a finite sequence of integers which describes the possible isotropy behaviour of ϕ under scalar extension to *arbitrary* overfields of k . A similarly important, but more accessible variant of this notion is that of the *Knebusch splitting pattern* of ϕ , which captures the isotropy behaviour of ϕ as one passes over a certain *prescribed tower* of k -overfields. In this paper, we determine all possible values of this latter invariant in the case where ϕ is *totally singular*. This includes an extension of Karpenko's theorem (formerly Hoffmann's conjecture) on the possible values of the first Witt index to the totally singular case. Contrary to the existing approaches to this problem (in the nonsingular case), our results are achieved by means of a new structural result on the higher anisotropic kernels of totally singular quadratic forms. Moreover, the methods used here readily generalise to give analogous results for arbitrary Fermat-type forms of degree p over fields of characteristic $p > 0$.

1. INTRODUCTION

Let k be a field, let ϕ be a nonzero quadratic form on a (nonzero) k -vector space V of finite dimension and let $X_\phi \subseteq \mathbb{P}(V)$ denote the projective k -scheme defined by the vanishing of ϕ . Given a k -linear subspace W of V , we write $\phi|_W$ for the form obtained by restricting ϕ to W . In the case where $\phi|_W$ is the zero form, W is said to be *totally isotropic* (with respect to ϕ). The largest integer among the dimensions of all totally isotropic subspaces of V is called the *isotropy index* of ϕ , and is denoted by $i_0(\phi)$. In the special case where ϕ is *nonsingular* (i.e., where X_ϕ is smooth), $i_0(\phi)$ is more commonly known as the *Witt index* of ϕ , and is bounded from above by the integer part of $\dim \phi/2$ (where $\dim \phi$ denotes the dimension of the k -vector space V). In the opposite extreme where ϕ is *totally singular* (i.e., where X_ϕ has no smooth points at all), $i_0(\phi)$ may take any value between 0 and $\dim \phi - 1$.

Assume now that ϕ is *anisotropic* (i.e., that $i_0(\phi) = 0$). A simple, yet fundamentally important invariant of ϕ is its *(full) splitting pattern*, which may be defined as the increasing sequence of nonzero isotropy indices attained by ϕ under scalar extension to *every* overfield of k .¹ Although this sequence appears to be somewhat intractable in general, its first entry (assuming there is one) may be computed more explicitly as the isotropy index of the extension ϕ_L of ϕ to the function field $L = k(X_\phi)$ of the (integral) quadric X_ϕ . This (almost tautological) observation is the basic motivation underlying the following construction, originally due to Knebusch (cf. [Kne76]): Let $k_0 = k$, $\phi_0 = \phi$, and inductively define

2010 *Mathematics Subject Classification.* 11E04, 14E05, 15A03.

Key words and phrases. Quadratic forms, quasilinear p -forms, splitting patterns, canonical dimension.

¹This terminology is not standard – see, e.g., [HL04], where refined splitting pattern invariants are considered. Our definition does, however, agree (in content if not presentation) with those found in the literature when ϕ is nonsingular or totally singular (the only relevant cases here).

$k_r = k_{r-1}(X_{\phi_{r-1}})$, $\phi_r = (\phi_{k_r})_{\text{an}}$,² with the understanding that this (finite) process stops when we reach the first nonnegative integer $h(\phi)$ such that $\dim \phi_{h(\phi)} \leq 1$. The integer $h(\phi)$ and the tower of fields $k = k_0 \subset k_1 \subset \dots \subset k_{h(\phi)}$ are known as the *height* and *Knebusch splitting tower* of ϕ , respectively. For $1 \leq r \leq h(\phi)$, the anisotropic form ϕ_r is called the *r-th higher anisotropic kernel* of ϕ . The *r-th higher isotropy index* of ϕ , denoted $i_r(\phi)$, is defined as the difference $i_0(\phi_{k_r}) - i_0(\phi_{k_{r-1}})$. The sequence $\mathbf{i}(\phi) = (i_1(\phi), \dots, i_{h(\phi)}(\phi))$ is called the *Knebusch splitting pattern* of ϕ .³ Note that we have $i_r(\phi) = i_1(\phi_{r-1})$ for every $r \geq 2$ by the inductive nature of the construction.

If ϕ is nonsingular, then its full and Knebusch splitting patterns are easily seen to determine one another (cf. [EKM08, Prop. 25.1]). This need not be the case for (totally) singular forms (see Example 2.47 below), but Knebusch's construction still offers a meaningful and practical way to pre-classify quadratic forms according to some notion of "algebraic complexity". By virtue of its definition, the Knebusch splitting pattern thus embodies a fundamental link between intrinsic algebraic properties of quadratic forms and the geometry of the algebraic varieties which are naturally associated to them. In recent years, the advent of effective new tools with which to study algebraic cycles on projective homogeneous varieties has, in this way, led to dramatic progress on many long-standing problems within the algebraic theory of quadratic forms. The impact of these developments has been felt most deeply in characteristic $\neq 2$, where (1) anisotropic forms of dimension ≥ 2 are necessarily nonsingular, and (2) the geometric methods are better developed, even if we restrict our considerations to nonsingular forms only – see [EKM08] for a thorough exposition of much of the recent work which has been done in this area.

One of the central problems in the investigation of splitting properties of quadratic forms over general fields is the following:

Question 1.1. Let ϕ be an anisotropic quadratic form of dimension ≥ 2 over a field. What are the possible values of the sequence $\mathbf{i}(\phi)$?

Since $i_r(\phi) = i_1(\phi_{r-1})$ for all $2 \leq r \leq h(\phi)$, the natural first approximation to this problem is to determine the possible values of the invariant i_1 among all forms of a given dimension. To this end, we have the following general conjecture:

Conjecture 1.2 (Hoffmann⁴). *Let ϕ be an anisotropic quadratic form of dimension ≥ 2 over a field. Then $i_1(\phi) - 1$ is the remainder modulo 2^s of $\dim \phi - 1$ for some $s < \log_2(\dim \phi)$.*⁵

In characteristic $\neq 2$, the first major result in the direction of Conjecture 1.2 was established by Hoffmann himself, who showed that if $\dim \phi = 2^n + m$ for nonnegative integers n and $1 \leq m \leq 2^n$, then $i_1(\phi) \leq m$ ([Hof95, Cor. 1]). A few years later, Izhboldin ([Izh04, Cor. 5.12]) proved that one cannot have $i_1(\phi) = m - 1$ here unless $m = 2$. Izhboldin's paper combined an elaboration of the algebraic methods conceived in [Hof95] with emerging work of Vishik ([Vis98],[Vis99]), who developed a systematic approach to the study of the splitting pattern using the (integral) motive of the given

²For any form ψ over a field K , ψ_{an} denotes the *anisotropic kernel* of ψ , an anisotropic K -form uniquely determined up to isomorphism by the (refined) Witt decomposition of ψ – see [HL04, §2].

³The term *standard splitting pattern* is also used in the literature. Footnote 1 again applies here.

⁴This conjecture was originally stated by Hoffmann under the additional hypothesis that $\text{char}(k) \neq 2$, but we expect that the assertion is also valid in characteristic 2.

⁵After passing to a purely transcendental extension of k if necessary, all values of $i_1(\phi)$ which are not excluded by the conjecture can be realised by making an appropriate choice of ϕ . In all cases, ϕ may, in fact, be chosen to be either nonsingular or (if $\text{char}(k) = 2$) totally singular – see [Vis04, §7.4] and Proposition 6.4 below, respectively.

quadric (see [Vis04], where motivic methods were used to verify Hoffmann's conjecture in all dimensions ≤ 22 in this setting). Going further, Vishik later formulated a very general conjecture concerning the complete motivic decomposition a smooth anisotropic quadric (the *Excellent Connections* conjecture) which subsumed the nonsingular case of Conjecture 1.2 in a conceptual way. Not long after this, Karpenko ([Kar03]) used a similar approach to prove the characteristic $\neq 2$ case of the conjecture in its entirety, the key new ingredient being the use of (reduced power) Steenrod operations on modulo-2 Chow groups. More recently, Vishik ([Vis11, Thm. 1.3]) proved the Excellent Connections conjecture in characteristic $\neq 2$, thus yielding another proof of Hoffmann's conjecture in this setting. This work also makes essential use of Steenrod squares in Chow theory.

In characteristic 2, the general picture is more complicated. In the *nonsingular* case, the Karpenko-Vishik approach to Conjecture 1.2 is still valid, and progress is only hindered here by the fact that the total mod-2 Steenrod operation is not yet available when 2 is not invertible in the base field. To this end, weak forms of the first three Steenrod squares have been constructed by Houton, and these suffice to prove nontrivial partial results towards Conjecture 1.2 (see [Hau13, Thm. 6.2],[Hau15, Thm. 5.8]). Houton's results are further supplemented by earlier work of Hoffmann and Laghribi, who extended Hoffmann's upper bound on $i_1(\phi)$ to the characteristic-2 setting, irrespective of whether ϕ is nonsingular or not ([HL06, Lem. 4.1]). For *singular* forms, however, the situation is different, and almost nothing is known in the direction of Conjecture 1.2 beyond Hoffmann and Laghribi's bound. In fact, the only real exception to this general state of affairs lies in the extreme case where ϕ is *totally singular*. Here, it was recently shown in [Scu14, Thm. 9.4] that if $\dim \phi = 2^n + m$ for nonnegative integers n and $1 \leq m \leq 2^n$, and if $i_1(\phi) \neq m$ (that is, if Hoffmann and Laghribi's bound is *not* met), then $i_1(\phi) \leq \frac{m}{2}$. In the present article, we will settle this case completely by proving:

Theorem 1.3. *Conjecture 1.2 is true in the case where ϕ is totally singular.*

Contrary to the existing approaches to the nonsingular case of Conjecture 1.2, our proof of Theorem 1.3 does not involve the study of Chow correspondences on the quadric X_ϕ . Indeed, although we also make use of the computation of the canonical dimension⁶ of X_ϕ (see [KM03, Tot08]), it is exploited here in a rather more direct and algebraic way. This point of view begins with the following observation:

Proposition 1.4 (see Proposition 4.3 below). *Let ϕ be an anisotropic totally singular quadratic form of dimension ≥ 2 over a field k of characteristic 2 and let $\psi \subset \phi$ be a subform of codimension $i_1(\phi)$. Suppose furthermore that $h(\psi) < h(\phi)$.⁷ Then there exist a quasi-Pfister quadratic form π ,⁸ a subform $\sigma \subset \pi$, an element $\lambda \in k^*$ and a form τ over k such that $\psi \simeq \pi \otimes \tau$ and $\phi \simeq \psi \perp \lambda\sigma$.*

In the situation of Proposition 1.4, Hoffmann's conjecture is immediately verified. Indeed, (since $\sigma \subset \pi$) the integer $\dim \pi$ is a power of 2 strictly greater than $i_1(\phi) - 1 = \dim \sigma - 1$, and (since ψ is divisible by π) we have $\dim \phi - 1 = \dim \psi + i_1(\phi) - 1 \equiv i_1(\phi) - 1 \pmod{\dim \pi}$. It is not always possible, however, to decompose the form ϕ in the manner intimated by the proposition. In fact, Vishik (see [Tot09, Lem. 7.1]) has given examples of 16-dimensional anisotropic quadratic forms in characteristic $\neq 2$ which have first higher isotropy index equal to 2, but which do not decompose in this way, and the same examples

⁶Here, the *canonical dimension* of an algebraic variety X over a field k should be understood as the minimal dimension $\text{cdim}(X)$ of the image of a rational self-map $X \dashrightarrow X$.

⁷A weaker condition will suffice – see the statement of Proposition 4.3.

⁸That is, π is the diagonal part of a bilinear Pfister form over k – see §2.3 below.

carry over into the totally singular setting (see Lemma 4.4 below). Thus, the picture is, in general, more complicated than that suggested by Proposition 1.4. Perhaps surprisingly, however, the main result of this paper shows that the next best thing happens:

Theorem 1.5. *Let ϕ be an anisotropic totally singular quadratic form of dimension ≥ 2 over a field of characteristic 2 and let s be the smallest nonnegative integer such that $2^s \geq i_1(\phi)$. Then ϕ_1 is divisible by an s -fold quasi-Pfister form.*

This is a new kind of statement of which no analogue is known in the nonsingular theory (even in characteristic $\neq 2$). As remarked above, a key ingredient needed for its proof is Totaro's computation of the canonical dimension of a totally singular quadric (see [Tot08, Thm. 5.1]). In [Scu13], this computation was extended to the wider class of Fermat-type hypersurfaces of degree p over fields of characteristic $p > 0$, and this enables us to also prove a direct analogue of Theorem 1.5 for totally singular forms of any prime degree $p > 2$ (known here as *quasilinear p -forms*) - see Theorem 5.1 below. Subsequently, we also get an analogue of Theorem 1.3 in higher degrees. As a corollary, this yields a complete solution to the problem of determining the possible values of the canonical dimension of a degree- p Fermat-type hypersurface in characteristic $p > 0$ (Theorem 6.6); it is worth noting here that no such result is known for Fermat-type hypersurfaces of prime degree $p > 2$ over fields of characteristic *not* p . Returning to the case where $p = 2$, let us explain more precisely how Theorem 1.5 implies the totally singular case of Hoffmann's conjecture:

Proof of Theorem 1.3. Since ϕ is totally singular, we have $\dim \phi_1 = \dim \phi - i_1(\phi)$ (see Remark 2.36 (2) below). Thus, if s is as in Theorem 1.5, then

$$\dim \phi - 1 = \dim \phi_1 + i_1(\phi) - 1 \equiv i_1(\phi) - 1 \pmod{2^s}.$$

Since $i_1(\phi) - 1 < 2^s$, the result follows. \square

Of course, our main result goes somewhat deeper than this. In fact, Theorem 1.5 (resp. Theorem 5.1 below) yields a complete answer to Question 1.1 in the totally singular case (resp. its analogue for arbitrary quasilinear p -forms). In other words, all restrictions on the possible values of the Knebusch splitting pattern of ϕ are explained here by the presence of certain divisibilities among its higher anisotropic kernels - precise statements are given in §6.1 below (see Theorem 6.1, Proposition 6.4). With this result in hand, it then becomes natural to try to understand the general discrepancy which exists between the Knebusch and *full* splitting patterns in the totally singular case. An immediate challenge here concerns the determination of all non-trivial restrictions which the former invariant imposes on the latter. In §7 below, we initiate this process by conjecturing that the gaps in the full splitting pattern established in characteristic $\neq 2$ by Vishik as a consequence of his proof of the Excellent Connections conjecture (see [Vis11, Prop. 2.6] or Theorem 7.1 below) are also present in the totally singular theory. A particular case of this conjecture was already proved in [Scu14, Thm. 9.2], and we provide some further evidence for its general veracity here.⁹

Finally, the main results of this paper should have valuable implications for the study of symmetric bilinear forms over fields of characteristic 2. Indeed, in characteristic 2, the diagonal parts of symmetric bilinear forms are nothing else but the totally singular quadratic forms discussed above. This will be investigated in a later text.

The remainder of this text is organised as follows: In sections 2 and 3, we recall the basic theory of quasilinear p -forms and introduce the key notions and results which will

⁹To the author's knowledge, there is no conjectural description of the possible values of the full splitting pattern, even in the nonsingular case (in any characteristic).

be needed in the main part of the text. As a warm-up for the proof of our main result, we prove in §4 (a stronger version of) Proposition 1.4 above and consider some situations in which it may be applied. The proof of Theorem 1.5 (and its generalisation to higher degrees) is then given in §5, and, in §6, we apply this result to determine all possible Knebusch splitting patterns of quasilinear p -forms and settle another conjecture of Hoffmann concerning quasilinear p -forms with “maximal splitting”. Lastly, in §7, we consider the aforementioned problem of establishing totally singular analogues of the results obtained by Vishik in [Vis11].

Notation and Terminology. Unless stated otherwise, p will denote an arbitrary prime integer and F will denote an arbitrary field of characteristic p . If L is a field of characteristic p and a_1, \dots, a_n are elements of L , then L_{a_1, \dots, a_n} will denote the field $L(\sqrt[p]{a_1}, \dots, \sqrt[p]{a_n})$. Finally, if k is a field and $T = (T_1, \dots, T_m)$ is a tuple of algebraically independent variables over k , then we will write $k[T]$ for the polynomial ring $k[T_1, \dots, T_m]$ and $k(T)$ for its fraction field.

2. QUASILINEAR p -FORMS AND QUASILINEAR p -HYPERSURFACES

The basic material presented in this section was originally developed in a series of papers by Hoffmann and Laghribi (see [HL04, Lag04a, Lag04b, Lag06] and especially [Hof04]). Additional elementary results which will be needed in the sequel are also included here. For any details which are omitted from our exposition of the basic theory, we refer the reader to [Hof04].

2.1. Basic notions. Let $\phi: V \rightarrow F$ be an F -valued form on a finite-dimensional F -vector space V . We say that ϕ is a *quasilinear p -form* (on V) if ϕ is homogeneous of degree p and the equation $\phi(v + w) = \phi(v) + \phi(w)$ holds for all $(v, w) \in V \times V$. By a *quasilinear p -form over F* (or sometimes simply a *form over F* or *F -form*), we will mean a quasilinear p -form on some finite-dimensional F -vector space. By a *quasilinear p -hypersurface over F* we will mean a projective hypersurface defined by the vanishing of a nonzero quasilinear p -form on some F -vector space of dimension ≥ 2 . In the special case where $p = 2$, we will speak of *quasilinear quadratic forms* rather than quasilinear 2-forms.

Remark 2.1. Let $T = (T_1, \dots, T_n)$ be a tuple of $n \geq 2$ algebraically independent variables over F , and let $f \in F[T] \setminus \{0\}$. Then the projective hypersurface $X_f = \{f = 0\} \subset \mathbb{P}^{n-1}$ is nowhere smooth if and only if $f \in F[T_1^p, \dots, T_n^p]$. In particular, if $p = 2$, then a nonzero quadratic form of dimension ≥ 2 over F is totally singular (in the sense of §1) if and only if it is quasilinear.

Let ϕ be a quasilinear p -form over F . The underlying F -vector space of ϕ will be denoted by V_ϕ . Its dimension will be called the *dimension* of ϕ and will be denoted by $\dim \phi$. If $\dim \phi \geq 2$ and ϕ is nonzero, then the quasilinear p -hypersurface $\{\phi = 0\} \subset \mathbb{P}(V_\phi)$ (which is nowhere smooth by Remark 2.1) will be denoted by X_ϕ . The set $\{\phi(v) \mid v \in V_\phi\}$ of elements of F represented by ϕ will be denoted by $D(\phi)$. Given a field extension L of F , we will write ϕ_L for the unique quasilinear p -form on the L -vector space $V_\phi \otimes_F L$ such that $\phi_L(v \otimes 1) = \phi(v)$ for all $v \in V_\phi$. If R is a subring of L containing F , then $D(\phi_R)$ will denote the subset $\{\phi(w) \mid w \in V_\phi \otimes_F R\}$ of $D(\phi_L)$ (which lies in R). Given $a \in F$, we will write $a\phi$ for the form $v \mapsto a\phi(v)$ on the vector space V_ϕ .

Let ψ be another quasilinear p -form over F . If there exists an injective (resp. bijective) F -linear map $f: V_\psi \rightarrow V_\phi$ such that $\phi(f(v)) = \psi(v)$ for all $v \in V_\psi$, then we will say that ψ is a *subform* of (resp. is *isomorphic* to) ϕ and write $\psi \subset \phi$ (resp. $\psi \simeq \phi$). If $\psi \simeq a\phi$ for

some $a \in F^*$, then we will say that ψ and ϕ are *similar*. The *sum* $\psi \oplus \phi$ (resp. *product* $\psi \otimes \phi$) is defined as the unique quasilinear p -form on $V_\psi \oplus V_\phi$ (resp. $V_\psi \otimes_F V_\phi$) such that $(\psi \oplus \phi)((v, w)) = \psi(v) + \phi(w)$ (resp. $(\psi \otimes \phi)(v \otimes w) = \psi(v)\phi(w)$) for all $(v, w) \in V_\psi \times V_\phi$. Given a positive integer n , $n \cdot \phi$ will denote the sum of n copies of ϕ (note that we have $n \cdot \phi \neq n\phi$ for $n > 1$). If there exists a form τ over F such that $\phi \simeq \psi \otimes \tau$, then we will say that ϕ is *divisible* by ψ .

Given elements $a_1, \dots, a_n \in F$, we will write $\langle a_1, \dots, a_n \rangle$ for the quasilinear p -form $(\lambda_1, \dots, \lambda_n) \mapsto \sum_{i=1}^n a_i \lambda_i^p$ on the F -vector space $F^{\oplus n}$. By definition, every quasilinear p -form of dimension n over F is isomorphic to $\langle a_1, \dots, a_n \rangle$ for some $a_i \in F$.

A vector $v \in V_\phi$ is said to be *isotropic* if $\phi(v) = 0$. We will say that ϕ is *isotropic* if V_ϕ contains a nonzero isotropic vector, and *anisotropic* otherwise. By the additivity of ϕ , the subset V_ϕ^0 of all isotropic vectors in V_ϕ is, in fact, an F -linear subspace of V_ϕ . Its dimension will be called the *isotropy index* of ϕ , and will be denoted by $\mathfrak{i}_0(\phi)$ (note that in the case where $p = 2$ and ϕ is nonzero, this agrees with the definition given in §1). The additivity of ϕ also implies that $D(\phi)$ is a finite-dimensional F^p -linear subspace of F (where F is equipped with its natural F^p -vector space structure). Conversely, if U is a nonzero finite-dimensional F^p -linear subspace of F , and if a_1, \dots, a_n is a basis of U , then $\sigma = \langle a_1, \dots, a_n \rangle$ is a quasilinear p -form over F satisfying $D(\sigma) = U$. In fact, it is easy to see that, up to isomorphism, σ is the unique *anisotropic* quasilinear p -form with this property:

Lemma 2.2 (cf. [Hof04, Prop. 2.12]). *Let U be a finite-dimensional F^p -linear subspace of F . Then, up to isomorphism, there exists a unique anisotropic quasilinear p -form ϕ over F such that $D(\phi) = U$.*

In particular, Pfister's quadratic subform theorem ([EKM08, Thm. 17.12]) takes the following simplified form in this setting:

Proposition 2.3 (cf. [Hof04, Prop. 2.6]). *Let ψ and ϕ be anisotropic quasilinear p -forms over F . Then $\psi \subset \phi$ if and only if $D(\psi) \subseteq D(\phi)$. In particular, $\psi \simeq \phi$ if and only if $D(\psi) = D(\phi)$.*

In view of these observations, we can define (up to isomorphism) the *anisotropic kernel* of ϕ as the unique anisotropic quasilinear p -form ϕ_{an} over F such that $D(\phi_{\text{an}}) = D(\phi)$. If we view $D(\phi)$ as an F -vector space via the Frobenius $F \mapsto F^p$, then $\phi: V_\phi \rightarrow D(\phi)$ is a surjective F -linear map with kernel V_ϕ^0 . We thus obtain:

Proposition 2.4 (cf. [Hof04, Lem 2.10]). *Let ϕ be a quasilinear p -form over F . Then ϕ is anisotropic if and only if $\phi \simeq \phi_{\text{an}}$. If ϕ is isotropic, then $\phi \simeq \phi_{\text{an}} \oplus \mathfrak{i}_0(\phi) \cdot \langle 0 \rangle$. In particular, $\dim \phi_{\text{an}} = \dim \phi - \mathfrak{i}_0(\phi)$.*

In summary, we see that ϕ is determined up to isomorphism by the set $D(\phi)$ and the integer $\mathfrak{i}_0(\phi)$. If $\dim \phi_{\text{an}} \leq 1$, then we will say that ϕ is *split*. Given another form ψ over F , we will write $\psi \sim \phi$ whenever $\psi_{\text{an}} \simeq \phi_{\text{an}}$. If F is perfect (that is, if $F = F^p$), then every form over F is split and the theory is vacuous. As such, we are essentially only interested in the case where F is imperfect. Unless indicated otherwise, we will assume henceforth that all quasilinear p -forms are *nonzero* (i.e., of anisotropic dimension ≥ 1).

Remark 2.5. Let ϕ be a quasilinear p -form over F . Choose a basis v_1, \dots, v_n of V_ϕ and let $a_i = \phi(v_i)$ for all $1 \leq i \leq n$, so that $\phi \simeq \langle a_1, \dots, a_n \rangle$. Suppose that $v \in V_\phi$ is an isotropic vector, and write $v = \sum_{i=1}^n \lambda_i v_i$. If $\lambda_j \neq 0$ for $1 \leq j \leq n$, then the F^p -vector space $D(\phi)$ is spanned by the elements $a_1, \dots, a_{j-1}, a_{j+1}, \dots, a_n$. In particular, we have $\phi \sim \langle a_1, \dots, a_{j-1}, a_{j+1}, \dots, a_n \rangle$.

2.2. Function fields of quasilinear p -hypersurfaces and their products. Let ϕ be a quasilinear p -form of dimension ≥ 2 over F . If ϕ is not split, then the quasilinear p -hypersurface X_ϕ is an integral scheme (cf. [Hof04, Lem. 7.1]), as is its affine cone $\{\phi = 0\} \subset \mathbb{A}(V_\phi)$. In this case, we will write $F(\phi)$ for the function field of the former and $F[\phi]$ for that of the latter. If L is a field extension of F , then we will simply write $L(\phi)$ instead of $L(\phi_L)$ whenever it is defined. In general, given a finite collection ϕ_1, \dots, ϕ_n of quasilinear p -forms of dimension ≥ 2 over F , we will write $F(\phi_1 \times \dots \times \phi_n)$ for the function field of the scheme $X_{\phi_1} \times \dots \times X_{\phi_n}$, provided that it is integral. This notation will be further simplified where possible; for example, if $\phi_1 = \dots = \phi_n = \phi$, then we will simply write $F(\phi^{\times n})$ instead of $F(\phi_1 \times \dots \times \phi_n)$.

Remarks 2.6. Let ϕ be a quasilinear p -form over F . Assume that ϕ is not split. We make the following basic observations:

- (1) Let $a_0, \dots, a_n \in F$ be such that $\phi \simeq \langle a_0, \dots, a_n \rangle$ and $a_0, a_1 \neq 0$. Then we have F -isomorphisms

$$F(\phi) \simeq F(S) \left(\sqrt[p]{a_1^{-1}(a_0 + a_2 S_2^p + \dots + a_n S_n^p)} \right)$$

and

$$F[\phi] \simeq \text{Frac}(F[T]/(a_0 T_0^p + \dots + a_n T_n^p)) \simeq F(U) \left(\sqrt[p]{a_0^{-1}(a_1 U_1^p + \dots + a_n U_n^p)} \right),$$

where $S = (S_2, \dots, S_n)$, $T = (T_0, \dots, T_n)$ and $U = (U_1, \dots, U_n)$ are tuples of algebraically independent variables over F .

- (2) $F[\phi]$ is F -isomorphic to a degree-1 purely transcendental extension of $F(\phi)$.
 (3) $F(\phi)$ is F -isomorphic to a degree- $i_0(\phi)$ purely transcendental extension of $F(\phi_{\text{an}})$ – see Proposition 2.4.
 (4) The form $\phi_{F(\phi)}$ is evidently isotropic. Furthermore, if $a_1, \dots, a_n \in F$ are such that $\phi \simeq \langle a_1, \dots, a_n \rangle$, then consideration of the generic point in $X_\phi(F(\phi))$ shows that $\phi_{F(\phi)} \sim \langle a_1, \dots, a_{i-1}, a_{i+1}, \dots, a_n \rangle$ for every $1 \leq i \leq n$ – see Remark 2.5.

2.3. Quasi-Pfister p -forms. Let ϕ be a quasilinear p -form over F and let n be a positive integer. We say that ϕ is an n -fold *quasi-Pfister p -form* if there exist $a_1, \dots, a_n \in F$ such that $\phi \simeq \langle \langle a_1, \dots, a_n \rangle \rangle := \otimes_{i=1}^n \langle 1, a_i, a_i^2, \dots, a_i^{p-1} \rangle$. For convenience, we also say that ϕ is a 0 -fold *quasi-Pfister p -form* if $\phi \simeq \langle 1 \rangle$. Note, in particular, that if ϕ is an n -fold quasi-Pfister p -form for some $n \geq 0$, then we have $\dim \phi = p^n$. The basic observation concerning quasi-Pfister p -forms is found in the following proposition (which follows easily from Lemma 2.2):

Proposition 2.7 (cf. [Hof04, Prop. 4.6]). *Let ϕ be a quasilinear p -form over F . Then ϕ_{an} is a quasi-Pfister p -form if and only if $D(\phi)$ is a subfield of F .*

Since the set of elements represented by an arbitrary quasi-Pfister p -form is, by definition, a subfield of the base field, we obtain:

Corollary 2.8 (cf. [Hof04, Prop. 4.6]). *Let ϕ be a quasi-Pfister p -form over F . Then ϕ_{an} is a quasi-Pfister p -form. In particular, if ϕ is isotropic, then $\dim \phi_{\text{an}} = \frac{1}{p^k} \dim \phi$ for some $k \geq 1$.*

Remark 2.9. More explicitly, let $\phi = \langle \langle a_1, \dots, a_n \rangle \rangle$ for some $n \geq 1$ and $a_i \in F$. Then $D(\phi) = F^p(a_1, \dots, a_n)$. Let m be such that $[F^p(a_1, \dots, a_n) : F^p] = p^m$. If $m = 0$ (i.e., $D(\phi) = F^p$), then $\phi_{\text{an}} \simeq \langle 1 \rangle$. If $m \geq 1$, then $\phi_{\text{an}} \simeq \langle \langle b_1, \dots, b_m \rangle \rangle$ for any m elements $b_1, \dots, b_m \in F$ such that $F^p(b_1, \dots, b_m) = F^p(a_1, \dots, a_n)$.

Quasi-Pfister p -forms have a central role to play in the general theory of quasilinear p -forms. As shown by Hoffmann ([Hof04]), these forms are distinguished here by the very same properties which distinguish the classical Pfister forms among nonsingular quadratic forms. For this reason, it will be useful to define the *divisibility index* of a given form ϕ , denoted $\mathfrak{d}_0(\phi)$, as the largest nonnegative integer s such that ϕ_{an} is divisible by an s -fold quasi-Pfister p -form. Clearly we have $\mathfrak{d}_0(\phi) \leq \log_p(\dim \phi_{\text{an}})$, with equality holding if and only if ϕ_{an} is similar to a quasi-Pfister p -form. An alternative description of this invariant will be given in Corollary 2.19 below.

2.4. The norm form. Let ϕ be a quasilinear p -form over F . The *norm field* of ϕ , denoted $N(\phi)$, is defined (cf. [Hof04, Def. 4.1]) as the smallest subfield of F which contains all ratios of nonzero elements of $D(\phi)$. Note, in particular, that we have $N(a\phi) = N(\phi) = N(\phi_{\text{an}})$ for all $a \in F^*$. In spite of its simple nature, this invariant has an important role to play in the whole theory. A more explicit description of the norm field may be given as follows:

Remark 2.10. If $a_1, \dots, a_n \in F$ are such that $\phi \simeq \langle a_1, \dots, a_n \rangle$ and $a_1 \neq 0$, then we have $N(\phi) = F^p(\frac{a_2}{a_1}, \dots, \frac{a_n}{a_1})$.

In particular, we see that $N(\phi)$ is a nonzero finite-dimensional F^p -linear subspace of F . By Lemma 2.2, it follows that, up to isomorphism, there exists a unique anisotropic quasilinear p -form ϕ_{nor} over F such that $D(\phi_{\text{nor}}) = N(\phi)$. The form ϕ_{nor} is called the *norm form* of ϕ (cf. [Hof04, Def. 4.9]). By Proposition 2.7, ϕ_{nor} is a quasi-Pfister p -form. Its dimension (which is necessarily equal to a power of p) is called the *norm degree* of ϕ , and is denoted by $\text{ndeg}(\phi)$ (cf. [Hof04, Def. 4.1]). The following lemma characterises the norm form as the smallest anisotropic quasi-Pfister p -form which contains ϕ_{an} as a subform up to multiplication by a scalar (again, this is a simple consequence of Proposition 2.3):

Lemma 2.11 (cf. [Scu14, Lem. 2.10]). *Let ϕ be a quasilinear p -form (resp. a quasilinear p -form such that $1 \in D(\phi)$) and π an anisotropic quasi-Pfister p -form over F . Then ϕ_{an} is similar to a subform of π (resp. $\phi_{\text{an}} \subset \pi$) if and only if $\phi_{\text{nor}} \subset \pi$. In particular, ϕ_{an} is similar to a subform of ϕ_{nor} (resp. $\phi_{\text{an}} \subset \phi_{\text{nor}}$).*

Example 2.12. Let ϕ be a quasilinear p -form over F . The following are equivalent:

- (1) ϕ_{an} is similar (resp. isomorphic) to a quasi-Pfister p -form.
- (2) $\phi_{\text{nor}} \simeq a\phi_{\text{an}}$ for some $a \in F^*$ (resp. $\phi_{\text{nor}} \simeq \phi_{\text{an}}$).
- (3) $N(\phi) = aD(\phi)$ for some $a \in F^*$ (resp. $N(\phi) = D(\phi)$).

2.5. Similarity factors. Let ϕ be a quasilinear p -form over F . By a *similarity factor* of ϕ , we mean an element $a \in F^*$ such that $a\phi \simeq \phi$. The set of all similarity factors of ϕ will be denoted by $G(\phi)^*$, and we will write $G(\phi)$ for the set $G(\phi)^* \cup \{0\}$. Note that $G(a\phi) = G(\phi) = G(\phi_{\text{an}})$ for all $a \in F^*$, the second equality being an obvious consequence of Proposition 2.4. Thus, in view of Proposition 2.3, we have:

Lemma 2.13 (cf. [Hof04, Lem. 6.3]). *Let ϕ be a quasilinear p -form over F and let $a \in F^*$. Then $a \in G(\phi)^*$ if and only if $aD(\phi) \subseteq D(\phi)$.*

Example 2.14. Let ϕ be a quasi-Pfister p -form over F . Then, since $D(\phi)$ is a subfield of F , we have $G(\phi) = D(\phi)$.

More generally, Lemma 2.13 immediately implies the following:

Corollary 2.15 (cf. [Hof04, Prop. 6.4]). *Let ϕ be a quasilinear p -form over F . Then $G(\phi)$ is a subfield of $N(\phi)$ containing F^p .*

In particular, $G(\phi)$ is a nonzero finite-dimensional F^p -linear subspace of F . By Lemma 2.2, it follows that, up to isomorphism, there exists a unique anisotropic quasilinear p -form ϕ_{sim} over F such that $D(\phi_{\text{sim}}) = G(\phi)$. The form ϕ_{sim} is called the *similarity form* of ϕ (cf. [Hof04, Def. 6.5]). By Proposition 2.7, ϕ_{sim} is a quasi-Pfister p -form. Taken together, Examples 2.12 and 2.14 yield:

Example 2.16. Let ϕ be a quasilinear p -form over F . The following are equivalent:

- (1) ϕ_{an} is similar (resp. isomorphic) to a quasi-Pfister p -form.
- (2) $\phi_{\text{sim}} \simeq \phi_{\text{nor}} \simeq a\phi_{\text{an}}$ for some $a \in F^*$ (resp. $\phi_{\text{sim}} \simeq \phi_{\text{nor}} \simeq \phi_{\text{an}}$).
- (3) $G(\phi) = N(\phi) = aD(\phi)$ for some $a \in F^*$ (resp. $G(\phi) = N(\phi) = D(\phi)$).

The basic observation concerning similarity factors is the following:

Proposition 2.17. *Let ϕ and ψ be quasilinear p -forms over F . Then $G(\psi) \subseteq G(\phi)$ if and only if ϕ_{an} is divisible by ψ_{sim} .*

Proof. We may assume that $1 \in D(\phi)$. Suppose that $G(\psi) \subseteq G(\phi)$. By Corollary 2.15, $G(\psi)$ and $G(\phi)$ are subfields of F . By Lemma 2.13, $D(\phi)$ is naturally a (finite-dimensional) vector space over $G(\phi)$, and hence over $G(\psi)$. If a_1, \dots, a_m is a basis of $D(\phi)$ over $G(\psi)$, then (since $D(\psi_{\text{sim}}) = G(\psi)$), Lemma 2.2 implies that $\phi_{\text{an}} \simeq \psi_{\text{sim}} \otimes \langle a_1, \dots, a_m \rangle$. Conversely, if ϕ_{an} is divisible by ψ_{sim} , then it is clear that $G(\psi) \subseteq G(\phi)$, since $G(\psi) = D(\psi_{\text{sim}}) = G(\psi_{\text{sim}})$ by Example 2.16. \square

We thus obtain the following characterisation of the similarity form:

Corollary 2.18 (cf. [Hof04, Prop. 6.4]). *Let ϕ be a quasilinear p -form and π an anisotropic quasi-Pfister p -form over F . Then ϕ_{an} is divisible by π if and only if ϕ_{sim} is divisible by π . In particular, ϕ_{an} is divisible by ϕ_{sim} .*

This enables us to reinterpret the divisibility index $\mathfrak{d}_0(\phi)$ (see §2.3) as follows:

Corollary 2.19. *Let ϕ be a quasilinear p -form over F . Then $\mathfrak{d}_0(\phi) = \log_p(\dim \phi_{\text{sim}}) = \log_p([G(\phi) : F^p])$.*

We also get the following:

Corollary 2.20. *Let ϕ and ψ be quasilinear p -forms over F . Then ϕ_{an} is divisible by ψ_{nor} if and only if $N(\psi) \subseteq G(\phi)$. If, additionally, $1 \in D(\psi)$, then the latter condition may be replaced by $D(\psi) \subseteq G(\phi)$.*

Proof. For the second statement, we simply recall that $N(\phi)$ is the smallest subfield of F containing all ratios of nonzero elements of $D(\phi)$ and that $G(\phi)$ is a subfield of F (Corollary 2.15). For the first, we can replace ψ by its norm form to arrive at the case where $N(\psi) = G(\psi)$ and $\psi_{\text{nor}} \simeq \psi_{\text{sim}}$ (see Example 2.16). The result is therefore a particular case of Proposition 2.17. \square

2.6. A criterion for a quasilinear p -form to be quasi-Pfister. Let ϕ be a quasilinear p -form over F such that $1 \in D(\phi)$, let L be a field extension of F and let $\alpha \in D(\phi_L) \setminus \{0\}$. Consider the set $S_\alpha = \{a \in F \mid \alpha a \in D(\phi_L)\}$. Since $D(\phi_L)$ is an L^p -linear subspace of L , we have the following observation:

$$(2.1) \quad \sum \lambda_i^p a_i \in S_\alpha \text{ for all } \lambda_i \in F \text{ and all } a_i \in S_\alpha.$$

Lemma 2.21. *In the above situation, let $P \in F^p[T]$ be a polynomial of degree $< p$ in a single variable T such that $P(b) \in S_\alpha$ for all $b \in D(\phi)$. Then $b^n \in S_\alpha$ for all $b \in D(\phi)$ and all $n \leq \deg(P)$.*

Proof. We proceed by induction on $d = \deg(P)$. Since $\alpha \in D(\phi_L)$, the case where $d = 0$ is trivial. Suppose now that $d > 0$, and let $\lambda \in F$ be such that $P(T + \lambda^p) = P(T) + Q(T)$ for some $Q \in F^p[T]$ of degree $d-1$. Since $F^p \subseteq D(\phi)$ by hypothesis, our assumption and (2.1) imply that $Q(b) = P(b + \lambda^p) - P(b) \in S_\alpha$ for all $b \in D(\phi)$. By the induction hypothesis, it follows that $b^n \in S_\alpha$ for all $b \in D(\phi)$ and all $n < d$. Finally, since $P(b) = \sum_{i=0}^d \lambda_i^p b^i$ for some $\lambda_i \in F$ with $\lambda_d \neq 0$, (2.1) implies that, for any $b \in D(\phi)$, we also have $b^d \in S_\alpha$. This proves the lemma. \square

Suppose now that there exists a polynomial $P \in F^p[T]$ as in the statement of Lemma 2.21 with $\deg(P) \geq 2$ (in particular, we necessarily have $p > 2$). A first application of the lemma shows that we have $D(\phi) \subseteq S_\alpha$. Since $D(\phi_L)$ is spanned by $D(\phi)$ as an L^p -vector space, this implies that $\alpha D(\phi_L) \subseteq D(\phi_L)$, and hence (for dimension reasons) that $\alpha D(\phi_L) = D(\phi_L)$. Another application of Lemma 2.21 then shows that $b^n \in D(\phi_L)$ for all $b \in D(\phi)$ and all $n \leq \deg(P)$. In particular, since $\deg(P) \geq 2$, we have $2bc = (b+c)^2 - b^2 - c^2 \in D(\phi_L)$ for all $b, c \in D(\phi)$. Since $p > 2$, and since $D(\phi_L)$ is spanned by $D(\phi)$ as an L^p -vector space, this implies that $D(\phi_L)$ is closed under multiplication, i.e., that $D(\phi_L) = N(\phi_L)$ (see Remark 2.10). By Example 2.12, this means that $(\phi_L)_{\text{an}} \simeq (\phi_L)_{\text{nor}}$. We have thus proved:

Lemma 2.22. *Assume that $p > 2$. Let ϕ be a quasilinear p -form over F such that $1 \in D(\phi)$ and let L be a field extension of F . Suppose that there exists a polynomial $P \in F^p[T]$ in a single variable T , and an element $\alpha \in D(\phi_L) \setminus \{0\}$ such that $2 \leq \deg(P) < p$ and $\alpha P(b) \in D(\phi_L)$ for all $b \in D(\phi)$. Then $(\phi_L)_{\text{an}}$ is a quasi-Pfister p -form.*

2.7. The Cassels-Pfister representation theorem. Let ϕ be a quadratic form over a field k and let $f \in k[T]$ be a polynomial in a single variable T which is represented by the form $\phi_{k(T)}$. One of the foundational results of the classical algebraic theory of quadratic forms is the Cassels-Pfister representation theorem, which asserts that, in this case, ϕ already represents f over the polynomial ring $k[T]$ (see [EKM08, Thm. 17.3]). In the present setting, the original argument of Cassels may be readily adapted to prove the analogous statement for quasilinear p -forms. However, as pointed out by Hoffmann ([Hof04]), the additivity property of these forms enables one to prove a stronger multi-variable statement taking the following form:

Theorem 2.23 (cf. [Hof04, Cor. 3.4]). *Let ϕ be a quasilinear p -form over F , let $T = (T_1, \dots, T_m)$ be a tuple of algebraically independent variables over F and let $f \in F[T]$. Then $f \in D(\phi_{F(T)})$ if and only if $f \in D(\phi_{F(T)})$, if and only if $f \in D(\phi)[T_1^p, \dots, T_m^p]$.*

Now, in the situation of Theorem 2.23, the $F(T)^p$ -vector space $D(\phi_{F(T)})$ is (evidently) spanned by elements of $D(\phi)$. Thus, in view of Lemma 2.13, we immediately obtain the following result concerning rational similarity factors:

Corollary 2.24 (cf. [Hof04, Prop. 6.7]). *Let ϕ be a quasilinear p -form over F , let $T = (T_1, \dots, T_m)$ be a tuple of algebraically independent variables over F and let $f \in F[T]$. Then $f \in G(\phi_{F(T)})$ if and only if $f \in G(\phi)[T_1^p, \dots, T_m^p]$.*

2.8. Isotropy of quasilinear p -forms under scalar extension. We now collect some basic facts regarding the isotropy of quasilinear p -forms under scalar extension.

Let K and L be extensions of a field k . Recall that a k -place $K \dashrightarrow L$ is a pair (R, f) consisting of a valuation subring $k \subseteq R \subseteq K$ and a local k -algebra homomorphism $f: R \rightarrow L$. For example, given an inclusion $i: K \hookrightarrow L$, the pair (K, i) defines a k -place $K \dashrightarrow L$. If there exist k -places $K \dashrightarrow L$ and $L \dashrightarrow K$, then we say that K and L are *equivalent over k* , and write $K \sim_k L$. For instance, this is easily seen to be the case

whenever L (resp. K) is a purely transcendental extension of K (resp. L) (see [EKM08, §103] for further details). We have here the following basic lemma, which is a consequence of the completeness of X_ϕ (see [Gro61, (7.3.8)]):

Lemma 2.25 (cf. [Scu14, Lem. 3.4]). *Let ϕ be a quasilinear p -form over F and let K and L be field extensions of F such that there exists an F -place $K \dashrightarrow L$. Then $\mathfrak{i}_0(\phi_L) \geq \mathfrak{i}_0(\phi_K)$. In particular, if $K \sim_F L$, then $\mathfrak{i}_0(\phi_K) = \mathfrak{i}_0(\phi_L)$.*

Note, in particular, that passage to rational extensions of the base field does not affect the isotropy index of a quasilinear p -form. By MacLane's theorem ([Lan02, Prop. VIII.4]), the same is, in fact, true of arbitrary *separable* extensions:¹⁰

Lemma 2.26 (cf. [Hof04, Prop. 5.3]). *Let ϕ be an anisotropic quasilinear p -form over F and let L be a field extension of F . If L is separable over F , then ϕ_L is anisotropic and $\text{ndeg}(\phi_L) = \text{ndeg}(\phi)$.*

Thus, in order to study the isotropy behaviour of quasilinear p -forms under scalar extension, we are effectively reduced to considering the case of purely inseparable algebraic extensions. In degree p , we have the following basic observations, all of which can be easily verified using the results which have been discussed thus far (recall here that, given $a_1, \dots, a_n \in F$, F_{a_1, \dots, a_n} denotes the field $F(\sqrt[p]{a_1}, \dots, \sqrt[p]{a_n})$):

Lemma 2.27 (see [Hof04, §5], [Scu14, Lem. 3.8]). *Let ϕ be a quasilinear p -form over F and let $a \in F \setminus F^p$. Then:*

- (1) $D(\phi_{F_a}) = D(\langle\langle a \rangle\rangle \otimes \phi) = \sum_{i=0}^{p-1} a^i D(\phi)$.
- (2) $\mathfrak{i}_0(\phi_{F_a}) = \frac{1}{p} \mathfrak{i}_0(\langle\langle a \rangle\rangle \otimes \phi)$.
- (3) $\text{ndeg}(\phi_{F_a}) = \begin{cases} \frac{1}{p} \text{ndeg}(\phi) & \text{if } a \in N(\phi), \\ \text{ndeg}(\phi) & \text{if } a \notin N(\phi). \end{cases}$
- (4) *If ϕ is anisotropic and $a \notin N(\phi)$, then ϕ_{F_a} is anisotropic.*
- (5) $\dim(\phi_{F_a})_{\text{an}} \geq \frac{1}{p} \dim \phi_{\text{an}}$.
- (6) *Equality holds in (5) if and only if ϕ_{an} is divisible by $\langle\langle a \rangle\rangle$, if and only if $a \in G(\phi)$.*

Remark 2.28. The second equivalence in (6) holds by Corollary 2.20.

As an application of the first part of the lemma, we have:

Corollary 2.29. *Let ϕ be a quasilinear p -form over F and let $a \in F \setminus F^p$. Then $G(\phi_{F_a}) = G(\langle\langle a \rangle\rangle \otimes \phi)$. In particular, $\mathfrak{d}_0(\langle\langle a \rangle\rangle \otimes \phi) = \mathfrak{d}_0(\phi_{F_a}) + 1$.*

Proof. More specifically, the first statement is an immediate consequence of Lemma 2.27 (1) and Lemma 2.13. The second then follows from Corollary 2.19. \square

Suppose now that $\pi = \langle\langle a_1, \dots, a_n \rangle\rangle$ is an *anisotropic* quasi-Pfister p -form over F . By Remark 2.9, we have $[F^p(a_1, \dots, a_n) : F^p] = p^n$, which means that $a_i \notin F_{a_1, \dots, a_{i-1}}$ for every $1 \leq i \leq n$. Repeated applications of Lemma 2.27 (2) and 2.29 therefore yield the following proposition:

Proposition 2.30. *Let ϕ be a quasilinear p -form over F , let π be as above and let $\psi = \pi \otimes \phi$. Then $\mathfrak{i}_0(\psi) = p^n \mathfrak{i}_0(\phi_{F_{a_1, \dots, a_n}})$ and $\mathfrak{d}_0(\psi) = \mathfrak{d}_0(\phi_{F_{a_1, \dots, a_n}}) + n$.*

¹⁰Recall that an extension of fields $k \subseteq L$ is called *separable* if, for any algebraic closure \bar{k} of k , the ring $L \otimes_k \bar{k}$ has no nontrivial nilpotent elements.

Now, in view of Remark 2.6 (1), one may combine the above results in order to study the isotropy behaviour of quasilinear p -forms under scalar extension to function fields of quasilinear p -hypersurfaces. More specifically, let ψ be a quasilinear p -form over F which is not split, and let $a_0, \dots, a_n \in F$ be such that $\psi \simeq \langle a_0, \dots, a_n \rangle$, with $a_0, a_1 \neq 0$. Then, by Remark 2.6 (1), we have an F -isomorphism of fields

$$F(\psi) \simeq F(T) \left(\sqrt[p]{a_1^{-1}(a_0 + a_2 T_2^p + \dots + a_n T_n^p)} \right),$$

where $T = (T_2, \dots, T_n)$ is an $(n-1)$ -tuple of algebraically independent variables over F . Thus, putting Lemmas 2.26 and 2.27 and together, we obtain:

Lemma 2.31 (see [Hof04, §§7.3,7.4]). *Let ϕ be an anisotropic quasilinear p -form over F , and let ψ be as above. Then:*

- (1) $\dim(\phi_{F(\psi)})_{\text{an}} \geq \frac{1}{p} \dim \phi$.
- (2) Equality holds in (1) if and only if $a_1^{-1}(a_0 + a_2 T_2^p + \dots + a_n T_n^p) \in G(\phi_{F(T)})$.
- (3) $\text{ndeg}(\phi_{F(\psi)}) \geq \frac{1}{p} \text{ndeg}(\phi)$.
- (4) Equality holds in (3) if and only if $a_1^{-1}(a_0 + a_2 T_2^p + \dots + a_n T_n^p) \in N(\phi_{F(T)})$.
- (5) The equivalent conditions of (4) are satisfied if $\phi_{F(\psi)}$ is isotropic.

As a basic application, we have:

Corollary 2.32 (see [Hof04, §§7.3,7.4]). *Let ϕ and ψ be quasilinear p -forms over F such that ϕ is anisotropic and ψ is not split. Then:*

- (1) $\dim(\phi_{F(\psi)})_{\text{an}} \geq \frac{1}{p} \dim \phi$, with equality holding if and only if $N(\psi) \subseteq G(\phi)$.
- (2) If $\phi_{F(\psi)}$ is isotropic, then $N(\psi) \subseteq N(\phi)$. In particular, $\text{ndeg}(\psi) \leq \text{ndeg}(\phi)$.

Proof. We may assume that ψ is as in Lemma 2.31. In this case, we have $N(\psi) = F^p(\frac{a_0}{a_1}, \dots, \frac{a_n}{a_1})$ (see Remark 2.9), and so (1) follows from the first two parts of the former lemma and Corollary 2.24. Similarly, since $N(\phi_L) = D((\phi_{\text{nor}})_L)$ for any field extension L of F , (2) follows from Lemma 2.31 (4,5) and Theorem 2.23. \square

Finally, it will be useful to record in this section another basic application of the Cassels-Pfister theorem. To state it, let $T = (T_1, \dots, T_m)$ be a tuple of algebraically independent variables over F , let $g \in F[T]$ be an irreducible polynomial and let $F[g]$ denote the field $\text{Frac}(F[T]/(g))$ (i.e. the function field of the integral hypersurface $\{g=0\} \subset \mathbb{A}_F^m$). Given $f \in F[T]$, we write $\text{mult}_g(f)$ for the multiplicity of g in f , i.e., the largest nonnegative integer s such that $f = g^s h$ for some $h \in F[T]$.¹¹

Proposition 2.33. *In the above situation, let ϕ be a quasilinear p -form over F and let $f \in F[T]$. Suppose that $f \in D(\phi_{F(T)})$ and that $\phi_{F[g]}$ is anisotropic. Then $\text{mult}_g(f) \equiv 0 \pmod{p}$.*

Proof. Let $s = \text{mult}_g(f)$. After replacing f by $f/g^{kp} \in D(\phi_{F(T)})$ for a suitable integer $k \geq 0$, we may assume that $s < p$. Our goal is then to prove that $s = 0$. To see this, note first that there exists $v \in V_\phi \otimes_F F[T]$ such that $\phi_{F(T)}(v) = f$ by Theorem 2.23. If $s \neq 0$, then the image \bar{v} of v in $V_\phi \otimes_F F[g]$ is an isotropic vector for $\phi_{F[g]}$. By hypothesis, it follows that $\bar{v} = 0$, which means that $v = gw$ for some $w \in V_\phi \otimes_F F[T]$. But this implies that $f = g^p \phi_{F(T)}(w)$, which contradicts the fact that $s < p$. We conclude that $s = 0$, and so the proposition is proved. \square

¹¹With the added convention that $\text{mult}_g(0) = +\infty$.

2.9. The divisibility index and scalar extension. Let ϕ be a quasilinear p -form over F . We make some brief remarks concerning the behaviour of the divisibility index $\mathfrak{d}_0(\phi)$ (see §2.3) under scalar extension.

Lemma 2.34. *Let ϕ be a quasilinear p -form over F and let L be a field extension of F . If $(\phi_{\text{sim}})_L$ is anisotropic, then $\mathfrak{d}_0(\phi_L) \geq \mathfrak{d}_0(\phi)$.*

Proof. As an L^p -vector space, $D((\phi_{\text{sim}})_L)$ is spanned by $D(\phi_{\text{sim}}) = G(\phi)$. Since we evidently have $G(\phi) \subseteq G(\phi_L) = D((\phi_L)_{\text{sim}})$, and since $(\phi_{\text{sim}})_L$ is anisotropic by hypothesis, Proposition 2.3 implies that $(\phi_{\text{sim}})_L \subset (\phi_L)_{\text{sim}}$. The desired assertion now follows from Corollary 2.19. \square

In particular, this applies in the case where L is a separable extension of F (see Lemma 2.26). In the case where L is purely transcendental over F , we can say more:

Lemma 2.35. *Let ϕ be a quasilinear p -form over F and let L be a purely transcendental extension of F . Then $(\phi_L)_{\text{sim}} \simeq (\phi_{\text{sim}})_L$ and $\mathfrak{d}_0(\phi_L) = \mathfrak{d}_0(\phi)$.*

Proof. Continuing with the proof of Lemma 2.34, it is sufficient to show that in this case $G(\phi_L)$ is generated by $G(\phi)$ over L^p . If L is finitely generated over F , then this follows from Corollary 2.24. On the other hand, the general case reduces easily to the finitely generated case in view of Lemma 2.13, so the lemma is proved. \square

2.10. The Knebusch splitting pattern. Let ϕ be a quasilinear p -form over F . Following the construction outlined in §1 (see also [Hof04, §7.5]), set $F_0 = F$, $\phi_0 = \phi_{\text{an}}$, and recursively define

- $F_r = F_{r-1}(\phi_{r-1})$ (provided ϕ_{r-1} is not split), and
- $\phi_r = (\phi_{F_r})_{\text{an}}$ (provided F_r is defined).

Note here that if ϕ_r is defined, then we have $\dim \phi_r < \dim \phi_{r-1}$ by Remark 2.6 (4). As such, the whole process is finite, terminating at the first nonnegative integer $h(\phi)$ for which $\dim \phi_{h(\phi)} \leq 1$. The integer $h(\phi)$ will be called the *height* of ϕ , and the tower of fields $F_0 \subset F_1 \subset \dots \subset F_{h(\phi)}$ will be called the *Knebusch splitting tower* of ϕ . For each $0 \leq r \leq h(\phi)$, we set $\mathfrak{j}_r(\phi) = \mathfrak{i}_0(\phi_{F_r})$. If ϕ is not split and $r \geq 1$, then the difference $\mathfrak{j}_r(\phi) - \mathfrak{j}_{r-1}(\phi)$ will be called the *r -th higher isotropy index* of ϕ , and will be denoted by $\mathfrak{i}_r(\phi)$. In this case, the form ϕ_r will be called the *r -th higher anisotropic kernel* of ϕ . Finally, the sequence $\mathfrak{i}(\phi) = (\mathfrak{i}_1(\phi), \dots, \mathfrak{i}_{h(\phi)}(\phi))$ (understood to be empty if ϕ is split) will be called the *Knebusch splitting pattern* of ϕ .¹²

Remarks 2.36. Let ϕ be a quasilinear p -form over F .

- (1) By the recursive nature of the above construction, we have $\mathfrak{i}_r(\phi) = \mathfrak{i}_1(\phi_{r-1})$ for every $1 \leq r \leq h(\phi)$.
- (2) By Proposition 2.4, we have $\mathfrak{i}_r(\phi) = \dim \phi_{r-1} - \dim \phi_r$ for all $1 \leq r \leq h(\phi)$.
- (3) Let L be a field extension of F . As already remarked in §1, it is not true in general that $\mathfrak{i}_0(\phi_L) = \mathfrak{j}_r(\phi)$ for some $0 \leq r \leq h(\phi)$ - see Example 2.47 below.

Note that by Remark 2.6 (3), we have the following:

Lemma 2.37. *Let ϕ be a quasilinear p -form form of dimension ≥ 2 and let (F_r) denote its Knebusch splitting tower. Then $F_r \sim_F F(\phi^{\times r})$ for every $0 \leq r \leq h(\phi)$.*

In light of Lemma 2.37, we therefore have:

¹²See Footnote 3. We omit the term $\mathfrak{i}_0(\phi)$ from the sequence because we are ultimately interested in the case where ϕ is anisotropic (i.e., where $\mathfrak{i}_0(\phi) = 0$).

Corollary 2.38. *Let ϕ be a quasilinear p -form over F . Then, for every $0 \leq r \leq h(\phi)$, we have $j_r(\phi) = i_0(\phi_{F(\phi^{\times r})})$.*

Given the results of §2.8, we are now in a position to prove the following characterisation of anisotropic quasi-Pfister p -forms:

Proposition 2.39 (cf. [HL04, Thm. 8.11]). *Let ϕ be an anisotropic quasilinear p -form of dimension ≥ 2 over F . Then $\dim \phi_1 \geq \frac{1}{p} \dim \phi$, and the following conditions are equivalent:*

- (1) $\dim \phi_1 = \frac{1}{p} \dim \phi$.
- (2) $i(\phi) = (p^{h(\phi)} - p^{h(\phi)-1}, p^{h(\phi)-1} - p^{h(\phi)-2}, \dots, p^2 - p, p - 1)$.
- (3) ϕ is similar to a quasi-Pfister p -form.

Proof. The inequality $\dim \phi_1 \geq \frac{1}{p} \dim \phi$ holds by Corollary 2.32. The same result shows that equality holds here if and only if $N(\phi) \subseteq G(\phi)$. By Corollary 2.20, the latter condition holds if and only if ϕ is divisible by ϕ_{nor} . In view of Lemma 2.11, this proves the equivalence of (1) and (3), as well as the implication (2) \Rightarrow (3). On the other hand, if ϕ is similar to a quasi-Pfister p -form of dimension p^n , then ϕ_1 is similar to a quasi-Pfister p -form of dimension p^{n-1} by Corollary 2.8 and (1). Since $i_1(\phi) = \dim \phi - \dim \phi_1$ (see Remark 2.36 (2)), an easy induction on $h(\phi)$ then shows that (3) implies (2). \square

Finally, a repeated application of Lemma 2.31 (3-5) (with $\psi = \phi$) yields the following computation of the height $h(\phi)$:

Corollary 2.40 (cf. [Hof04, Thm. 7.25 (ii)]). *Let ϕ be a quasilinear p -form over F . Then $h(\phi) = \log_p(\text{ndeg}(\phi))$.*

Together with Corollary 2.32 (2), this implies the following useful result:

Corollary 2.41 (cf. [Scu14, Prop. 4.12]). *Let ϕ and ψ be quasilinear p -forms over F such that ϕ is anisotropic and ψ is not split. If $\phi_{F(\psi)}$ is isotropic, then $h(\psi) \leq h(\phi)$.*

2.11. The quasi-Pfister height and higher divisibility indices. Let ϕ be a quasilinear p -form over F . As in [Scu14, §4.2], we define the *quasi-Pfister height* of ϕ , denoted $h_{\text{qp}}(\phi)$, to be the smallest nonnegative integer l such that ϕ_l is similar to a quasi-Pfister p -form (this is well defined, since $\phi_{h(\phi)}$, being of dimension 1, is similar to a 0-fold quasi-Pfister p -form). We have:

Lemma 2.42. *Let ϕ be a quasilinear p -form over F and let $d = h(\phi) - h_{\text{qp}}(\phi)$. Then $i(\phi) = (i_1(\phi), \dots, i_{h_{\text{qp}}(\phi)}(\phi), p^d - p^{d-1}, p^{d-1} - p^{d-2}, \dots, p^2 - p, p - 1)$ and $i_{h_{\text{qp}}(\phi)}(\phi) = \dim(\phi) - j_{h_{\text{qp}}(\phi)-1}(\phi) - p^d < p^{d+1} - p^d$.*

Proof. The first statement is an immediate consequence of Proposition 2.39. The point here is that $\phi_{h_{\text{qp}}(\phi)}$ is similar to a quasi-Pfister p -form of dimension p^d . By Remark 2.36 (2), we therefore have $i_{h_{\text{qp}}(\phi)}(\phi) = \dim \phi_{h_{\text{qp}}(\phi)-1} - \dim \phi_{h_{\text{qp}}(\phi)} = \dim(\phi) - j_{h_{\text{qp}}(\phi)-1}(\phi) - p^d$. Finally, since $\phi_{h_{\text{qp}}(\phi)-1}$ is (by the definition of $h_{\text{qp}}(\phi)$) not similar to a quasi-Pfister p -form, it must have dimension $< p^{d+1}$, again by Proposition 2.39. This proves the inequality in the second statement, and hence the lemma. \square

The Knebusch splitting pattern of a quasilinear p -form ϕ is therefore determined by $h(\phi)$, $h_{\text{qp}}(\phi)$ and the truncated sequence $(i_1(\phi), \dots, i_{h_{\text{qp}}(\phi)}(\phi))$. With a view to studying the latter invariant, we now introduce new invariants of ϕ which will be of central interest in the sequel. More specifically, for each $1 \leq r \leq h(\phi)$, we define the *r -th higher divisibility index* of ϕ , denoted $\mathfrak{d}_r(\phi)$, as the integer $\mathfrak{d}_0(\phi_r)$ (see §2.3). In other words, $\mathfrak{d}_r(\phi)$ is the largest integer s such that ϕ_r is divisible by an s -fold quasi-Pfister p -form

(over the corresponding field of the Knebusch splitting tower of ϕ). The sequence of integers $(\mathfrak{d}_0(\phi), \dots, \mathfrak{d}_{h(\phi)}(\phi))$ will be denoted by $\mathfrak{d}(\phi)$. As per Lemma 2.42 (and the proof of Proposition 2.39), we have:

Lemma 2.43. *Let ϕ be a quasilinear p -form over F and let $d = h(\phi) - h_{\text{qp}}(\phi)$. Then $\mathfrak{d}(\phi) = (\mathfrak{d}_0(\phi), \dots, \mathfrak{d}_{h_{\text{qp}}(\phi)-1}(\phi), d, d-1, \dots, 1, 0)$.*

We also, however, have the following information concerning the “nontrivial part” of the sequence $\mathfrak{d}(\phi)$:

Lemma 2.44. *Let ϕ be a quasilinear p -form over F . Then $\mathfrak{d}_0(\phi) \leq \dots \leq \mathfrak{d}_{h_{\text{qp}}(\phi)}$.*

Proof. We may assume that ϕ is anisotropic and not split. We need to show that if ϕ is not similar to a quasi-Pfister p -form, then $\mathfrak{d}_1(\phi) \geq \mathfrak{d}_0(\phi)$. By Lemma 2.34 it will be sufficient to check that ϕ_{sim} remains anisotropic over $F(\phi)$. Suppose otherwise. Then, by Corollary 2.32 (1), we have $N(\phi) \subseteq G(\phi)$. By Corollary 2.15 it follows that $N(\phi) = G(\phi)$, or, equivalently, that $\phi_{\text{nor}} \simeq \phi_{\text{sim}}$ (see Lemma 2.2). But, in view of Lemma 2.11 and Corollary 2.18, this implies that ϕ is similar to a quasi-Pfister p -form, thus contradicting our assumption. The lemma follows. \square

In particular, since $\mathfrak{i}_r(\phi) = \dim \phi_r - \dim \phi_{r-1}$ for all $1 \leq r \leq h(\phi)$ (see Remark 2.36 (2)), we obtain the following result concerning the integers $\mathfrak{i}_r(\phi)$:

Corollary 2.45. *Let ϕ be a quasilinear p -form over F . Then $\mathfrak{i}_r(\phi) \equiv 0 \pmod{\mathfrak{d}_{r-1}(\phi)}$ for all $1 \leq r \leq h_{\text{qp}}(\phi)$.*

2.12. Some examples. We now conclude this section with two basic computations which will be needed in the sequel, beginning with:

Lemma 2.46. *Let ϕ be an anisotropic quasilinear p -form over F and let $\psi = \phi_{F(T)} \perp \langle T \rangle$, where T is an algebraically independent variable over F . Then $\mathfrak{i}(\psi) = (1, \mathfrak{i}_1(\phi), \mathfrak{i}_2(\phi), \dots, \mathfrak{i}_{h(\phi)}(\phi))$ and $\mathfrak{d}(\psi) = (0, \mathfrak{d}_0(\phi), \mathfrak{d}_1(\phi), \dots, \mathfrak{d}_{h(\phi)}(\phi))$.*

Proof. The form ψ is clearly anisotropic. Now, the field $F(T)(\psi)$ is F -isomorphic to a purely transcendental extension of F (see the presentation of Remark 2.6 (1), for example). In particular, $\phi_{F(T)(\psi)}$ is anisotropic (Lemma 2.26), and so $\mathfrak{i}_1(\psi) = 1$ and $\psi_1 \simeq \phi_{F(T)(\psi)}$ (see Remark 2.5). Since $F(T)(\psi)$ is purely transcendental over F , the first statement now follows immediately from Lemma 2.26. In a similar way, Lemma 2.35 implies that $\mathfrak{d}_r(\psi) = \mathfrak{d}_{r-1}(\phi)$ for all $1 \leq r \leq h(\psi)$. Thus, to prove the second statement, it only remains to check that $\mathfrak{d}_0(\psi) = 0$. But, since $\mathfrak{i}_1(\psi) = 1$, this is an immediate consequence of Corollary 2.45. \square

Given this result, we can give an example of a quasilinear p -form whose (full) splitting pattern is *not* determined its Knebusch splitting pattern:

Example 2.47 (cf. [HL04, Ex. 8.15]). Let $T = (T_1, \dots, T_{n+1})$ be a tuple of algebraically independent variables over a field F_0 of characteristic p , and let $F = F_0(T)$. Consider the form $\phi = \langle\langle T_1, \dots, T_n \rangle\rangle \perp \langle T_{n+1} \rangle$ over F . By Lemma 2.46 and Proposition 2.39, we have $\mathfrak{i}(\phi) = (1, p^n - p^{n-1}, p^{n-1} - p^{n-2}, \dots, p^2 - p, p - 1)$, so that $\mathfrak{j}_r(\phi) = p^n - p^{n-r+1} + 1$ for all $1 \leq r \leq n + 1$. On the other hand, the full splitting pattern of ϕ also contains all the integers $p^n - p^{n-r+1}$ ($1 \leq r \leq n + 1$). Indeed, if (L_s) denotes the Knebusch splitting tower of $\langle\langle a_1, \dots, a_n \rangle\rangle_L$, then we clearly have $\mathfrak{i}_0(\phi_{L_{r-1}}) = p^n - p^{n-r+1}$ for all $1 \leq r \leq n + 1$ (again, we are using Lemma 2.26 and Proposition 2.39 here).

Our second computation is the following:

Lemma 2.48. *Let ϕ be a quasilinear p -form over F and let $\psi = \langle\langle T_1, \dots, T_n \rangle\rangle \otimes \phi_{F(T)}$, where $T = (T_1, \dots, T_n)$ is a tuple of algebraically independent variables over F . Then $\mathfrak{i}(\psi) = (p^n \mathfrak{i}_1(\phi), \dots, p^n \mathfrak{i}_{h(\phi)}(\phi), p^n - p^{n-1}, p^{n-1} - p^{n-2}, \dots, p^2 - p, p - 1)$ and $\mathfrak{d}(\psi) = (\mathfrak{d}_0(\phi) + n, \mathfrak{d}_1(\phi) + n, \dots, \mathfrak{d}_{h(\phi)-1}(\phi) + n, n, n - 1, \dots, 1, 0)$.*

Proof. It is enough to treat the case where $n = 1$. To simplify the notation, let us write T for the variable T_1 and L for the rational function field $F(T)$. Now, by construction, we have $\text{ndeg}(\psi) = p(\text{ndeg}(\phi))$ (see Remark 2.10). In view of Corollary 2.40, it follows that $h(\psi) = h(\phi) + 1$. Let (L_r) and (F_r) denote the Knebusch splitting towers of ψ and ϕ respectively. We claim that, for every $0 \leq r \leq h(\phi)$, $(L_r)_T$ is F -isomorphic to a purely transcendental extension of F_r . The case where $r = 0$ is evident. In general, we have $(L_r)_T = (L_T)_r$, where $((L_T)_r)$ denotes the Knebusch splitting tower of ψ_{L_T} . But, since $\langle\langle T \rangle\rangle_{L_T} \sim \langle 1 \rangle$, and since L_T is purely transcendental over F (the $r = 0$ case), we have $(\psi_{L_T})_{\text{an}} \simeq \phi_{L_T}$. By Remark 2.5, it follows that $(L_r)_T$ is L -isomorphic to a purely transcendental extension of the free composite $F_r \cdot L_T$. Again, since L_T is purely transcendental over F , the claim follows. Given this, Proposition 2.30 and Lemma 2.26 together imply that

$$\mathfrak{j}_r(\psi) = \mathfrak{i}_0(\psi_{L_r}) = p\mathfrak{i}_0(\phi_{(L_r)_T}) = p\mathfrak{i}_0(\phi_{F_r}) = p\mathfrak{j}_r(\phi)$$

for all $0 \leq r \leq h(\phi)$, which proves the first statement of the lemma. Similarly, our claim, Proposition 2.30 and Lemma 2.35 together imply that

$$\mathfrak{d}_r(\psi) = \mathfrak{d}_0(\psi_{L_r}) = \mathfrak{d}_0(\phi_{(L_r)_T}) + 1 = \mathfrak{d}_0(\phi_{F_r}) + 1 = \mathfrak{d}_r(\phi) + 1$$

for all $0 \leq r \leq h(\phi)$, and so the second statement also holds. \square

3. AN INCOMPRESSIBILITY THEOREM AND RELATED RESULTS

In this section, we collect some of the farther-reaching results on the isotropy behaviour of quasilinear p -forms over function fields of quasilinear p -hypersurfaces which have been obtained in recent years. These results will have an essential role to play in the sequel. We do not provide full details here, but the interested reader is referred to the original articles ([HL04, Tot08, Scu14]) for further information.

3.1. The incompressibility theorem. Let ϕ be an anisotropic quasilinear p -form of dimension ≥ 2 over F with associated quasilinear p -hypersurface X_ϕ . As in [Scu13, §5], we define the *Izhboldin dimension* of X_ϕ , denoted $\dim_{\text{Izh}}(X_\phi)$, to be the integer $\dim X_\phi - \mathfrak{i}_1(\phi) + 1$. The following result was proved in *loc. cit.*:

Theorem 3.1 ([Scu13, Thm. 5.12]). *Let X be an anisotropic quasilinear p -hypersurface over F and let Y be an algebraic variety over F such that $Y(F_{\text{sep}}) = \emptyset$. If $\dim Y < \dim_{\text{Izh}}(X)$, then there cannot exist a rational map $X \dashrightarrow Y$.*

Remarks 3.2. (1) In the case where $p = 2$, Theorem 3.1 is due to Totaro ([Tot08, Thm. 5.1]). In fact, if $X_\phi = \{\phi = 0\}$ is an anisotropic projective quadric over a field k of any characteristic, and if Y is any complete k -variety possessing no closed points of odd degree, then it is known that the existence of a rational map $X \dashrightarrow Y$ necessarily implies that $\dim Y \geq \dim_{\text{Izh}}(X_\phi)$, where $\dim_{\text{Izh}}(X_\phi) = \dim X_\phi - \mathfrak{i}_1(\phi) + 1$. This result was first proved by Karpenko and Merkurjev in the case where X_ϕ is smooth ([KM03, Thm 4.1], [EKM08, Thm. 76.5]), and was later extended by Totaro (*loc. cit.*) to the singular case.

(2) The special case where Y is a closed subvariety of X shows that the canonical dimension of X (see §1) is equal to $\dim_{\text{Izh}}(X)$ (the inequality $\text{cdim}(X) \leq \dim_{\text{Izh}}(X)$ is trivial – see [Scu13, Cor. 5.14]).

In the remainder of this section, we will recall some of the main applications of Theorem 3.1 (and its proof). Here, we mention the following:

Corollary 3.3 (cf. [Scu14, Cor. 5.4]). *Let ϕ and ψ be anisotropic quasilinear p -forms of dimension ≥ 2 over F , and let $\sigma \subset \phi$ be a subform of dimension $\leq \dim \psi - \mathbf{i}_1(\psi)$. Then $\sigma_{F(\psi)} \subset (\phi_{F(\psi)})_{\text{an}}$. In particular, $\sigma_{F(\psi)}$ is anisotropic.*

Proof. We trivially have $D(\sigma_{F(\psi)}) \subseteq D(\phi_{F(\psi)})$. In light of Proposition 2.3, it therefore suffices to check that $\sigma_{F(\psi)}$ is anisotropic, or, equivalently, that there does not exist a rational map $X_\psi \dashrightarrow X_\sigma$. But, since $X_\sigma(F_{\text{sep}}) = \emptyset$ (see Lemma 2.26), this is an immediate consequence of Theorem 3.1. \square

In particular, we have the following fundamental observation:

Corollary 3.4. *Let ϕ be an anisotropic quasilinear p -form of dimension ≥ 2 over F and let $\psi \subset \phi$ be a subform of codimension $\mathbf{i}_1(\phi)$. Then $\phi_1 \simeq \psi_{F(\phi)}$.*

Proof. By Corollary 3.3, we have $\psi_{F(\phi)} \subset \phi_1$. Since both forms have the same dimension by hypothesis, the result follows. \square

3.2. Neighbours and near neighbours, I. Let ψ and ϕ be anisotropic quasilinear p -forms of dimension ≥ 2 over F . We will say that ψ is a *neighbour* (resp. *near neighbour*) of ϕ if ψ is similar to a subform of codimension $< \mathbf{i}_1(\phi)$ (resp. codimension $\mathbf{i}_1(\phi)$) of ϕ . Our motivation here is the following extension of Corollary 3.4:

Lemma 3.5. *Let ϕ and ψ be anisotropic quasilinear p -forms over F such that ϕ is anisotropic of dimension ≥ 2 and ψ is similar to a subform of ϕ . Then $(\psi_{F(\phi)})_{\text{an}}$ is similar to ϕ_1 if and only if ψ is a neighbour or near neighbour of ϕ .*

Proof. We may assume that $\psi \subset \phi$. Again, we trivially have $D(\psi_{F(\phi)}) \subseteq D(\phi_{F(\phi)}) = D(\phi_1)$. By Lemma 2.3, it therefore suffices to check that $\dim(\psi_{F(\phi)})_{\text{an}} = \dim \phi - \mathbf{i}_1(\phi)$ if and only if ψ has codimension $\leq \mathbf{i}_1(\phi)$ in ϕ . The left-to-right implication here is trivial. Conversely, if ψ has codimension $\leq \mathbf{i}_1(\phi)$ in ϕ , then we have $\dim(\psi_{F(\phi)})_{\text{an}} \geq \dim \phi - \mathbf{i}_1(\phi)$ by Theorem 3.1. Since the reverse inequality holds here by obvious dimension reasons, the lemma is proved. \square

Note here that while neighbours of ϕ become anisotropic over $F(\phi)$, its *near* neighbours do not (Corollary 3.3). This enables us to compute:

Proposition 3.6 (cf. [Scu13, Prop. 6.1]). *Let ϕ and ψ be anisotropic quasilinear p -forms of dimension ≥ 2 over F such that ψ is a codimension- d neighbour of ϕ . Then $\mathbf{i}_1(\psi) = \mathbf{i}_1(\phi) - d$.*

3.3. The Ruledness theorem. Another key application of Theorem 3.1 is the following extension of Proposition 3.6, which shows in a precise way that anisotropic quasilinear p -hypersurfaces having first higher isotropy index larger than 1 are ruled.

Theorem 3.7 (cf. [Scu13, Thm. 7.6]). *Let ϕ and ψ be anisotropic quasilinear p -forms of dimension ≥ 2 over F such that ψ is a codimension- d neighbour of ϕ . Then X_ϕ is birationally isomorphic to $X_\psi \times_F \mathbb{P}^d$.*

Proof. By Proposition 3.6, we have $\mathbf{i}_1(\psi) = \mathbf{i}_1(\phi) - d$. It is therefore enough to treat the case where $d = \mathbf{i}_1(\psi) - 1$, and this is covered by [Scu13, Thm. 7.6]. \square

Remark 3.8. Again, in the case where $p = 2$, this result is due to Totaro ([Tot08, Thm. 6.4]). Unlike Theorem 3.1, however, the analogous assertion remains open for generically smooth quadrics (in any characteristic - see [Tot08, Tot09]).

It is worth mentioning the following explicitly:

Corollary 3.9. *Let ϕ and ψ be an anisotropic quasilinear p -forms of dimension ≥ 2 over F such that $\mathbf{i}_1(\phi) > 1$ and ψ is similar to a codimension-1 subform of ϕ . Then we have an F -isomorphism of fields $F(\phi) \simeq F[\psi]$.*

Proof. By Theorem 3.7, $F(\phi)$ is F -isomorphic to a degree-one purely transcendental extension of $F(\psi)$. In view of Remark 2.6 (3), the result follows. \square

3.4. Neighbours and near neighbours, II. Given Theorem 3.7, one can extend Proposition 3.6 as follows:

Proposition 3.10 (cf. [Scu14, Prop. 6.2]). *Let ϕ and ψ be anisotropic quasilinear p -forms of dimension ≥ 2 over F such that ψ is a codimension- d neighbour of ϕ . Then $\mathbf{i}(\psi) = (\mathbf{i}_1(\phi) - d, \mathbf{i}_2(\phi), \dots, \mathbf{i}_{h(\phi)}(\phi))$ and $\mathfrak{d}(\psi) = (\mathfrak{d}_0(\psi), \mathfrak{d}_1(\phi), \dots, \mathfrak{d}_{h(\phi)}(\phi))$.*

Proof. By Theorem 3.7, $F(\phi)$ is F -isomorphic to a purely transcendental extension of $F(\psi)$. Thus, if (F_r) and $(F(\phi)_r)$ denote the Knebusch splitting towers of ψ and $\psi_{F(\phi)}$, respectively, then $F(\phi)_r$ is F_r -isomorphic to a purely transcendental extension of F_{r+1} for every $0 \leq r \leq h(\psi_{F(\phi)})$. In particular, we have $\mathbf{i}_r(\psi_{F(\phi)}) = \mathbf{i}_{r+1}(\psi)$ and $\mathfrak{d}_r(\psi_{F(\phi)}) = \mathfrak{d}_{r+1}(\psi)$ for any such r by Lemmas 2.26 and 2.35, respectively. On the other hand, Lemma 3.5 shows that $(\psi_{F(\phi)})_{\text{an}}$ is similar to ϕ_1 . Since $\mathbf{i}_r(\phi) = \mathbf{i}_{r-1}(\phi_1)$ and $\mathfrak{d}_r(\phi) = \mathfrak{d}_{r-1}(\phi_1)$ for all $1 \leq r \leq h(\phi)$, the proposition follows immediately. \square

Let ϕ be a quasilinear p -form over F . We say that ϕ is a *quasi-Pfister p -neighbour* (of π) if there exists a quasi-Pfister p -form π over F such that ϕ is similar to a subform of π and $\dim \phi > \frac{1}{p} \dim \pi$. If ϕ is anisotropic, then it follows from Proposition 2.39 that ϕ is quasi-Pfister p -neighbour if and only if it is a neighbour of some quasi-Pfister p -form in the sense of §3.2. Forms of this type are of special importance in the general theory of quasilinear p -forms. Putting Lemma 2.11, Proposition 2.39, Corollary 2.40, Lemma 3.5 and Proposition 3.10 together, we obtain the following classification of anisotropic quasi-Pfister p -neighbours:

Corollary 3.11 (cf. [Scu14, Thm. 6.4]). *Let ϕ be an anisotropic quasilinear p -form of dimension ≥ 2 over F and let n be the smallest nonnegative integer such that $p^{n+1} \geq \dim \phi$. Then the following are equivalent:*

- (1) ϕ is a quasi-Pfister p -neighbour.
- (2) ϕ is a neighbour of ϕ_{nor} .
- (3) $\phi_{F(\phi_{\text{nor}})}$ is isotropic.
- (4) $\text{ndeg}(\phi) = p^{n+1}$.
- (5) $h(\phi) = n + 1$.
- (6) $\mathbf{i}_1(\phi) = \dim \phi - p^n$ and $\mathbf{i}_2(\phi) = p^n - p^{n-1}$.
- (7) $\mathbf{i}(\phi) = (\dim \phi - p^n, p^n - p^{n-1}, p^{n-1} - p^{n-2}, \dots, p^2 - p, p - 1)$.
- (8) ϕ_1 is similar to a quasi-Pfister p -form (i.e., $h_{\text{qp}}(\phi) \leq 1$).

Remark 3.12. In the case where $p = 2$, this result was proved earlier by Hoffmann and Laghribi (cf. [HL04, Thm. 8.1]) using different methods.

For arbitrary subforms, the situation is naturally more complicated, but we can nevertheless appeal to the following general result which was proved in [Scu14] (and whose proof again makes essential use of Theorems 3.1 and 3.7):

Proposition 3.13 (cf. [Scu14, Prop. 8.6]). *Let ϕ and ψ be anisotropic quasilinear p -forms of dimension ≥ 2 over F such that $\phi_{F(\psi)}$ is isotropic. Then, either*

- (1) $(\psi_r)_{F_r(\phi)}$ is anisotropic for all $0 \leq r < h(\psi)$ and $\mathbf{i}(\psi_{F(\phi)}) = \mathbf{i}(\psi)$, or
- (2) $\mathbf{i}(\psi_{F(\phi)}) = (\mathbf{i}_1(\psi), \dots, \mathbf{i}_{s-1}(\psi), \mathbf{i}_s(\psi) + \mathbf{i}_{s+1}(\psi), \mathbf{i}_{s+2}(\psi), \dots, \mathbf{i}_{h(\psi)}(\psi))$, where $s < h(\psi)$ is the smallest nonnegative integer such that $(\psi_s)_{F_s(\phi)}$ is isotropic.

Note here that in the special situation where ψ is a neighbour of ϕ , we are necessarily in case (2) with s being equal to 0. Since $(\psi_{F(\phi)})_{\text{an}} \simeq \phi_1$ in this instance (Lemma 3.5), we thus recover the computation of Proposition 3.10. By contrast, if ψ is a *near* neighbour of ϕ , then we can be in either of cases (1) and (2) (see Remark 3.15 below). Nevertheless, we still have $\psi_{F(\phi)} \simeq \phi_1$ (Corollary 3.4), and so we get:

Corollary 3.14 (cf. [Scu14, Cor. 6.10]). *Let ϕ be an anisotropic quasilinear p -form of dimension ≥ 2 over F and let ψ be a near neighbour of ϕ . Then, either*

- (1) $(\psi_r)_{F_r(\phi)}$ is anisotropic for all $0 \leq r < h(\psi)$ and $\mathbf{i}(\psi) = \mathbf{i}(\phi_1)$, or
- (2) $\mathbf{i}(\psi) = (\mathbf{i}_2(\phi), \mathbf{i}_3(\phi), \dots, \mathbf{i}_s(\phi), \mathbf{i}_s(\psi), \mathbf{i}_{s+1}(\phi) - \mathbf{i}_s(\psi), \mathbf{i}_{s+2}(\phi) \dots, \mathbf{i}_{h(\phi)}(\phi))$, where $s < h(\psi)$ is the smallest positive integer such that $(\psi_s)_{F_s(\phi)}$ is isotropic.

Remark 3.15. As per the comments above, neither of cases (1) and (2) can be ruled out here. Indeed, case (1) describes the situation where $h(\psi) = h(\phi) - 1$, while case (2) describes that where $h(\psi) = h(\phi)$. As the reader will easily verify using Corollary 2.40, both these situations can arise in practice.

3.5. A comparison result. We now conclude this section by recalling the following comparison result for isotropy indices of quasilinear quadratic forms which was obtained in [Scu14] with the help of Theorem 3.1:

Proposition 3.16 (cf. [Scu14, Thm. 7.13, Rem. 9.1]). *Assume that $p = 2$. Let ϕ be an anisotropic quasilinear quadratic form of dimension ≥ 2 over F and let L be a field extension of F such that ϕ_L is not split. Then:*

$$\mathbf{i}_0(\phi_L) - \mathbf{i}_1(\phi) \geq \min \left\{ \mathbf{i}_0(\phi_L), \left\lceil \frac{\dim \phi - \mathbf{i}_1(\phi) + 1}{2} \right\rceil \right\}.$$

Remark 3.17. A similar statement also holds for $p > 2$ (see [Scu14, Thm. 7.13]), but this will not be needed below.

Finally, it will be convenient to record here the following application of this result which was also obtained in [Scu14]:

Theorem 3.18 ([Scu14, Thm. 9.2]). *Let ϕ be an anisotropic quasilinear quadratic form of dimension ≥ 2 over F and write $\dim \phi = 2^n + m$ for uniquely determined integers $n \geq 0$ and $1 \leq m \leq 2^n$. Then, for any field extension L of F , we either have $\mathbf{i}_0(\phi_L) \geq m$ or $\mathbf{i}_0(\phi_L) \leq m - \mathbf{i}_1(\phi)$.*

Remark 3.19. Taking $L = F(\phi)$ here, we see that if $\mathbf{i}_1(\phi) < m$, then $\mathbf{i}_1(\phi) \leq \frac{m}{2}$. This result will now be subsumed in our Theorem 1.3.

4. A MOTIVATIONAL EXAMPLE

As a warm-up for the proof of our main result, we will now prove Proposition 1.4. Throughout this section, we assume that $p = 2$. By a *quasi-Pfister form*, we will mean a quasi-Pfister 2-form. Our assumption on the prime p is imposed for simplicity, but also because we will make use of the following fact already mentioned in the introduction, the analogue of which is unknown when $p > 3$ (see [Scu14, §4.1]):

Lemma 4.1 (cf. [Hof04, Cor. 7.22]). *Let ϕ be an anisotropic quasilinear quadratic form of dimension ≥ 2 over F and let L be a field extension of F such that ϕ_L is isotropic. Then $\mathfrak{i}_0(\phi_L) \geq \mathfrak{i}_1(\phi)$.*

Now, let ϕ be an anisotropic quasilinear quadratic form of dimension ≥ 2 over F and let $\psi \subset \phi$ be a subform of codimension $\mathfrak{i}_1(\phi)$. By Corollary 3.4, $\psi_{F(\phi)}$ is isomorphic to the first higher anisotropic kernel of ϕ . In the next section, we will prove Theorem 1.5, which asserts that the latter form is divisible by a quasi-Pfister form of dimension $\geq \mathfrak{i}_1(\phi)$. Here, we will consider some special situations in which this divisibility property is already visible over the base field F . To this end, we will be interested in the following technical condition on the pair (ϕ, ψ) :

- (\star) There exist elements $a \in D(\phi) \setminus \{0\}$ and $b \in D(\psi) \setminus \{0\}$ such that $a \neq c(bd + ef)$ for any $c, d, e, f \in D(\psi)$.

Example 4.2. If, in the above situation, we have $\text{ndeg}(\psi) < \text{ndeg}(\phi)$ (equivalently, if $h(\psi) < h(\phi)$ – see Corollary 2.40), then (\star) holds for the pair (ϕ, ψ) . Indeed, in this case, $D(\phi)$ is (evidently) not contained in $cN(\psi)$ for any $c \in F$. Since $bd + ef \in N(\psi)$ for every $b, d, e, f \in D(\psi)$, the validity of (\star) is immediately verified.

In light of Example 4.2, Proposition 1.4 is subsumed in the following result:

Proposition 4.3. *Let ϕ be an anisotropic quasilinear quadratic form of dimension ≥ 2 over F and let $\psi \subset \phi$ be subform of codimension $\mathfrak{i}_1(\phi)$ such that the pair (ϕ, ψ) satisfies (\star). Then there exist a quasi-Pfister form π , a subform $\sigma \subset \pi$, an element $\lambda \in D(\phi)$ and a form τ over F such that $\psi \simeq \pi \otimes \tau$ and $\phi \simeq \psi \perp \lambda\sigma$.*

Proof. Let $b \in D(\psi) \setminus \{0\}$ be as in (\star). Then (\star) also holds for the pair $(b\phi, b\psi)$. Indeed, if this were not the case, then, for every $a \in D(\phi)$, we could find $c, d, e, f \in D(\psi)$ such that $ba = bc(b^3d + beb^2f)$, or, equivalently, such that $a = b^2c(bd + ef)$. But, since $D(\psi)$ is an F^2 -linear subspace of F , we have $b^2c \in D(\psi)$, and so this would contradict the fact that (\star) holds for the original pair (ϕ, ψ) . Since the exchange $(\phi, \psi) \rightarrow (b\phi, b\psi)$ does not affect the statement of the proposition, we can therefore assume that $b = 1$. In other words, we can assume that $1 \in D(\psi)$ and that:

- (\star') There exists an element $a \in D(\phi) \setminus \{0\}$ such that $a \neq c(d + ef)$ for any $c, d, e, f \in D(\psi)$.

Let us fix $a \in D(\phi) \setminus \{0\}$ as in (\star'). We will now prove that the statement of the proposition holds with $\lambda = a$. First, we note that (\star') implies:

- (1) $a \notin D(\psi)$.
- (2) $\psi_{F_{au}}$ is anisotropic for every $u \in D(\psi) \setminus \{0\}$.

Indeed, if a were in $D(\psi)$, then we could contradict (\star') by taking $c = 1$, $d = a$ and $e = f = 0$. Similarly, if $\psi_{F_{au}}$ were isotropic in (2), then (since ψ is anisotropic) we could find nonzero elements $x, y \in D(\psi)$ such that $au = xy$ (see Lemma 2.27 (2)); taking $c = x^{-1}$, $d = 0$, $e = y$ and $f = u$, this would again contradict (\star').

Now, (1) implies that we have $\psi \perp \langle a \rangle \subset \phi$. If $\mathfrak{i}_1(\phi) = 1$, then the latter inclusion is an isomorphism, and the statement of the proposition holds with $\pi = \sigma = \langle 1 \rangle$ and $\tau = \psi$. Assume now that $\mathfrak{i}_1(\phi) > 1$. Since $1, a \in D(\phi)$, ϕ_{F_a} is isotropic by Lemma 2.27 (2). By Lemma 4.1, it then follows that $\mathfrak{i}_0(\phi_{F_a}) \geq \mathfrak{i}_1(\phi)$. On the other hand, (2) (with $u = 1 \in D(\psi)$) shows that ψ_{F_a} is anisotropic, and, since $\dim \psi = \dim \phi - \mathfrak{i}_1(\phi)$, we conclude that $(\phi_{F_a})_{\text{an}} \simeq \psi_{F_a}$. In other words, we have $D(\phi_{F_a}) = D(\psi_{F_a}) = D(\psi) + aD(\psi)$ (where the latter equality holds by Lemma 2.27 (1)). By Lemma 2.2, it follows that we can write

$\phi \simeq \psi \perp \langle a \rangle \perp a\sigma'$ for some form $\sigma' \subset \psi$. Let $\sigma = \langle 1 \rangle \perp \sigma'$, so that $\phi \simeq \psi \perp a\sigma$. As $1 \in D(\psi)$, we have $\sigma \subset \psi$. Since σ is anisotropic and represents 1, Lemma 2.11 shows that $\sigma \subset \sigma_{\text{nor}}$. Thus, in order to complete the proof, it will be enough to prove that ψ is divisible by $\pi = \sigma_{\text{nor}}$. By Corollary 2.20, this amounts to showing that $D(\sigma) \subset G(\psi)$. Let $x \in D(\sigma) \setminus \{0\}$. In order to show that $x \in G(\psi)$, we must check that $xy \in D(\psi)$ for all $y \in D(\psi)$ (see Lemma 2.13). If $y = 0$, then there is nothing to prove. Suppose now that $y \neq 0$. Then, by Lemma 2.27 (2), $\phi_{F_{ay}}$ is isotropic. On the other hand, $\psi_{F_{ay}}$ is anisotropic by (2). Using Lemma 4.1 in the same way as before, we see that $(\phi_{F_{ay}})_{\text{an}} \simeq \psi_{F_{ay}}$, or, equivalently, that $D(\phi_{F_{ay}}) = D(\psi_{F_{ay}}) = D(\psi) + ayD(\psi)$. But, letting $u = \sqrt{ay}$, we have $xy = (a^{-1}u)^2ax \in D(\phi_{F_{ay}})$. In particular, we can find $u, v \in D(\psi)$ such that $xy = u + ayv$. To complete the proof, it now only remains to show that $v = 0$. But, if $v \neq 0$, then we have $a = v^{-1}(x + uy^{-1})$, and we obtain a contradiction to (\star') by taking $c = v^{-1}$, $d = x$, $e = u$ and $f = y^{-1}$. The result follows. \square

In general, it is *not* always possible to find a subform $\psi \subset \phi$ of codimension $i_1(\phi)$ such that the pair (ϕ, ψ) satisfies condition (\star) . Indeed, note that if $i_1(\phi) = 2$ in the situation of Proposition 4.3, then ϕ is divisible by the binary (i.e., 2-dimensional) form σ . The following example, which is directly analogous to an example of Vishik from the characteristic $\neq 2$ theory of quadratic forms, shows that there exist anisotropic quasilinear quadratic forms which have first higher isotropy index equal to 2, but which are not divisible by a binary form: Let a, b, c, d, e be algebraically independent variables over a field F_0 of characteristic 2, let $F = F_0(a, b, c, d, e)$ and consider the anisotropic F -form $\phi = \langle\langle ab, ac, ad \rangle\rangle \perp bcd\langle 1, ab, ac, ad \rangle \perp e\langle a, b, c, d \rangle$.

Lemma 4.4 (Vishik, cf. [Tot09, Lem. 7.1]). *In the above situation, we have $i_1(\phi) = 2$, but ϕ is not divisible by a binary form.*

Proof. We will freely use some basic facts from the theory of symmetric bilinear forms over fields of characteristic 2. If B is such a form, then ϕ_B will denote the totally singular quadratic form $v \mapsto B(v, v)$. For all other notation and terminology, the reader is referred to [EKM08, Ch. I].

Let \mathfrak{b} denote the bilinear form $\langle\langle ab, ac, ad \rangle\rangle_b \perp bcd\langle 1, ab, ac, ad \rangle_b \perp e\langle a, b, c, d \rangle_b$ over F , so that $\phi = \phi_{\mathfrak{b}}$. As the reader will immediately verify, we have

$$\mathfrak{b} \sim \mathfrak{c} := \langle\langle a, b, c, d \rangle\rangle_b \perp \langle\langle e \rangle\rangle_b \otimes \langle a, b, c, d \rangle_b,$$

where the symbol \sim denotes Witt equivalence. Let $\pi = \langle\langle a, b, c, d \rangle\rangle_b$, and let π' denote the pure subform of π . Since $\text{ndeg}(\phi_{\pi}) = 16 < 32 = \text{ndeg}(\phi)$, Corollary 2.32 (2) implies that $\pi_{F(\phi)}$ is anisotropic. At the same time, it follows from Remark 2.6 (4) and the definition of ϕ that $e \in D(\pi'_{F(\phi)})$. Thus, by [EKM08, Lem. 6.1], there exists a 3-fold bilinear Pfister form η over $F(\phi)$ such that $\pi_{F(\phi)} \simeq \langle\langle e \rangle\rangle_b \otimes \eta$. In particular, $\mathfrak{c}_{F(\phi)}$ is divisible by $\langle\langle e \rangle\rangle_b$, and so $i_W(\mathfrak{c}_{F(\phi)})$ is even (see [EKM08, Prop. 6.22]; i_W denotes here the Witt index). Since $\dim \mathfrak{c} - \dim \mathfrak{b} = 8 \equiv 0 \pmod{4}$, it follows that $i_W(\mathfrak{b}_{F(\phi)})$ is also even. In particular, this shows that $i_1(\phi) \geq 2$ (see [Lag07, Prop. 5.15]), and to prove that $i_1(\phi) = 2$, it suffices to find a field extension L of F such that ϕ_L is isotropic but $i_0(\phi_L) \leq 2$ (see Lemma 4.1). We claim that $L = F_{cde}$ is such an extension. First, note that since $cde = (bcd)(b^{-1}e)$ is a product of two nonzero elements of $D(\phi)$, ϕ_L is isotropic by Lemma 2.27 (2). On the other hand, it is easy to see that the codimension-2 subform $\psi = \langle\langle ab, ac, ad \rangle\rangle \perp bcd\langle 1, ab, ac, ad \rangle \perp e\langle c, d \rangle \subset \phi$ remains anisotropic over L . Indeed, since

$$\psi_L \simeq \langle\langle ab, ac, ad \rangle\rangle \perp bcd\langle 1, ab, ac, ad \rangle \perp \langle d, c \rangle \subset \langle\langle a, b, c, d \rangle\rangle_L,$$

it suffices to check that $\langle\langle a, b, c, d \rangle\rangle$ remains anisotropic over L . But, since $cde \notin F^2(a, b, c, d) = N(\langle\langle a, b, c, d \rangle\rangle)$, this follows from Lemma 2.27 (4). Since the anisotropy of ψ_L readily implies that $i_0(\phi_L) \leq 2$, we have proved our claim.

It now remains to check that ϕ is not divisible by a binary form. For the sake of contradiction, suppose instead that ϕ is divisible by $\langle\langle u \rangle\rangle$ for some $u \in F \setminus F^2$. We claim that $u \in F^2(ab, ac, ad)$. Again, let us assume that this is not the case. Then the quasi-Pfister form $\tau = \langle\langle ab, ac, ad \rangle\rangle$ remains anisotropic over F_u by Lemma 2.27 (4). Let $\sigma = \tau \perp \langle bcd \rangle \subset \phi$ and $\eta = \tau \perp \langle ae \rangle \subset \phi$. Since $i_0(\phi_{F_u}) = \frac{1}{2} \dim \phi = 8$ (Lemma 2.27 (6)), and since $\dim \sigma = \dim \eta = 9$, both σ_{F_u} and η_{F_u} are necessarily isotropic. Since τ_{F_u} is anisotropic, this means that we have $bcd, ae \in D(\tau_{F_u}) = F^2(ab, ac, ad, u)$. But this implies that $F^2(a, b, c, d, e) \subseteq F^2(ab, ac, ad, u)$, thus contradicting the fact that the elements a, b, c, d, e are algebraically independent over F_0 . This proves our claim, and so we can write $u = v + w$ for some $v \in D(\langle\langle 1, ab, ac, ad \rangle\rangle)$ and $w \in D(\langle bc, bd, cd, abcd \rangle)$. Now, Lemma 2.27 (6) implies that $u \in G(\phi)$. In particular, applying Lemma 2.13 to the elements $ae, acd, abd \in D(\phi)$, we see that

- (1) $aeu \in D(\phi)$,
- (2) $acdu \in D(\phi)$, and
- (3) $abdu \in D(\phi)$.

We can now complete the proof: First, note that since $ae\langle 1, ab, ac, ad \rangle \simeq e\langle a, b, c, d \rangle \subset \phi$, we have $aeu \in D(\phi)$. By (1), this implies that $aeu \in D(\phi)$. Note however that $ae\langle bc, bd, cd, abcd \rangle \simeq \langle abce, abde, acde, bcde \rangle$, and the latter form does not represent any nonzero element of $D(\phi)$. It follows that $w = 0$, and so $u \in D(\langle\langle 1, ab, ac, ad \rangle\rangle)$. Next, consider the form $\rho = acd\langle 1, ab, ac, ad \rangle \simeq \langle acd, bcd, c, d \rangle$. By (2), we have $acdu \in D(\rho) \cap D(\phi)$. Since the elements a, b, c, d, e are algebraically independent over F_0 , direct inspection shows that the former intersection is equal to $acdD(\langle 1, ab \rangle)$, and so $u \in D(\langle 1, ab \rangle)$. Finally, we can use (3) in a similar way to show that $u \in F^2$, thus providing us with the needed contradiction. The lemma is proved. \square

In fact, Vishik's example shows more: It is generally not possible to find a subform $\psi \subset \phi$ of codimension $i_1(\phi)$ such that the pair (ϕ, ψ) satisfies condition (\star) , *even after making arbitrary rational extensions of F* – this stronger assertion follows from Lemma 2.35. This leads us to consider the possibility that decompositions of the kind suggested by Proposition 4.3 may be found by passing to suitable transcendental extensions of the base field, and, ultimately, to our main result. Nevertheless, it is still interesting to ask for conditions on ϕ (or, more specifically, on the Knebusch splitting pattern of ϕ) which automatically ensure the existence of the needed subform ψ . To this end, it is natural to look to the extremities where ϕ is “far from generic” (e.g., where $i_1(\phi)$ is “large”). The basic example is provided here by Proposition 2.39, which characterises (scalar multiples of) anisotropic quasi-Pfister forms in terms of i_1 , and one can hope that similar characterisations exist for “sufficiently simple” forms (cf. also Theorem 6.8 below). A better understanding of all these problems would have important implications for the study of symmetric bilinear forms of “low complexity” (e.g., of “small” height) in characteristic 2.

5. MAIN THEOREM

We are now ready to give the proof of Theorem 1.5. In order to treat the case where $p > 2$, the statement needs to be modified as follows:

Theorem 5.1. *Let ϕ be an anisotropic quasilinear p -form of dimension ≥ 2 over F and let s be the smallest nonnegative integer such that $p^s \geq \mathbf{i}_1(\phi)$. If ϕ is not a quasi-Pfister p -neighbour, then ϕ_1 is divisible by an s -fold quasi-Pfister p -form.*

Remark 5.2. Nothing is lost here by assuming that ϕ is not a quasi-Pfister p -neighbour. Indeed, if ϕ is a quasi-Pfister p -neighbour, and n denotes the smallest nonnegative integer such that $p^{n+1} \geq \dim \phi$, then ϕ_1 is similar to an n -fold quasi-Pfister p -form by Corollary 3.11. If $p = 2$, then we have $\mathbf{i}_1(\phi) \leq \frac{1}{2} \dim \phi \leq 2^n$ by Proposition 2.39, so that $n \geq s$, where s is the integer defined in the statement of the theorem. Note, however, that if $p > 2$, then n may be strictly smaller than s (again, see Corollary 3.11). This explains why the additional hypothesis is needed here, but not in the statement of Theorem 1.5 (i.e., the case where $p = 2$).

Proof. To simplify the notation, we will write \mathbf{i}_1 instead of $\mathbf{i}_1(\phi)$ in what follows. If $\mathbf{i}_1 = 1$, then the statement of the theorem holds trivially. We therefore assume henceforth that $\mathbf{i}_1 > 1$. After multiplying ϕ by a nonzero scalar if necessary, we may also assume that $1 \in D(\phi)$. In particular, we can find $a_1, \dots, a_n \in F$ such that $\phi \simeq \langle 1, a_1, \dots, a_n \rangle$. For the remainder of the proof, we let $\phi' = \langle a_1, \dots, a_n \rangle$, and we write $\phi'(T)$ for the “generic value” of ϕ' , i.e., $\phi'(T) = \sum_{i=1}^n a_i T_i^p \in F[T]$, where $T = (T_1, \dots, T_n)$ is a tuple of algebraically independent variables over F . By Corollary 3.9, the function field $F(\phi)$ is F -isomorphic to $F[\phi']$, and may therefore be identified with $\text{Frac}(F[T]/(\phi'(T)))$ (see Remark 2.6 (1)). Fixing this identification henceforth, we will write \bar{f} for the image of a polynomial $f \in F[T]$ under the canonical F -algebra homomorphism $F[T] \rightarrow F(\phi)$. We will also write $m(f)$ for the multiplicity $\text{mult}_{\phi'(T)}(f)$ of $\phi'(T)$ in f , i.e., the largest integer k such that $f = \phi'(T)^k h$ for some $h \in F[T]$. Note here that we have $\bar{f} \neq 0$ if and only if $m(f) = 0$.

Now, let $\psi \subset \phi$ be any subform of codimension $\mathbf{i}_1(\phi)$ such that $1 \in D(\psi)$. We then have the following lemma:

Lemma 5.3. *In the above situation, we can find elements $g_{i,j} \in D(\psi_{F[T]})$ ($1 \leq i < \mathbf{i}_1$, $1 \leq j < p$) such that*

$$\phi_{F(T)} \simeq \psi_{F(T)} \oplus \phi'(T) \langle 1, f_1, \dots, f_{\mathbf{i}_1-1} \rangle,$$

where, for each $1 \leq i < \mathbf{i}_1$, $f_i = \sum_{j=1}^{p-1} g_{i,j} \phi'(T)^{j-1}$.

Proof. First, let us note that $\phi'(T) \notin D(\psi_{F(T)})$. Indeed, if $\psi_{F(T)}$ were to represent $\phi'(T)$, then it would follow from Theorem 2.23 that $a_1, \dots, a_n \in D(\psi)$. Since $1 \in D(\psi)$ by hypothesis, this would imply that $D(\phi) \subseteq D(\psi)$, or, equivalently, that $\phi \subset \psi$ (see Proposition 2.3), which is impossible for dimension reasons (recall here that $\mathbf{i}_1 > 1$ by assumption). It follows that $\psi_{F(T)} \oplus \langle \phi'(T) \rangle$ is anisotropic, and so $\psi_{F(T)} \oplus \langle \phi'(T) \rangle \subset \phi_{F(T)}$ by Proposition 2.3. Now, the affine function field $F[\phi]$ may be identified (over F) with the field $K = F(T)_{\phi'(T)}$ (see Remark 2.6 (1)). Since $F[\phi]$ is F -isomorphic to a purely transcendental extension of $F(\phi)$ (Remark 2.6 (2)), Corollary 3.4 and Lemma 2.26 together imply that $\psi_K \simeq (\phi_K)_{\text{an}}$. In particular, we have $D(\phi_{F(T)}) \subset D(\phi_K) = D(\psi_K) = \sum_{j=0}^{p-1} D(\psi_{F(T)}) \phi'(T)^j$ (where the last equality holds by Lemma 2.27 (1)). Thus, by Lemma 2.2, we can complete the subform inclusion $\psi_{F(T)} \oplus \langle \phi'(T) \rangle \subset \phi_{F(T)}$ to an isomorphism $\phi_{F(T)} \simeq \psi_{F(T)} \oplus \langle \phi'(T) \rangle \oplus \langle f'_1, \dots, f'_{\mathbf{i}_1-1} \rangle$, where, for each i , we have $f'_i = \sum_{j=0}^{p-1} g_{i,j} \phi'(T)^j$ for some $g_{i,j} \in D(\psi_{F(T)})$. Note, however, that every element of $D(\psi_{F(T)})$ is (trivially) the ratio of an element of $D(\psi_{F[T]})$ and a p -th power in $F[T]$. Since multiplying the f'_i by p -th powers in $F[T]$ does not change the $F(T)$ -form $\langle f'_1, \dots, f'_{\mathbf{i}_1-1} \rangle$ up to isomorphism (see Lemma 2.2), we can arrange it so that the $g_{i,j}$ belong to $D(\psi_{F[T]})$. Similarly, since subtracting elements of $D(\psi_{F(T)})$ from the f'_i does not change the isomorphism class of

$\psi_{F(T)} \oplus \langle f'_1, \dots, f'_{i_1-1} \rangle$ (again, see Lemma 2.2), we can also arrange it so that $g_{i,0} = 0$ for all i . The remaining $g_{i,j}$ then satisfy the statement of the lemma. \square

Let us now fix elements $g_{i,j} \in D(\psi_{F[T]})$ (and the associated polynomials f_i) satisfying the statement of Lemma 5.3. We are searching here for a sufficiently large quasi-Pfister divisor of ϕ_1 , and we would like to try to build this quasi-Pfister p -form from the elements $g_{i,1}$. The basic point here is the following:

Lemma 5.4. *In the above situation, we have $\overline{g_{i,1}}b \in D(\phi_1)$ for all $b \in D(\psi)$ and all $1 \leq i < i_1$.*

Proof. If $b = 0$, then the statement is trivial. Let us now fix $b \in D(\psi) \setminus \{0\}$. We will need another lemma:

Lemma 5.5. *In the above situation, there exist elements $s_{i,j} \in D(\psi_{F[T]})$ and $t_{i,j} \in F[T] \setminus \{0\}$ ($1 \leq i < i_1$, $0 \leq j < p$) such that, for every $1 \leq i < i_1$, we have:*

- (1) $bf_i = \sum_{j=0}^{p-1} \frac{s_{i,j}}{t_{i,j}^p} \frac{\phi'(T)^j}{b^j}$ in $F(T)$.
- (2) For each $0 \leq j < p$, at least one of $\overline{s_{i,j}}$ and $\overline{t_{i,j}}$ is nonzero.

Proof. Let $1 \leq i < i_1$, and consider the field $L = F(T)_u$, where $u = \frac{\phi'(T)}{b}$. In view of Remark 2.6 (1), L is F -isomorphic to the affine function field $F[\eta]$, where η denotes the F -form $\langle b \rangle \perp \phi'$. Now, since $b \in D(\psi)$, and since $\psi \subset \phi$, we have $D(\phi') \subseteq D(\eta) \subseteq D(\phi)$. In particular, if $\eta \not\subseteq \phi$, then it follows from Lemma 2.2 that $\eta_{\text{an}} \simeq \phi'$. Either way, we see that L is F -isomorphic to a degree-1 purely transcendental extension of $F(\phi)$ – in the first case, see Remark 2.6 (2); in the second, see Remark 2.6 (3) and Corollary 3.9. By Corollary 3.4 and Lemma 2.26, it follows that $\psi_L \simeq (\phi_L)_{\text{an}}$. In other words, we have $D(\phi_L) = D(\psi_L) = \sum_{j=0}^{p-1} D(\psi_{F(T)})u^j$ (again, see Lemma 2.27 (1) for the final equality). Now, since u is a p -th power in L , we have $bf_i = \frac{\phi'(T)f_i}{u} \in D(\phi_L)$. We can therefore write $bf_i = \sum_{i=0}^{p-1} q_j u^j$ for some $q_j \in D(\psi_{F(T)})$. Since every element of $D(\psi_{F(T)})$ is the quotient of an element of $D(\psi_{F[T]})$ and a p -th power in $F[T]$, and since $u = \frac{\phi'(T)}{b}$, this shows that we can find elements $s_{i,j} \in D(\psi_{F[T]})$ and $t_{i,j} \in F[T] \setminus \{0\}$ such that (1) holds. Finally, since $\psi_{F(\phi)}$ is anisotropic, Proposition 2.33 implies that $m(s_{i,j}) \equiv 0 \pmod{p}$ for all $0 \leq j < p$. For each such j , let $m_j = \min(m(s_{i,j}), pm(t_{i,j}))$, and put $s'_{i,j} = \frac{s_{i,j}}{\phi'(T)^{m_j}}$ and $t'_{i,j} = \frac{t_{i,j}}{\phi'(T)^{m_j/p}}$. Then, by Theorem 2.23, we again have $s'_{i,j} \in D(\psi_{F[T]})$. Thus, replacing $s_{i,j}$ by $s'_{i,j}$ and $t_{i,j}$ by $t'_{i,j}$ (for each j), we arrive at the situation where, for any j , either $m(s_{i,j}) = 0$ or $m(t_{i,j}) = 0$. In other words at least one of $\overline{s_{i,j}}$ and $\overline{t_{i,j}}$ is nonzero, as we wanted. \square

Returning now to the proof of Lemma 5.4, let $s_{i,j} \in D(\psi_{F[T]})$ and $t_{i,j} \in F[T] \setminus \{0\}$ be as in Lemma 5.5. In particular, we have the equation

$$\sum_{l=1}^{p-1} bg_{i,l} \phi'(T)^{l-1} = bf_i = \sum_{j=0}^{p-1} \frac{s_{i,j}}{t_{i,j}^p} \frac{\phi'(T)^j}{b^j}$$

in $F(T)$. Clearing denominators, we obtain

$$(5.1) \quad \prod_k t_{i,k}^p \sum_{l=1}^{p-1} bg_{i,l} \phi'(T)^{l-1} = \sum_{j=0}^{p-1} \prod_{k \neq j} t_{i,k}^p s_{i,j} \frac{\phi'(T)^j}{b^j}$$

Now, we claim that, for all $0 \leq j < p$, we have $\overline{t_{i,j}} \neq 0$, or, equivalently, $m(t_{i,j}) = 0$. To see this, let $m = \min\{\sum_{k \neq j} m(t_{i,k}) \mid 0 \leq j < p\}$. Then our claim amounts to the assertion

that $m = \sum_{k=0}^{p-1} m(t_{i,k})$. Suppose that this is not the case, and let $0 \leq j < p$ be minimal so that $\sum_{k \neq j} m(t_{i,k}) = m$. Then, reducing both sides of (5.1) modulo $\phi'(T)^{pm+j+1}$, we see that $s_{i,j} \equiv 0 \pmod{\phi'(T)}$. In other words, we have $\overline{s_{i,j}} = 0$. By the choice of the $s_{i,j}$ and $t_{i,j}$, this implies that $\overline{t_{i,j}} \neq 0$, or, equivalently, that $m(t_{i,j}) = 0$. But then $m = \sum_{k \neq j} m(t_{i,k}) = \sum_{k=0}^{p-1} m(t_{i,k})$, which contradicts our assumption. The claim is therefore proved, and so, reducing (5.1) modulo $\phi'(T)$ and dividing through by $\prod_k \overline{t_{i,k}}^p$, we obtain the equality $\overline{g_{i,1}}b = \overline{s_{i,0}}/\overline{t_{i,0}}^p$ in $F(\phi)$. As $s_{i,0} \in D(\psi_{F[T]})$, this shows that $\overline{g_{i,1}}b \in D(\psi_{F(\phi)})$. But since $\psi_{F(\phi)} \simeq \phi_1$, we have $D(\psi_{F(\phi)}) = D(\phi_1)$, and the lemma is therefore proved. \square

Continuing with the proof of Theorem 5.1, let us now choose elements $g_{i,j} \in D(\psi_{F[T]})$ as in the statement of Lemma 5.3 so that the integer $\sum_{i=1}^{i_1-1} \deg_{T_1}(g_{i,1})$ is minimal, where, for any $g \in F[T]$, $\deg_{T_1}(g)$ denotes the degree of g viewed as an element of the ring $F(T_2, \dots, T_n)[T_1]$, i.e., as a polynomial in the single variable T_1 (with the added convention that $\deg_{T_1}(0) = 0$). Consider the form $\sigma = \langle 1, \overline{g_{1,1}}, \dots, \overline{g_{i_1-1,1}} \rangle$ over $F(\phi)$. The final step in the proof of the theorem will be to prove the following statement:

Lemma 5.6. *In the above situation, σ is anisotropic.*

Before proving the lemma, let us explain how this concludes the proof of Theorem 5.1. First, we claim that ϕ_1 is divisible by the quasi-Pfister p -form σ_{nor} . By Corollary 2.20, this amounts to checking that $D(\sigma) \subseteq G(\phi_1)$. Since $G(\phi_1)$ is a subfield of F containing F^p (Corollary 2.15), it suffices to show here that $\overline{g_{i,1}} \in G(\phi_1)$ for all $1 \leq i < i_1$. But by Lemma 2.13, this is equivalent to showing that, for all such i , we have $\overline{g_{i,1}}D(\phi_1) \subseteq D(\phi_1)$. Since $D(\phi_1) = D(\psi_{F(\phi)})$ is spanned as an $F(\phi)^p$ -vector space by $D(\psi)$, this follows immediately from Lemma 5.4. The claim is therefore proved, and to finish the proof of the theorem, it only remains to check that $\dim \sigma_{\text{nor}} \geq p^s$. But, σ is anisotropic by Lemma 5.6, and so $\sigma \subset \sigma_{\text{nor}}$ by Lemma 2.11. In particular, we have $\dim \sigma_{\text{nor}} \geq \dim \sigma = i_1$, which is precisely the assertion that $\dim \sigma_{\text{nor}} \geq p^s$ (because $\dim \sigma_{\text{nor}}$ is necessarily a power of p). Now, in order to prove Lemma 5.6, we need another auxiliary statement:

Lemma 5.7. *If σ is isotropic, then $p > 2$, and there exist polynomials $g_j \in D(\psi_{F[T]})$ ($1 \leq j < p$) and an integer $2 \leq k < p$ such that*

- (1) $\sum_{j=1}^{p-1} g_j \phi'(T)^j \in D(\psi_{F[T]})$.
- (2) $m(g_l) > 0$ for all $1 \leq l < k$.
- (3) $m(g_k) = 0$.

Proof. If σ is isotropic, then since $\phi'(T)$ is a Fermat-type polynomial of degree p , we can find an integer $1 \leq m \leq p$ and polynomials $h_0, \dots, h_{i_1-1}, h \in F[T]$ such that:

- (i) $h_0^p + g_{1,1}h_1^p + \dots + g_{i_1-1,1}h_{i_1-1}^p = \phi'(T)^m h$ in $F[T]$.
- (ii) $\deg_{T_1}(h_i) < p$ for all $0 \leq i < i_1$.
- (iii) $\overline{h_i} \neq 0$ for some $1 \leq i < i_1$.

First, let us note that we have $\phi'(T)^m h \in D(\psi_{F[T]})$ by (i) and the definition of the elements $g_{i,1}$. Since $\psi_{F(\phi)}$ is anisotropic, it follows from Proposition 2.33 that $h = 0$ or $m = p$. Either way, we can assume henceforth that $m = p$. Now, by (iii), there exists $1 \leq l < i_1$ such that $h_l \neq 0$. Among all such integers l , let us fix one so that $\deg_{T_1}(g_{l,1}h_l^p)$ is maximal. Consider now the polynomial $f = h_0^p + \sum_{k=1}^{i_1-1} f_k h_k^p \in F[T]$, where the f_k are as in the statement of Lemma 5.3. Since $h_l \neq 0$, Lemma 2.2 implies that $\langle 1, f_1, \dots, f_{i_1-1} \rangle \simeq \langle 1, f_1, \dots, f_{l-1}, f, f_{l+1}, \dots, f_{i_1-1} \rangle$ as $F(T)$ -forms. In particular, we

have

$$(5.2) \quad \phi_{F(T)} \simeq \psi_{F(T)} \oplus \phi'(T)\langle 1, f_1, \dots, f_{l-1}, f, f_{l+1}, \dots, f_{i_1-1} \rangle.$$

Now, by definition, $f = \sum_{j=1}^{p-1} g'_j \phi'(T)^{j-1}$, where $g'_1 = \phi'(T)^m h \in D(\psi_{F(T)})$ and $g'_j = \sum_{k=1}^{i_1-1} g_{k,j} h_k^p \in D(\psi_{F(T)})$ for all $2 \leq j < p$. Let $r = \min\{m(g'_j) \mid 1 \leq j < p\}$. Since $g'_j \in D(\psi_{F(T)})$ for all j , and since $\psi_{F(\phi)}$ is anisotropic, another application of Proposition 2.33 shows that $r \equiv 0 \pmod{p}$. In particular, for each $j \geq 1$, we have $g_j := \frac{g'_j}{\phi'(T)^r} \in D(\psi_{F(T)})$. In view of (5.2), it follows that the exchange $g_{l,j} \rightarrow g_j$ does not alter the statement of Lemma 5.3. By our choice of the $g_{i,j}$, we therefore have

$$(5.3) \quad \deg_{T_1}(\phi'(T)^{p-r} h) = \deg_{T_1}(g_1) \geq \deg_{T_1}(g_{l,1}).$$

Now, we claim that the elements g_j (together with an appropriate integer k) satisfy the conditions of the lemma. We have already seen here the validity of (1). At the same time, we have $m(g_j) = 0$ for some $j \geq 1$ by construction. Thus, in order to prove the existence of an integer k such that (2) and (3) are satisfied, we just need to check that $m(g_1) > 0$. Recall again that we have $g_1 = \phi'(T)^{p-r} h$. If $h = 0$, then there is nothing to prove. Suppose now that $h \neq 0$. By (i) and the choice of the integer l , we have $\deg_{T_1}(g_{l,1} h_l^p) \geq \deg_{T_1}(\phi'(T)^p h) \geq p^2 + \deg_{T_1}(h)$. Since $\deg_{T_1}(h_l) < p$ (by (ii)), it follows that $\deg_{T_1}(g_{l,1}) > \deg_{T_1}(h)$. In view of (5.3), we see that $r = 0$ in this case. In particular, we have $g_1 = \phi'(T)^p h$, and so $m(g_1) \geq p > 0$, as we wanted. \square

We are now ready to prove Lemma 5.6 and thus complete the proof of Theorem 5.1. If $p = 2$, then the statement was already proved in Lemma 5.7. Suppose now that $p > 2$, and assume for the sake of contradiction that σ is isotropic. Let g_j ($1 \leq j < p$) and k be as in the statement of Lemma 5.7. By condition (2) of the lemma, we can, for each $l < k$, write $g_l = \phi'(T)^{m_l} h_l$ for some positive integer m_l and some polynomial h_l . By a now familiar application of Proposition 2.33, we have $m_l \equiv 0 \pmod{p}$ for every such l . In particular, the m_l are all strictly larger than k . Now, by condition (1) of the lemma, the element

$$\phi'(T)^k (g_k + g_{k-1} \phi'(T) + \dots + g_{i_1-1} \phi'(T)^{p-k-1} + h_1 \phi'(T)^{m_1-k} + \dots + h_{k-1} \phi'(T)^{m_{k-1}-k})$$

lies in $D(\phi_{F(T)})$. Using the very same argument as that used to prove Lemma 5.4 above (and the fact that the integers $m_l - k$ ($l < k$) are all positive), one readily shows that $b^k \overline{g_k} \in D(\phi_1)$ for every $b \in D(\psi)$. Note, however, that $\overline{g_k} \neq 0$ by condition (3) of Lemma 5.7. Since $2 \leq k < p$, and since $\phi_1 \simeq \psi_{F(\phi)}$, it follows from Lemma 2.22 that ϕ_1 is a quasi-Pfister p -form. But, by Corollary 3.11, this in turn implies that ϕ is a quasi-Pfister p -neighbour, thus contradicting our original hypothesis. The lemma and theorem are therefore proved. \square

6. FIRST APPLICATIONS OF THE MAIN THEOREM

We now give the basic applications of Theorem 1.5.

6.1. Possible values of the Knebusch splitting pattern. Let ϕ be a quasilinear p -form of dimension ≥ 2 over F . In the previous section we have shown that the first higher anisotropic kernel ϕ_1 of ϕ is divisible by a quasi-Pfister p -form of dimension $\geq i_1(\phi)$, provided that ϕ_{an} is not a quasi-Pfister p -neighbour. In the terminology of §2.11, this amounts to the assertion that if $h_{\text{qp}}(\phi) \geq 2$, then $\mathfrak{d}_1(\phi) \geq \log_p(i_1(\phi))$ (here we are also making use Corollary 3.11). By virtue of the inductive nature of the Knebusch splitting tower construction, we also obtain analogous restrictions on the higher isotropy indices

$\mathbf{i}_r(\phi)$ ($2 \leq r < h_{\text{qp}}(\phi)$) in terms of the corresponding higher divisibility indices $\mathfrak{d}_r(\phi)$. Taking the observations of §2.11 into account, our results may be summarised as follows:

Theorem 6.1. *Let ϕ be a quasilinear p -form over F and let $d = h(\phi) - h_{\text{qp}}(\phi)$. Then:*

- (1) $\mathbf{i}(\phi) = (\mathbf{i}_1(\phi), \dots, \mathbf{i}_{h_{\text{qp}}(\phi)}(\phi), p^d - p^{d-1}, p^{d-1} - p^{d-2}, \dots, p^2 - p, p - 1)$.
- (2) $\mathbf{i}_{h_{\text{qp}}(\phi)}(\phi) = \dim \phi - \mathbf{j}_{h_{\text{qp}}(\phi)-1}(\phi) - p^d < p^{d+1} - p^d$.
- (3) $\mathfrak{d}(\phi) = (\mathfrak{d}_0(\phi), \dots, \mathfrak{d}_{h_{\text{qp}}(\phi)-1}(\phi), d, d - 1, \dots, 1, 0)$.
- (4) $\mathfrak{d}_0(\phi) \leq \mathfrak{d}_1(\phi) \leq \dots \leq \mathfrak{d}_{h_{\text{qp}}(\phi)}(\phi) = d$.
- (5) $\mathbf{i}_r(\phi) \equiv 0 \pmod{p^{\mathfrak{d}_{r-1}(\phi)}}$ for all $1 \leq r < h_{\text{qp}}(\phi)$.
- (6) $\mathfrak{d}_r(\phi) \geq \log_p(\mathbf{i}_r(\phi))$ for all $1 \leq r < h_{\text{qp}}(\phi)$.
- (7) For every $1 \leq r < h_{\text{qp}}(\phi)$, $\mathbf{i}_r(\phi) - 1$ is the remainder of $\dim \phi - \mathbf{j}_{r-1}(\phi) - 1$ modulo $p^{\mathfrak{d}_r(\phi)}$.

Proof. Parts (1), (2), (3), (4) and (5) are the statements comprising Lemmas 2.42, 2.43, 2.44 and Corollary 2.45. Since $\mathbf{i}_r(\phi) = \mathbf{i}_1(\phi_{r-1})$, $\mathfrak{d}_r(\phi) = \mathfrak{d}_1(\phi_{r-1})$ and $\dim \phi_{r-1} = \dim \phi - \mathbf{j}_{r-1}(\phi)$ for all $1 \leq r \leq h(\phi)$, parts (6) and (7) follow immediately from Theorem 5.1 and Corollary 3.11. \square

In order to highlight the general shape of the Knebusch splitting pattern exposed by Theorem 6.1, it is worth writing down the following result explicitly (here the notation $a \mid b$ means that a divides b):

Corollary 6.2. *Let ϕ be a quasilinear p -form over F . Then $\mathbf{i}_1(\phi) \leq p^{\mathfrak{d}_1(\phi)} \mid \mathbf{i}_2(\phi) \leq p^{\mathfrak{d}_2(\phi)} \mid \dots \mid \mathbf{i}_{h_{\text{qp}}(\phi)-1}(\phi) \leq p^{\mathfrak{d}_{h_{\text{qp}}(\phi)-1}(\phi)} \mid \mathbf{i}_{h_{\text{qp}}(\phi)}(\phi)$.*

Remark 6.3. In the special case where $p = 2$, the chain of inequalities $\mathbf{i}_1(\phi) \leq \mathbf{i}_2(\phi) \leq \dots \leq \mathbf{i}_{h_{\text{qp}}(\phi)}(\phi)$ was previously obtained in [Scu14, Thm. 9.5] using Proposition 3.16. Here, we have given a more precise and natural explanation of this phenomenon.

We now show that, as far as the Knebusch splitting pattern is concerned, one cannot do any better than Theorem 6.1 in general:

Proposition 6.4. *Let n be any positive integer. Suppose that we are given a nonnegative integer $k \leq n$ and two sequences $(\mathfrak{d}_0, \mathfrak{d}_1, \dots, \mathfrak{d}_k = d)$ and $(\mathbf{i}_1, \dots, \mathbf{i}_k)$ of $k + 1$ and k nonnegative integers, respectively, such that the following conditions hold:*

- (i) $\mathbf{i}_k = n - \sum_{j=1}^{k-1} \mathbf{i}_j - p^d < p^{d+1} - p^d$.
- (ii) $\mathfrak{d}_0 \leq \mathfrak{d}_1 \leq \dots \leq \mathfrak{d}_k = d$.
- (iii) $1 \leq \mathbf{i}_r \equiv 0 \pmod{p^{\mathfrak{d}_{r-1}}}$ for all $1 \leq r < k$.
- (iv) $\mathfrak{d}_r \geq \log_p(\mathbf{i}_r)$ for all $1 \leq r < k$.
- (v) For every $1 \leq r < k$, $\mathbf{i}_r - 1$ is the remainder of $n - (\sum_{j=1}^{r-1} \mathbf{i}_j) - 1$ modulo $p^{\mathfrak{d}_r}$.

Then there exists a (purely transcendental) field extension L of F and an anisotropic quasilinear p -form ϕ of dimension n over L such that:

- (1) $h_{\text{qp}}(\phi) = k$.
- (2) $h(\phi) = k + d$.
- (3) $\mathfrak{d}(\phi) = (\mathfrak{d}_0(\phi), \dots, \mathfrak{d}_{k-1}(\phi), d, d - 1, \dots, 1, 0)$.
- (4) $\mathfrak{d}_r(\phi) \geq \mathfrak{d}_r$ for all $0 \leq r < k$.
- (5) $\mathbf{i}(\phi) = (\mathbf{i}_1, \dots, \mathbf{i}_k, p^d - p^{d-1}, p^{d-1} - p^{d-2}, \dots, p^2 - p, p - 1)$.

Proof. We argue by induction on k . If $k = 0$, then Proposition 2.39 shows that we can take $L = F(T)$ and $\phi = \langle\langle T_1, \dots, T_d \rangle\rangle$, where $T = (T_1, \dots, T_d)$ is a d -tuple of algebraically independent variables over F . Suppose now that $k > 0$, and let $n' = \frac{n - \mathbf{i}_1}{p^{\mathfrak{d}_1}}$ and $\mathbf{i}'_r = \frac{\mathbf{i}_r + 1}{p^{\mathfrak{d}_1}}$

for all $1 \leq r < k$. By our hypotheses, these ratios are, in fact, positive integers. Setting $\mathfrak{d}'_r = \mathfrak{d}_{r+1} - \mathfrak{d}_1$ for all $0 \leq r < k$, and putting $d' = \mathfrak{d}'_{k-1}$, conditions (i)-(v) then imply the following:

- (i') $i'_{k-1} = n' - \sum_{j=1}^{k-2} i'_j - p^{d'} < p^{d'+1} - p^{d'}$.
- (ii') $\mathfrak{d}'_0 \leq \mathfrak{d}'_1 \leq \dots \leq \mathfrak{d}'_{k-1} = d'$.
- (iii') $1 \leq i'_r \equiv 0 \pmod{p^{d'_r-1}}$ for all $1 \leq r < k-1$.
- (iv') $\mathfrak{d}'_r \geq \log_p(i'_r)$ for all $1 \leq r < k-1$.
- (v') For every $1 \leq r < k-1$, $i'_r - 1$ is the remainder of $n' - (\sum_{j=1}^{r-1} i'_j) - 1$ modulo $p^{d'_r}$.

By the induction hypothesis, there exists a (purely transcendental) field extension L_0 of F and an anisotropic quasilinear p -form ψ of dimension n' over L_0 such that

- (1') $h_{\text{qp}}(\psi) = k - 1$.
- (2') $h(\psi) = k - 1 + d'$.
- (3') $\mathfrak{d}(\psi) = (\mathfrak{d}_0(\psi), \mathfrak{d}_1(\psi), \dots, \mathfrak{d}_{k-2}(\psi), d', d' - 1, \dots, 1, 0)$.
- (4') $\mathfrak{d}_r(\psi) \geq \mathfrak{d}'_r$ for all $0 \leq r < k-1$.
- (5') $\mathfrak{i}(\psi) = (i'_1, \dots, i'_{k-1}, p^{d'} - p^{d'-1}, p^{d'-1} - p^{d'-2}, \dots, p^2 - p, p - 1)$.

Consider now the form $\sigma = \psi_{L_1} \perp \langle T_0 \rangle$ over the rational function field $L_1 = L_0(T_0)$. By (3'), (5') and Lemma 2.46, we have

- (a) $\mathfrak{d}(\sigma) = (0, \mathfrak{d}_0(\psi), \mathfrak{d}_1(\psi), \dots, \mathfrak{d}_{k-2}(\psi), d', d' - 1, \dots, 1, 0)$.
- (b) $\mathfrak{i}(\sigma) = (1, i'_1, \dots, i'_{k-1}, p^{d'} - p^{d'-1}, p^{d'-1} - p^{d'-2}, \dots, p^2 - p, p - 1)$.

We would like to modify this further. Consider next the product $\tau = \langle\langle T_1, \dots, T_{\mathfrak{d}_1} \rangle\rangle \otimes \sigma_L$ over $L = L_1(T)$, where $T = (T_1, \dots, T_{\mathfrak{d}_1})$ is a \mathfrak{d}_1 -tuple of algebraically independent variables over L_1 . Then, by (a), (b) and Lemma 2.48, we have

- (c) $\mathfrak{d}(\tau) = (\mathfrak{d}_1, \mathfrak{d}_1(\psi) + \mathfrak{d}_1, \mathfrak{d}_2(\psi) + \mathfrak{d}_1, \dots, \mathfrak{d}_{k-2}(\psi) + \mathfrak{d}_1, d, d - 1, \dots, 1, 0)$.
- (d) $\mathfrak{i}(\tau) = (p^{\mathfrak{d}_1}, i_2, \dots, i_k, p^d - p^{d-1}, p^{d-1} - p^{d-2}, \dots, p^2 - p, p - 1)$.

Now, by (iv), we have $\mathfrak{i}_1 = p^{\mathfrak{d}_1} - s$ for some $0 \leq s < p^{\mathfrak{d}_1}$. By (ii) and (iii), s is divisible by $p^{\mathfrak{d}_0}$. Let ϕ be any codimension- s subform of τ which is divisible by $\langle\langle T_1, \dots, T_{\mathfrak{d}_0} \rangle\rangle$. Clearly ϕ is anisotropic, and by (c), (d) and Proposition 3.10, we have

- (e) $\mathfrak{d}(\phi) = (\mathfrak{d}_0(\phi), \mathfrak{d}_1(\psi) + \mathfrak{d}_1, \dots, \mathfrak{d}_{k-2}(\psi) + \mathfrak{d}_1, d, d - 1, \dots, 1, 0)$.
- (f) $\mathfrak{i}(\phi) = (i_1, i_2, \dots, i_k, p^d - p^{d-1}, p^{d-1} - p^{d-2}, \dots, p^2 - p, p - 1)$.

The second statement shows that ϕ satisfies conditions (2) and (5). At the same time, since $\mathfrak{d}_0(\phi) \geq \mathfrak{d}_0$ by construction, and since $\mathfrak{d}_2(\psi) + \mathfrak{d}_1 \geq \mathfrak{d}'_r + \mathfrak{d}_1 = \mathfrak{d}_{r+1}$ for all $0 \leq r < k-1$ by (4'), (e) shows that (3) and (4) are also satisfied. Finally, since $\mathfrak{i}_k < p^{d'+1} - p^d$, Proposition 2.39 shows that $h_{\text{qp}}(\phi) = k$, i.e., that (1) holds for ϕ . The pair (L, ϕ) therefore has all the desired properties. \square

Remark 6.5. In general, it is not possible to arrange it so that $\mathfrak{d}_r(\phi) = \mathfrak{d}_r$ for all $0 \leq r < k$ in the statement of Proposition 6.4. For example, suppose that $p = 2$, and take $n = 2^{s+1} - 2$ for some $s \geq 3$, $k = 1$, $\mathfrak{d}_0 = 0$, $\mathfrak{d}_1 = s$, $\mathfrak{i}_0 = 0$, $\mathfrak{i}_1 = 2^s - 2$. As the reader will readily verify, these integers satisfy conditions (i)-(v) of the Proposition. On the other hand, let (L, ϕ) be any pair consisting of a field extension L of F and an anisotropic form ϕ of dimension $2^{s+1} - 2$ over L such that $\mathfrak{i}_1(\phi) = 2^s - 2$. By Theorem 6.8 below (see also [Scu14, Thm. 9.6]), ϕ is necessarily a quasi-Pfister 2-neighbour, and therefore satisfies conditions (1)-(5) of the proposition (see Corollary 3.11). We claim, however, that $\mathfrak{d}_0(\phi) > 0$, i.e., that ϕ is divisible by a binary form. To see this, note that there exists an anisotropic $(s+1)$ -fold bilinear Pfister form π over L and a subform $\mathfrak{b} \subset \pi$ such that ϕ is given by the assignment $v \mapsto \mathfrak{b}(v, v)$. Since \mathfrak{b} (being a codimension-2 subform of π) becomes split over the extension

$K = L_{\det(\mathbf{b})}$ (where \det denotes the determinant), we have $i_0(\phi_K) \geq \frac{1}{2} \dim \phi$ (see [Lag07, Prop. 5.15]), whence ϕ is divisible by $\langle\langle \det(\mathbf{b}) \rangle\rangle$ (Lemma 2.27 (6)).

Theorem 6.1 and Proposition 6.4 thus give a complete solution to the problem of determining the possible values of the Knebusch splitting pattern for quasilinear p -forms. In particular, we have an answer to Question 1.1 in the totally singular case. As noted in Example 2.47, the Knebusch and full splitting patterns need not agree in general for quasilinear p -forms. In §7 below, we will consider the problem of determining the possible values of the full splitting pattern in the case where $p = 2$.

6.2. Canonical dimensions of quasilinear p -hypersurfaces. Since the canonical dimension of an anisotropic quasilinear p -hypersurface is determined by the Knebusch splitting pattern of its underlying form (this is the basic fact underlying our proof of Theorem 5.1, see Theorem 3.1 and Remark 3.2 (2)), we also obtain a list of all restrictions on the possible values of the former invariant:

Theorem 6.6. *Let X be an anisotropic quasilinear p -hypersurface over F . Then there exists a nonnegative integer s such that*

- (1) $p^s - 1 \leq \dim X < p^s + \text{cdim}(X)$.
- (2) $\text{cdim}(X) \equiv -1 \pmod{p^s}$.

There are no further restrictions on $\text{cdim}(X)$.

Proof. As per the above discussion, this follows readily from Theorem 3.1, Theorem 6.1 and Proposition 6.4 (see also Remark 3.2 (2)). \square

Remarks 6.7. (1) We emphasise again that $\text{cdim}(X)$ is to be understood here as the minimum dimension of the image of a rational self-map $X \dashrightarrow X$. For $p > 3$, we do not know if this agrees with the definition given in [EKM08, §90] (although this is certainly expected – cf. [Scu14, Q. 4.4, Prop. 4.7]).

(2) The problem of determining all possible values of the canonical dimension for smooth (projective) hypersurfaces X of prime degree $p > 2$ in characteristic $\neq p$ remains open in general. Using a degree formula, Merkurjev (cf. [Mer03, §7.3]) showed that if $\dim X \geq p^n - 1$ for some nonnegative integer n , then we also have $\text{cdim}(X) \geq p^n - 1$ provided that X has no points of degree *prime to* p . Little else is known. By way of specialization, our Theorem 6.6 may be of some use when considering specific examples over certain fields of characteristic zero.

6.3. Quasilinear p -forms with maximal splitting. Let ϕ be an anisotropic quasilinear p -form of dimension ≥ 2 over F and write $\dim \phi = p^n + m$ for uniquely determined integers $n \geq 0$ and $1 \leq m \leq p^{n+1} - p^n$. By Theorem 6.1 (see also [Scu13, Cor. 6.8]), we have $i_1(\phi) \leq m$. If equality holds here, then we say that ϕ has *maximal splitting*. The basic examples of forms having this property are given by anisotropic quasi-Pfister p -neighbours (see Corollary 3.11). It is interesting to ask here to what extent this property characterises quasi-Pfister p -neighbours. Given Theorem 5.1, we can now prove the following general result:

Theorem 6.8. *Let ϕ be an anisotropic quasilinear p -form of dimension ≥ 2 over F , and n be the smallest nonnegative integer such that $p^{n+1} \geq \dim \phi$. If ϕ has maximal splitting, and if either*

- (1) $p > 2$ and $\dim \phi > p^n + p^{n-1}$, or
- (2) $p = 2$ and $\dim \phi > 2^n + 2^{n-2}$,

then ϕ is a quasi-Pfister p -neighbour.

Proof. By Theorem 5.1, we may assume that $\mathfrak{d}_1(\phi) \geq \mathfrak{i}_1(\phi)$. Suppose first that $p > 2$. Since ϕ has maximal splitting, we have $\mathfrak{i}_1(\phi) > p^{n-1}$ by (1), and so $\mathfrak{d}_1(\phi) \geq p^n$. On the other hand, we have $\dim \phi_1 = \dim \phi - \mathfrak{i}_1(\phi) = p^n$ (see Remark 2.36 (2)). It follows that ϕ_1 is similar to an n -fold quasi-Pfister p -form, and so ϕ is a quasi-Pfister p -neighbour by Corollary 3.11. If $p = 2$, the same argument (and (2)) shows that ϕ_1 is a form of dimension 2^n which is divisible by an $(n-1)$ -fold quasi-Pfister 2-form. Since every binary form is similar to a quasi-Pfister 2-form in this case, ϕ_1 is, in fact, similar to an n -fold quasi-Pfister 2-form, and we now conclude as before. \square

Remark 6.9. The statement of Theorem 6.8 was originally conjectured by Hoffmann in [Hof04, Rem. 7.32] (see also [Scu14, Que. 7.6]). The result is the best possible, in the sense that one has examples of anisotropic quasilinear p -forms with maximal splitting which are not quasi-Pfister p -neighbours in every dimension omitted in the statement of the theorem (see [Hof04, Ex. 7.31]). In the special case where $p = 2$, Theorem 6.8 was previously established in [Scu14, Thm. 9.6] using Proposition 3.16. Here, we obtain a more natural explanation of this phenomenon by way of Theorem 5.1. Note that the $p = 2$ case of the theorem is a direct analogue of a conjecture of Hoffmann in the theory of nonsingular quadratic forms which remains open, even over fields of characteristic different from 2 (see [Hof95, §4], [IV00, Conj. 1.6]).

7. FURTHER REMARKS ON THE SPLITTING OF QUASILINEAR QUADRATIC FORMS

Having determined all possible *standard* splitting patterns of quasilinear p -forms (Theorem 6.1, Proposition 6.4), we now turn our attention towards the problem of obtaining a similar result for the *full* splitting pattern. Here, we restrict our considerations to the case of quasilinear *quadratic* forms, where we can take direct inspiration from the following theorem of Vishik (which may be deduced from the existence of “excellent connections” in the integral Chow motives of anisotropic quadrics over fields of characteristic $\neq 2$ – see [Vis11, Thm. 1.3]):

Theorem 7.1 (Vishik, [Vis11]). *Let ϕ be an anisotropic quadratic form of dimension ≥ 2 over a field k of characteristic $\neq 2$, and write $\dim \phi - \mathfrak{i}_1(\phi) = 2^{r_1} - 2^{r_2} + \dots + (-1)^{s-1} 2^{r_s}$ for uniquely determined integers $r_1 > r_2 > \dots > r_{s-1} > r_s + 1 > 1$. Let $1 \leq l \leq s$, and put $D_l = \sum_{i=1}^{l-1} (-1)^{i-1} 2^{r_i-1} + \epsilon(l) \sum_{j=l}^s (-1)^{j-1} 2^{r_j}$, where $\epsilon(l) = 1$ (resp. $\epsilon(l) = 0$) if l is even (resp. odd). Then, for any field extension L of k , we either have $\mathfrak{i}_0(\phi_L) \geq D_l + \mathfrak{i}_1(\phi)$ or $\mathfrak{i}_0(\phi_L) \leq D_l$.*

Proof. This is nothing else but a restatement of [Vis11, Prop. 2.6] in terms of Witt indices. To see how it may be derived from [Vis11, Thm 2.1] in further detail, we refer the reader to [Scu14, Proof of Thm. 1.2]. \square

Examples 7.2. Let ϕ be an anisotropic quadratic form of dimension ≥ 2 over a field k of characteristic $\neq 2$, and write $\dim \phi = 2^n + m$ for uniquely determined integers $n \geq 0$ and $1 \leq m \leq 2^n$.

- (1) For $l = 1$, Theorem 7.1 asserts that $\mathfrak{i}_0(\phi_L) \geq \mathfrak{i}_1(\phi)$ whenever ϕ_L is isotropic.
- (2) For $l = 2$, Theorem 7.1 asserts that, for any field extension L of k , we either have $\mathfrak{i}_0(\phi_L) \geq m$ or $\mathfrak{i}_0(\phi_L) \leq m - \mathfrak{i}_1(\phi_L)$.
- (3) For $l = s$, Theorem 7.1 asserts that, for any field extension L of k , we either have $\mathfrak{i}_0(\phi_L) \geq \frac{1}{2}(\dim \phi + \mathfrak{i}_1(\phi) - 2^{r_s})$ or $\mathfrak{i}_0(\phi_L) \leq \frac{1}{2}(\dim \phi - \mathfrak{i}_1(\phi) - 2^{r_s})$. Taking $L = \bar{k}$ (so that $\mathfrak{i}_0(\phi_L) = \lceil \frac{1}{2} \dim \phi \rceil$), we see that $\mathfrak{i}_1(\phi) \leq 2^{r_s}$, which gives a proof of Hoffmann’s Conjecture 1.2 in this setting (see [Vis11, Thm 2.5.])

We expect, in fact, that Theorem 7.1 extends verbatim to our setting. Henceforth, let us assume that $p = 2$. We state the following:

Conjecture 7.3. *Let ϕ be an anisotropic quasilinear quadratic form of dimension ≥ 2 over F , and write $\dim \phi - \mathbf{i}_1(\phi) = 2^{r_1} - 2^{r_2} + \dots + (-1)^{s-1} 2^{r_s}$ for uniquely determined integers $r_1 > r_2 > \dots > r_{s-1} > r_s + 1 > 1$. Let $1 \leq l \leq s$, and put $D_l = \sum_{i=1}^{l-1} (-1)^{i-1} 2^{r_{i-1}} + \epsilon(l) \sum_{j=l}^s (-1)^{j-1} 2^{r_j}$, where $\epsilon(l) = 1$ (resp. $\epsilon(l) = 0$) if l is even (resp. odd). If L is any field extension of F , then we either have $\mathbf{i}_0(\phi_L) \geq D_l + \mathbf{i}_1(\phi)$ or $\mathbf{i}_0(\phi_L) \leq D_l$.*

This expectation is partly justified by:

Proposition 7.4. *Conjecture 7.3 holds for $l \leq 2$.*

Proof. As in Example 7.2 (1) (resp. (2)), the $l = 1$ (resp. $l = 2$) case is nothing else but Lemma 4.1 (resp. Theorem 3.18). \square

At present, we do not have a general approach to the $l > 2$ case of Conjecture 7.3. Using Theorem 1.3, however, we can provide further evidence for the $l = s$ case. First, it is worth stating here the following lemma:

Lemma 7.5. *In order to prove Conjecture 7.3, we may assume that $\mathbf{i}_1(\phi_L) \geq \mathbf{i}_1(\phi)$.*

Proof. Suppose that the statement of the conjecture fails to hold. In other words, suppose that $\mathbf{i}_1(\phi) > 1$ and $\mathbf{i}_0(\phi_L) = D_l + t$ for some $1 \leq t < \mathbf{i}_1(\phi)$. Note that we necessarily have $r_s > 0$ by Theorem 1.3. Now, let $\sigma = (\phi_L)_{\text{an}}$. Since $D(\sigma)$ is spanned by elements of $D(\phi)$, there exists a subform $\rho \subset \phi$ such that $\sigma \simeq \rho_L$ (see Lemma 2.2). Let ψ be any codimension- $(t-1)$ subform of ϕ containing ρ . By construction, we have $D(\psi_L) \subseteq D(\phi_L) = D(\sigma) = D(\rho_L) \subseteq D(\psi_L)$, whence $D(\psi_L) = D(\sigma)$, or, equivalently, $(\psi_L)_{\text{an}} \simeq \sigma$. In particular, we have $\mathbf{i}_0(\psi_L) = D_l + 1$. On the other hand, since $t < \mathbf{i}_1(\phi)$, ψ is a neighbour of ϕ , and so $\dim \psi - \mathbf{i}_1(\psi) = \dim \phi - \mathbf{i}_1(\phi) = 2^{r_1} - 2^{r_2} + \dots + (-1)^{s-1} 2^{r_s}$ (Proposition 3.10). We therefore conclude that the statement of the conjecture also fails for the triple (ψ, l, L) . Since we are looking to produce a contradiction, we can replace ϕ by ψ in order to arrive at the case where $t = 1$. In this case, we claim that $\mathbf{i}_1(\phi_L) \geq \mathbf{i}_1(\phi)$. To see this, note first that $L(\phi)$ is L -isomorphic to a purely transcendental extension of $L(\sigma)$ (Remark 2.6 (3)). In view of Lemma 2.26, it follows that $\mathbf{i}_1(\phi_L) = \mathbf{i}_1(\sigma_{L(\phi)})$. In particular, we have $\mathbf{i}_0(\phi_{L(\phi)}) = \mathbf{i}_0(\phi_L) + \mathbf{i}_0(\sigma_{L(\phi)}) = \mathbf{i}_0(\phi_L) + \mathbf{i}_1(\phi_L)$. The statement of Proposition 3.16 may therefore be rewritten here as

$$(7.1) \quad \mathbf{i}_1(\phi_L) + \mathbf{i}_0(\phi_L) - \mathbf{i}_1(\phi) \geq \min \left\{ \mathbf{i}_0(\phi_L), \left\lfloor \frac{\dim \phi - \mathbf{i}_1(\phi) + 1}{2} \right\rfloor \right\}.$$

But, since $r_s > 0$, we have $\mathbf{i}_0(\phi_L) = D_l + 1 \leq 2^{r_1-1} - 2^{r_2-1} + \dots + (-1)^{s-1} 2^{r_s-1} = \frac{1}{2}(\dim \phi - \mathbf{i}_1(\phi))$. Inequality (7.1) therefore yields the desired assertion, and so the lemma is proved. \square

We can now prove:

Proposition 7.6. *Conjecture 7.3 holds when $l = s$ and $L = F(Q)$ is the function field of any integral (affine or projective) quadric Q over F .*

Proof. In view of Lemma 2.26, the statement of the conjecture is stable under replacing F by any separable extension of itself. We may therefore assume that L is a purely inseparable quadratic extension of F . By Lemma 7.5, we may also assume that $\mathbf{i}_1(\phi_L) \geq$

$i_1(\phi)$. Suppose now that the statement fails to hold, so that $i_1(\phi) > 1$ and $i_0(\phi_L) = D_s + t$ for some $1 \leq t < i_1(\phi)$. We then have

$$\dim(\phi_L)_{\text{an}} = \begin{cases} 2^{r_1-1} - 2^{r_2-1} + \dots - 2^{r_{s-2}-1} + 2^{r_{s-1}-1} + i_1(\phi) - t & \text{if } s \text{ is even} \\ 2^{r_1-1} - 2^{r_2-1} + \dots - 2^{r_{s-1}-1} + 2^{r_s} + i_1(\phi) - t & \text{if } s \text{ is odd.} \end{cases}$$

Since $i_1(\phi_L) \geq i_1(\phi)$, and since $1 \leq i_1(\phi) - t < i_1(\phi)$, Theorem 1.3 implies that

$$i_1(\phi_L) \geq \begin{cases} 2^{r_{s-1}-1} + i_1(\phi) - t & \text{if } s \text{ is even} \\ 2^{r_s} + i_1(\phi) - t & \text{if } s \text{ is odd,} \end{cases}$$

from which we conclude that

$$i_0(\phi_L) + i_1(\phi_L) \geq \begin{cases} 2^{r_1-1} - 2^{r_2-1} + \dots + 2^{r_{s-1}-1} + i_1(\phi) & \text{if } s \text{ is even} \\ 2^{r_1-1} - 2^{r_2-1} + \dots + 2^{r_{s-2}-1} + i_1(\phi) & \text{if } s \text{ is odd.} \end{cases}$$

But $i_0(\phi_L) + i_1(\phi_L) = i_0(\phi_{L(\phi)}) = i_1(\phi) + i_0((\phi_1)_{L(\phi)})$ (see the proof of Lemma 7.5), and so we have

$$i_0((\phi_1)_{L(\phi)}) \geq \begin{cases} 2^{r_1-1} - 2^{r_2-1} + \dots + 2^{r_{s-1}-1} & \text{if } s \text{ is even} \\ 2^{r_1-1} - 2^{r_2-1} + \dots + 2^{r_{s-2}-1} & \text{if } s \text{ is odd.} \end{cases}$$

Either way, we see that $i_0((\phi_1)_{L(\phi)}) > 2^{r_1-1} - 2^{r_2-1} + \dots + (-1)^{s-1}2^{r_{s-1}} = \frac{1}{2}(\dim \phi - i_1(\phi)) = \frac{1}{2}\dim \phi_1$. Since $L(\phi)$ is a purely inseparable quadratic extension of $F(\phi)$, this is impossible by Lemma 2.27 (5), and so the result follows. \square

In general, it suffices to prove Conjecture 7.3 in the case where L is a finite purely inseparable extension of F . Proposition 7.6 shows that the $l = s$ case of the conjecture holds for degree 2 purely inseparable extensions. More generally, the proposition covers the case where $\text{ndeg}(\phi_L) = \frac{1}{2}\text{ndeg}(\phi)$. Indeed, using Lemma 2.26 and 2.27 (3), one can easily reduce this case to that of a quadratic extension.

Acknowledgements. This work was carried out while I was a visiting postdoctoral fellow at Max-Planck-Institut für Mathematik in Bonn. I would like to thank this institution for its support and hospitality throughout the duration of my stay.

REFERENCES

- [EKM08] R. Elman, N. Karpenko, and A. Merkurjev. *The algebraic and geometric theory of quadratic forms*. AMS Colloquium Publications, **56**, American Mathematical Society, 2008.
- [Gro61] A. Grothendieck. *Éléments de géométrie algébrique (rédigés avec la collaboration de Jean Dieudonné): II. Étude globale élémentaire de quelques classes de morphismes*. Publications Mathématiques de l’IHÉS, **8** (1961), 5–222.
- [Hau13] O. Houton. *On the first Steenrod square for Chow groups*. Amer. J. Math., **135** (2013), 53–63.
- [Hau15] O. Houton. *Detection by regular schemes in degree two*. Algebr. Geom. **2** (2015), no. 1, 44–61.
- [HL04] D.W. Hoffmann and A. Laghribi. *Quadratic forms and Pfister neighbors in characteristic 2*. Trans. Amer. Math. Soc., **356** (2004), no. 10, 4019–4053.
- [HL06] D.W. Hoffmann and A. Laghribi. *Isotropy of quadratic forms over the function field of a quadric in characteristic 2*. J. Algebra, **295** (2006), no. 2, 362–386.
- [Hof95] D.W. Hoffmann. *Isotropy of quadratic forms over the function field of a quadric*. Math. Z., **220** (1995), no.3, 461–476.

- [Hof04] D.W. Hoffmann. *Diagonal forms of degree p in characteristic p* . Algebraic and arithmetic theory of quadratic forms, 135–183, Contemp. Math., **344**, Amer. Math. Soc., 2004.
- [Izh04] O. Izhboldin. *Virtual Pfister neighbors and first Witt index. With an introduction by Nikita Karpenko*. Geometric methods in the algebraic theory of quadratic forms, 131–142, Lecture Notes in Math., **1835**, Springer, 2004.
- [IV00] Oleg Izhboldin and Alexander Vishik. *Quadratic forms with absolutely maximal splitting*. Quadratic forms and their applications (Dublin, 1999), 103–125, Contemp. Math., **272**, Amer. Math. Soc., 2000.
- [Kar03] N.A. Karpenko. *On the first Witt index of quadratic forms*. Invent. Math., **153** (2003), no.2, 455–462.
- [KM03] Nikita Karpenko and Alexander Merkurjev. *Essential dimension of quadrics*. Invent. Math., **153** (2003), no.2, 361–372.
- [Kne76] M. Knebusch. *Generic splitting of quadratic forms. I*. Proc. London Math. Soc. (3), **33** (1976), no. 1, 65–93.
- [Lag04a] A. Laghribi. *Quasi-hyperbolicity of totally singular quadratic forms*. Algebraic and arithmetic theory of quadratic forms, 237–248, Contemp. Math., **344**, Amer. Math. Soc., 2004.
- [Lag04b] A. Laghribi. *On splitting of totally singular quadratic forms*. Rend. Circ. Mat. Palermo (2), **53** (2004), no. 3, 325–336.
- [Lag06] A. Laghribi. *The norm theorem for totally singular quadratic forms*. Rocky Mountain J. Math., **36** (2006), no. 3, 575–592.
- [Lag07] A. Laghribi. *Sur le déploiement des formes bilinéaires en caractéristique 2*. Pacific J. Math., **232** (2007), no. 1, 207–232.
- [Lan02] S. Lang. *Algebra. Revised Third Edition*. Graduate Texts in Mathematics, **211**, Springer Verlag, 2002.
- [Mer03] A. Merkurjev. *Steenrod operations and degree formulas*. J. Reine Angew. Math., **565** (2003), 13–26.
- [Scu13] S. Scully. *Rational maps between quasilinear hypersurfaces*. Compos. Math., **149** (2013), no.3, 333–355.
- [Scu14] S. Scully. *On the splitting of quasilinear p -forms*. J. Reine Angew. Math., to appear, 2014, DOI: 10.1515/crelle-2013-0121.
- [Tot08] B. Totaro. *Birational geometry of quadrics in characteristic 2*. J. Algebraic Geom., **17** (2008), no. 3, 577–597.
- [Tot09] B. Totaro. *Birational geometry of quadrics*. Bull. Soc. Math. France, **137** (2009), no. 2, 253–276.
- [Vis98] A. Vishik. *Integral motives of quadrics*. MPIM Preprint, 1998-13.
- [Vis99] A. Vishik. *Direct summands in the motives of quadrics*. Preprint, 1999, <https://www.maths.nottingham.ac.uk/personal/av/papers.html>.
- [Vis04] A. Vishik. *Motives of quadrics with applications to the theory of quadratic forms*. Geometric methods in the algebraic theory of quadratic forms, 25–101, Lecture Notes in Math., **1835**, Springer, 2004.
- [Vis11] A. Vishik. *Excellent connections in the motives of quadrics*. Ann. Sci. Éc. Norm. Supér. (4), **44** (2011), no. 1, 183–195.

DEPARTMENT OF MATHEMATICAL AND STATISTICAL SCIENCES, UNIVERSITY OF ALBERTA, EDMONTON AB T6G 2G1, CANADA

E-mail address: stephenjscully@gmail.com