



A modular analysis of adaptive (non-)convex optimization: Optimism, composite objectives, variance reduction, and variational bounds

Pooria Joulani ^{a,*}, András György ^{a,2}, Csaba Szepesvári ^{a,b}

^a Deepmind, London, UK

^b University of Alberta, Edmonton, Alberta, Canada

ARTICLE INFO

Article history:

Available online 12 November 2019

Keywords:

Online learning

Stochastic optimization

Adaptive algorithms

Regret bound

Follow the regularized leader

Mirror descent

ABSTRACT

Recently, much work has been done on extending the scope of online learning and incremental stochastic optimization algorithms. In this paper we contribute to this effort in two ways: First, based on a generalization of Bregman divergences and a generic regret decomposition, we provide a self-contained, modular analysis of the two workhorses of online learning: (general) adaptive versions of Mirror Descent (MD) and the Follow-the-Regularized-Leader (FTRL) algorithms. The analysis is done with extra care so as not to introduce assumptions not needed in the proofs and allows to combine, in a straightforward way, different algorithmic ideas (e.g., adaptivity, optimism, implicit updates, variance reduction) and learning settings (e.g., strongly convex or composite objectives). This way we are able to reprove, extend and refine a large body of the literature, while keeping the proofs concise. The second contribution is a by-product of this careful analysis: We present algorithms with improved variational bounds for smooth, composite objectives, including a new family of optimistic MD algorithms with only one projection step per round. Furthermore, we provide a simple extension of adaptive regret bounds to a class of practically relevant non-convex problem settings (namely, star-convex loss functions and their extensions) with essentially no extra effort.

© 2019 Published by Elsevier B.V.

1. Introduction

Online and stochastic optimization algorithms form the underlying machinery in much of modern machine learning. Perhaps the most well-known example is Stochastic Gradient Descent (SGD) and its adaptive variants, the so-called ADAGRAD algorithms [1,2]. Other special cases include multi-armed and linear bandit algorithms, as well as algorithms for online control, tracking and prediction with expert advice [3–5].

There are numerous algorithmic variants in online and stochastic optimization, such as adaptive [2,1] and optimistic algorithms [6–10], implicit updates [11,12], composite objectives [13,2,14], or non-monotone regularization [15]. Each of these variants has been analyzed under a specific set of assumptions on the problem, e.g., smooth [16–18], convex [4,5,19,20],

* Corresponding author.

E-mail addresses: pjoulani@google.com (P. Joulani), agyorgy@google.com (A. György), szepe@google.com (Cs. Szepesvári).

¹ During part of this work, P. Joulani was a PhD student at the University of Alberta, Edmonton, Alberta, Canada.

² During part of this work, A. György was also with Imperial College London, London, UK.

or strongly convex [21,22,19,20] objectives. However, a useful property is typically missing from the analyses: modularity. It is typically not clear from the original analysis whether the algorithmic idea can be mixed with other techniques, or whether the effect of the assumptions extend beyond the specific setting considered. For example, based on the existing analyses it is very much unclear to what extent ADAGRAD techniques, or the effects of smoothness, or variational bounds in online learning, extend to new learning settings. Thus, for every new combination of algorithmic ideas, or under every new learning setting, the algorithms are typically analyzed from scratch.

A special new learning setting is non-convex optimization. While the bulk of results in online and stochastic optimization assume the convexity of the loss functions, online and stochastic optimization algorithms have been successfully applied in settings where the objectives are non-convex. In particular, the highly popular deep learning techniques [23] are based on the application of stochastic optimization algorithms to non-convex objectives. In the face of this discrepancy between the state of the art in theory and practice, an on-going thread of research attempts to generalize the analyses of stochastic optimization to non-convex settings. In particular, certain non-convex problems have been shown to actually admit efficient optimization methods, usually taking some form of a gradient method (one such problem is matrix completion, see, e.g., [24,25]).

The goal of this paper is to provide a flexible, modular analysis of online and stochastic optimization algorithms that allows to easily combine different algorithmic techniques and learning settings under as few assumptions as possible. We demonstrate the benefits of such an analysis by combining and transferring algorithmic ideas to create new algorithms, and by exploring the extent to which the convexity assumption can be relaxed in our analysis.

1.1. Contributions

First, building on previous attempts to unify the analyses of online and stochastic optimization [4,5,19,20], we provide a unified analysis of a large family of optimization algorithms in general Hilbert spaces. The analysis is crafted to exhibit modularity by decoupling the contribution of each assumption or algorithmic idea from the analysis, so as to enable us to combine different assumptions and techniques without analyzing the algorithms from scratch. In particular, we rely on a generalized Bregman divergence definition, followed by a careful decomposition of the optimization performance (optimization error or regret) into two parts: the first part captures the generic performance of the algorithm, whereas the second part connects the assumptions about the learning setting to the information given to the algorithm. Lemma 2 in Section 2.1 provides such a decomposition.³ Then, in Theorem 3, we bound the generic (first) part, using a careful analysis of the linear regret of generalized adaptive Follow-The-Regularized-Leader (FTRL) and Mirror Descent (MD) algorithms.

Second, we use this analysis framework to provide a concise summary of a large body of previous results. Section 4 provides the basic results, and Sections 5 to 7 present the relevant extensions and applications. A combination of our techniques with variance reduction techniques is explored in Section 8.

Third, building on the aforementioned modularity, we analyze new learning algorithms. In particular, in Section 7.4 we analyze a new adaptive, optimistic, composite-objective FTRL algorithm with variational bounds for smooth convex loss functions, which combines the best properties and avoids the limitations of the previous work. We also present a new class of optimistic MD algorithms with only one MD update per round (Section 7.2).

Finally, we extend the previous results to special classes of non-convex optimization problems in Section 9; as a result of relaxing the assumptions in our framework as much as possible, these results come without any extra effort just by examining to what non-convex problems our results generalize immediately. The family of non-convex functions we arrive at generalizes a small, but practically important class of functions (the set of star-convex functions) considered in previous work on non-convex optimization [26–28].

1.2. Notation and definitions

We work with a (possibly infinite-dimensional) Hilbert space $\mathcal{H}$ over the reals. That is, $\mathcal{H}$ is a real vector space equipped with an inner product $\langle \cdot, \cdot \rangle : \mathcal{H} \times \mathcal{H} \rightarrow \mathbb{R}$, such that $\mathcal{H}$ is complete with respect to (w.r.t.) the norm induced by $\langle \cdot, \cdot \rangle$. Examples include $\mathcal{H} = \mathbb{R}^d$ (for a positive integer d) where $\langle \cdot, \cdot \rangle$ is the standard dot-product, or $\mathcal{H} = \mathbb{R}^{m \times n}$, the set of $m \times n$ real matrices, where $\langle A, B \rangle = \text{tr}(A^\top B)$, or $\mathcal{H} = \ell^2(\mathcal{C})$, the set of square-integrable real-valued functions on $\mathcal{C} \subset \mathbb{R}^d$, where $\langle f, g \rangle = \int_{\mathcal{C}} f(x)g(x)dx$ for any $f, g \in \mathcal{H}$.

We denote the extended real line by $\overline{\mathbb{R}} := \mathbb{R} \cup \{-\infty, +\infty\}$, and work with functions of the form $f : \mathcal{H} \rightarrow \overline{\mathbb{R}}$. Given a set $C \subset \mathcal{H}$, the *indicatrix* of C is the function $\mathcal{I}_C : \mathcal{H} \rightarrow \overline{\mathbb{R}}$ given by $\mathcal{I}_C(x) = 0$ for $x \in C$ and $\mathcal{I}_C(x) = +\infty$ for $x \notin C$. The *effective domain* of a function $f : \mathcal{H} \rightarrow \overline{\mathbb{R}}$, denoted by $\text{dom}(f)$, is the set $\{x \in \mathcal{H} \mid f(x) < +\infty\}$ where f is less than infinity; conversely, we identify any function $f : C \rightarrow \overline{\mathbb{R}}$ defined only on a set $C \subset \mathcal{H}$ by the function $f + \mathcal{I}_C$. A function f is *proper* if $\text{dom}(f)$ is non-empty and $f(x) > -\infty$ for all $x \in \mathcal{H}$.

³ Our approach can be viewed as a refined version of the so-called “follow the leader/be the leader”-style analysis. Previous work (e.g., [20,4]) may give the impression that “follow-the-leader/be-the-leader” analyses lose constant factors while other methods such as primal-dual analysis do not. This is not the case about our analysis. In fact, we improve constants in optimistic online learning; see Section 7 for details.

Let $f : \mathcal{H} \rightarrow \overline{\mathbb{R}}$ be proper. We denote the set of all sub-gradients of f at $x \in \mathcal{H}$ by $\partial f(x)$, that is,

$$\partial f(x) := \{ u \in \mathcal{H} \mid \forall y \in \mathcal{H}, \langle u, y - x \rangle + f(x) \leq f(y) \}.$$

The function f is *sub-differentiable* at x if $\partial f(x) \neq \emptyset$; we use $f'(x)$ to denote any member of $\partial f(x)$. Note $\partial f(x) = \emptyset$ when $x \notin \text{dom}(f)$.

Let $x \in \text{dom}(f)$, assume that $f(x) > -\infty$, and let $z \in \mathcal{H}$. The *directional derivative* of f at x in the direction z is defined as $f'(x; z) := \lim_{\alpha \downarrow 0} \frac{f(x+\alpha z) - f(x)}{\alpha}$, provided that the limit exists in $[-\infty, +\infty]$. The function f is *differentiable* at x if it has a *gradient* at x , that is, a vector $\nabla f(x) \in \mathcal{H}$ such that $f'(x; z) = \langle \nabla f(x), z \rangle$ for all $z \in \mathcal{H}$. The function f is *locally sub-differentiable* at x if it has a *local sub-gradient* at x , that is, a vector $g_x \in \mathcal{H}$ such that $\langle g_x, z \rangle \leq f'(x; z)$ for all $z \in \mathcal{H}$. We denote the set of local sub-gradients of f at x by $\delta f(x)$. Note that if $f'(x; z)$ exists for all $z \in \mathcal{H}$, and f is sub-differentiable at x , then it is also locally sub-differentiable with $g_x = u$ for any $u \in \partial f(x)$. Similarly, if f is differentiable at x , then it is also locally sub-differentiable, with $g_x = \nabla f(x)$. The function f is called *directionally differentiable* at $x \in \text{dom}(f)$ if $f(x) > -\infty$ and $f'(x; z)$ exists in $[-\infty, +\infty]$ for all $z \in \mathcal{H}$; f is called *directionally differentiable* if it is directionally differentiable at every $x \in \text{dom}(f)$. Note that a directionally differentiable function is proper.

Next, we define a generalized⁴ notion of Bregman divergence:

Definition 1 (Bregman divergence). Let f be directionally differentiable at $x \in \text{dom}(f)$. The f -induced Bregman divergence from x is the function from $\mathcal{H} \rightarrow \overline{\mathbb{R}}$, given by

$$\mathcal{B}_f(y, x) := \begin{cases} f(y) - f(x) - f'(x; y - x) & \text{if } f(y) \text{ is finite;} \\ +\infty & \text{otherwise.} \end{cases} \quad (1)$$

A function $f : \mathcal{H} \rightarrow \overline{\mathbb{R}}$ is *convex* if for all $x, y \in \text{dom}(f)$ and all $\alpha \in (0, 1)$, $\alpha f(x) + (1 - \alpha)f(y) \geq f(\alpha x + (1 - \alpha)y)$. We can show that a proper convex function is always directionally differentiable, and the Bregman divergence it induces is always non-negative (see Appendix E). Let $\|\cdot\|$ denote a norm on $\mathcal{H}$ and let $L, \beta > 0$. A directionally differentiable function $f : \mathcal{H} \rightarrow \overline{\mathbb{R}}$ is β -strongly convex w.r.t. $\|\cdot\|$ if $\mathcal{B}_f(x, y) \geq \frac{\beta}{2} \|x - y\|^2$ for all $x, y \in \text{dom}(f)$. The function f is L -smooth w.r.t. $\|\cdot\|$ if for all $x, y \in \text{dom}(f)$, $|\mathcal{B}_f(x, y)| \leq \frac{L}{2} \|x - y\|^2$.

We use $\{c_t\}_{t=i}^j$ to denote the sequence $c_i, c_{i+1}, \dots, c_j$, and $c_{i:j}$ to denote the sum $\sum_{t=i}^j c_t$, with $c_{i:j} := 0$ for $i > j$.

2. Problem setting: online optimization

We study a general first-order iterative optimization setting that encompasses several common optimization scenarios, including online, stochastic, and full-gradient optimization. Consider a convex set $\mathcal{X} \subset \mathcal{H}$, a sequence of directionally differentiable functions $f_1, f_2, \dots, f_T$ from $\mathcal{H}$ to $\overline{\mathbb{R}}$ with $\mathcal{X} \subset \text{dom}(f_t)$ for all $t = 1, 2, \dots, T$, and a first-order iterative optimization algorithm. The algorithm starts with an initial point x_1 . Then, in each iteration $t = 1, 2, \dots, T$, the algorithm suffers a loss $f_t(x_t)$ from the latest point x_t , receives some feedback $g_t \in \mathcal{H}$, and selects the next point x_{t+1} . Typically, $\langle g_t, \cdot \rangle$ is supposed to be an estimate or lower bound on the directional derivative of f_t at x_t . This protocol is summarized in Fig. 1.

Unlike online convex optimization (OCO), at this stage we do not assume that the f_t are convex⁵ or differentiable, nor do we assume that g_t are gradients or sub-gradients. Our goal is to minimize the *regret* $R_T(x^*)$ against any $x^* \in \mathcal{X}$, defined as

$$R_T(x^*) = \sum_{t=1}^T (f_t(x_t) - f_t(x^*)). \quad (2)$$

More precisely, we want to guarantee that $\sup_{x^* \in \mathcal{X}} R_T(x^*)$ is small.

In this paper we are also concerned with the problem of stochastic optimization, where the goal is to minimize a function f defined by $f(x) := \mathbb{E}_{\xi \sim D} \{F(x, \xi)\}$. That is, the optimization algorithm has to find an estimate $\hat{x}_T$ such that $\mathbb{E}\{f(\hat{x}_T)\} - f(x^*)$ is small, given noisy gradient observations g_t such that $\mathbb{E}\{g_t | x_t\} \in \delta f(x_t)$. It is well-known (e.g., [4, Theorem 5.1]) that for any f , this equals $\mathbb{E}\{R_T(x^*)/T\}$ if $\hat{x}_T$ is selected uniformly at random from $x_1, \dots, x_T$. Also, if f is convex, $\mathbb{E}\{f(\hat{x}_T)\} - f(x^*) \leq \mathbb{E}\{R_T(x^*)/T\}$ if $\hat{x}_T$ is the average of $x_1, \dots, x_T$. Thus, the analysis of online algorithms that provide guarantees on the regret $R_T(x^*)$ can be used to provide performance bounds for iterative stochastic optimization, and we will analyze stochastic optimization algorithms using this approach. Note that this sometimes comes with a sacrifice.

⁴ If f is differentiable at x , then (1) matches the traditional definition of Bregman divergence. Previous work also considered generalized Bregman divergences, e.g., the works of Telgarsky and Dasgupta [29], Kiwiel [30] and the references therein. However, our definition is not limited to convex functions, allowing us to study convex and non-convex functions and regularizers under a unified theory; see, e.g., Sections 3 and 9.

⁵ There is a long tradition of non-convex assumptions in the Stochastic Approximation (SA) literature, see, e.g., the book of Bertsekas and Shreve [31]. Our results differ in that they apply to more recent advances in online learning (e.g., AdaGrad algorithms), and we derive any-time regret bounds, rather than asymptotic convergence results, for specific non-convex function classes.

Input: Convex set $\mathcal{X} \subset \mathcal{H}$; directionally differentiable functions $f_1, f_2, \dots, f_T$ from $\mathcal{H}$ to $\overline{\mathbb{R}}$.

- The algorithm selects an initial point $x_1 \in \mathcal{X}$.
- **For each** time step $t = 1, 2, \dots, T$:
 - The algorithm observes feedback $g_t \in \mathcal{H}$ and selects the next point $x_{t+1} \in \mathcal{X}$.

Goal: Minimize the regret $R_T(x^*)$ (given by (2)) against any $x^* \in \mathcal{X}$.

Fig. 1. Iterative optimization.

For example, in strongly-convex stochastic optimization, using suffix-averaging (averaging just the last αT iterates with $\alpha < 1$ instead of all T of them) [32,33] or successive tuning and restarting [34] leads to a slightly improved, optimal performance that is not available through the direct reduction mentioned above [34]. The regret framework is also not directly applicable to explain some refined notions, such as acceleration (see, e.g., [17]). Furthermore, in optimization one often cares about the performance of the last iterate x_T . In particular, in non-convex problems, the randomized reduction scheme mentioned above does not provide the high-probability guarantees that exist for the last iterate (e.g., [35]). Even in convex problems, the analysis of the last iterate follows only indirectly from a regret analysis [32,33]. Some of these problems can be addressed, for example, by considering a weighted regret where the terms in $R_T(x^*)$ have different weights (see, e.g., [36]) or by additional tricks applied on top of the regret analysis (such as done in [32–34]), which is beyond the scope of our paper. Finally, our analysis concerns global convergence, and hence applies only to convex and a small class of non-convex problems, while there are several other results in non-convex stochastic optimization that we do not recover (mostly about proving convergence rates to local optima, e.g., [35,37]).

2.1. Regret decomposition

Below, we provide a decomposition of $R_T(x^*)$, which holds for any sequence of points $x_1, x_2, \dots, x_{T+1}$ and any x^* . The decomposition is in terms of the *forward linear regret* $R_T^+(x^*)$, defined as

$$R_T^+(x^*) := \sum_{t=1}^T \langle g_t, x_{t+1} - x^* \rangle.$$

Intuitively, R_T^+ is the regret (in linear losses) of the “cheating” algorithm that uses action x_{t+1} at time t , and depends only on the choices of the algorithm and the feedback it receives.

Lemma 2 (Regret decomposition). *Let $x^*, x_1, x_2, \dots, x_{T+1}$ be any sequence of points in $\mathcal{X}$. For $t = 1, 2, \dots, T$, let $f_t : \mathcal{H} \rightarrow \overline{\mathbb{R}}$ be directionally differentiable with $\mathcal{X} \subset \text{dom}(f_t)$, and let $g_t \in \mathcal{H}$. Then,*

$$R_T(x^*) = R_T^+(x^*) + \sum_{t=1}^T \langle g_t, x_t - x_{t+1} \rangle - \sum_{t=1}^T \mathcal{B}_{f_t}(x^*, x_t) + \sum_{t=1}^T \delta_t, \quad (3)$$

where $\delta_t = -f'(x_t; x^* - x_t) + \langle g_t, x^* - x_t \rangle$.

Proof. By definition,

$$\begin{aligned} f_t(x_t) - f_t(x^*) &= -\mathcal{B}_{f_t}(x^*, x_t) - f'(x_t; x^* - x_t) \\ &= \langle g_t, x_t - x^* \rangle - \mathcal{B}_{f_t}(x^*, x_t) + \delta_t \\ &= \langle g_t, x_{t+1} - x^* \rangle + \langle g_t, x_t - x_{t+1} \rangle - \mathcal{B}_{f_t}(x^*, x_t) + \delta_t. \end{aligned}$$

Summing over t completes the proof. $\square$

Intuitively, the second term captures the regret due to the algorithm’s inability to look ahead into the future.⁶ The last two terms capture, respectively, the gain in regret that is possible due to the curvature of f_t , and the accuracy of the first-order (gradient) information g_t . In light of this lemma, controlling the regret reduces to controlling the individual terms in (3). A similar use of the forward regret appears in [40], though their analysis is limited to smooth loss functions.

Next, we provide upper bounds on $R_T^+(x^*)$ for a large class of online algorithms.

⁶ This is also related to the concept of “prediction drift”, which appears in learning with delayed feedback [38], and to the role of stability in online algorithms [39].

3. The algorithms: ADA-FTRL and ADA-MD

In this section, we analyze ADA-FTRL and ADA-MD. These two algorithms generalize the well-known core algorithms of online optimization: FTRL [4,5] and MD [41–43,14]. In particular, ADA-FTRL and ADA-MD capture variants of FTRL and MD such as Dual-Averaging [44,13], AdaGrad [2,1], composite-objective algorithms [13,2,14], implicit-update MD [11,12], strongly-convex and non-linearized FTRL [21,22,19,20], optimistic FTRL and MD [6–10], and even algorithms like AdaDelay [15] that violate the common non-decreasing regularization assumption existing in much of the previous work.⁷

3.1. ADA-FTRL: generalized adaptive follow-the-regularized-leader

The ADA-FTRL algorithm works with two sequences of regularizers, $p_1, p_2, \dots, p_T$ and $q_0, q_1, q_2, \dots, q_T$, where each p_t and q_t is a function from $\mathcal{H}$ to $\overline{\mathbb{R}}$. At time $t = 0, 1, 2, \dots, T$, having received $(g_s)_{s=1}^t$, ADA-FTRL uses $g_{1:t}, p_{1:t}$ and $q_{0:t}$ to compute the next point x_{t+1} . The regularizers p_t and q_t can be built by ADA-FTRL in an online adaptive manner using the information generated up to the end of time step t (including g_t and x_t). In particular, we use p_t to distinguish the “proximal” part of this adaptive regularization: for all $t = 1, 2, \dots, T$, we require that p_t (but not necessarily q_t) be minimized over $\mathcal{X}$ at x_t , that is,⁸

$$p_t(x_t) = \inf_{x \in \mathcal{X}} p_t(x) < +\infty. \quad (4)$$

With the definitions above, for $t = 0, 1, 2, \dots, T$, ADA-FTRL selects x_{t+1} such that

$$x_{t+1} \in \operatorname{argmin}_{x \in \mathcal{X}} \langle g_{1:t}, x \rangle + p_{1:t}(x) + q_{0:t}(x). \quad (5)$$

In particular, this means that the initial point x_1 satisfies⁹

$$x_1 \in \operatorname{argmin}_{x \in \mathcal{X}} q_0(x).$$

In addition, for notational convenience, we define $r_t := p_t + q_{t-1}, t = 1, 2, \dots, T$, so that

$$x_{t+1} \in \operatorname{argmin}_{x \in \mathcal{X}} \langle g_{1:t}, x \rangle + q_t(x) + r_{1:t}(x). \quad (6)$$

Finally, we need to make a minimal assumption to ensure that ADA-FTRL is well-defined.

Assumption 1 (Well-posed ADA-FTRL). The functions q_0 and $p_t, q_t, t = 1, 2, \dots, T$, are proper. In addition, for all $t = 0, 1, \dots, T$, the argmin sets that define x_{t+1} in (5) are non-empty, and their optimal values are finite. Finally, for all $t = 1, 2, \dots, T$, $r_{1:t}$ is directionally differentiable, and p_t satisfies (4).

Table 1 provides examples of several special cases of ADA-FTRL. In particular, ADA-FTRL combines, unifies and considerably extends the two major types of FTRL algorithms previously considered in the literature, that is, the so-called FTRL-CENTERED and FTRL-PROX algorithms [20] and their variants, as discussed in the subsequent sections.

3.2. ADA-MD: generalized adaptive mirror-descent

As in ADA-FTRL, the ADA-MD algorithm uses two sequences of regularizer functions from $\mathcal{H}$ to $\overline{\mathbb{R}}$: $r_1, r_2, \dots, r_T$ and $q_0, q_1, \dots, q_T$. Further, we assume that the domains of (r_t) are non-increasing, that is, $\operatorname{dom}(r_t) \subset \operatorname{dom}(r_{t-1})$ for $t = 2, 3, \dots, T$. Again, q_t, r_t can be created using the information generated by the end of time step t . The initial point x_1 of ADA-MD satisfies¹⁰

$$x_1 \in \operatorname{argmin}_{x \in \mathcal{X}} q_0(x).$$

Furthermore, at time $t = 1, 2, \dots, T$, having observed $(g_s)_{s=1}^t$, ADA-MD uses g_t, q_t and $r_{1:t}$ to select the point x_{t+1} such that

$$x_{t+1} \in \operatorname{argmin}_{x \in \mathcal{X}} \langle g_t, x \rangle + q_t(x) + \mathcal{B}_{r_{1:t}}(x, x_t). \quad (7)$$

⁷ AdaDelay [15] is an algorithm for delayed-feedback distributed stochastic optimization. While this setting is beyond the scope of the current paper and is left for future work, we discuss the potential implications of our framework for AdaDelay in Section 10.

⁸ Note that x_t does not depend on p_t , but is rather computed using only $p_{1:t-1}$. Once x_t is calculated, p_t can be chosen so that (4) holds (and then used in computing x_{t+1}).

⁹ The case of an arbitrary x_1 is equivalent to using, e.g., $q_0 \equiv 0$ (and changing q_1 correspondingly).

¹⁰ The case of an arbitrary x_1 is equivalent to using, e.g., $q_0 \equiv 0$ (and changing r_1 correspondingly).

Table 1

Some special instances of ADA-FTRL; for more examples, see also the survey of McMahan [20].

Algorithm	Regularization	Notes, conditions and assumptions
Online Gradient Descent (OGD)	$q_0 = \frac{1}{2\eta} \ \cdot\ _2^2$ $q_t = p_t = 0$	$\mathcal{X} = \mathbb{R}^d$, $\eta > 0$ Update: $x_{t+1} = x_t - \eta g_t$
Dual Averaging (DA)	$q_t = \frac{\alpha_t}{2} \ \cdot\ _2^2$ $p_t = 0$	$\alpha_{0:t} > 0$, $\alpha_t \geq 0$ ($t \geq 1$)
AdaGrad – Dual Averaging	$q_{0:t} = \frac{1}{2} \ x\ _{(t)}^2$ $p_t = 0$	$\ x\ _{(t)}^2 := \frac{1}{\eta} x^\top (Q_{0:t}^{1/2}) x$ $Q_0 = \gamma I$ $Q_{1:t} = \sum_{s=1}^t g_s g_s^\top$ (full-matrix update) $Q_{1:t}^{(j,j)} = \sum_{s=1}^t g_{s,j}^2$ (diagonal-matrix update)
FTRL-PROX	$q_t = 0$ $p_t = \frac{1}{2} \ x - x_t\ _{(t)}^2 - \frac{1}{2} \ x - x_t\ _{(t-1)}^2$	$Q_0 = 0$ Q_t and $\ \cdot\ _{(t)}$ as in AdaGrad-DA
Composite-Objective Online Learning	$q_0 = \tilde{q}_0$ $q_t = \psi_t + \tilde{q}_t$ $p_t = \tilde{p}_t$	For adding composite-objective learning to any instance of ADA-FTRL (see also Section 5) $x_{t+1} = \operatorname{argmin}_{\mathcal{X}} \langle g_{1:t}, x \rangle + \psi_{1:t}(x) + \tilde{p}_{1:t}(x) + \tilde{q}_{0:t}(x)$

In addition, similarly to ADA-FTRL, we define $p_t := r_t - q_{t-1}$, $t = 1, 2, \dots, T$, though we do not require p_t to be minimized at x_t in ADA-MD.¹¹

Finally, we present our assumption on the regularizers of ADA-MD. Compared to ADA-FTRL, we require a stronger assumption to ensure that ADA-MD is well-defined, and that the Bregman divergences in (7) have a controlled behavior.

Assumption 2 (Well-posed ADA-MD). The regularizers q_t, r_t , $t = 1, 2, \dots, T$, are proper, and q_0 is directionally differentiable. In addition, for all $t = 0, 1, \dots, T$, the argmin sets that define x_{t+1} in (7) are non-empty, and their optimal values are finite. Finally, for all $t = 1, 2, \dots, T$, $q_t, r_{1:t}$, and $r_{1:t} + q_t$ are directionally differentiable, $x_t \in \operatorname{dom}(r_{1:t})$, and $r'_{1:t}(x_t; \cdot)$ is linear in the directions inside $\operatorname{dom}(r_{1:t})$, that is, there is a vector in $\mathcal{H}$, denoted by $\nabla r_{1:t}(x_t)$, such that $r'_{1:t}(x_t, x - x_t) = \langle \nabla r_{1:t}(x_t), x - x_t \rangle$ for all $x \in \operatorname{dom}(r_{1:t})$.

Remark 1. Our results also hold under the weaker condition that $r'_{1:t}(x_t; \cdot - x_t)$ is *concave*¹² (rather than linear) on $\operatorname{dom}(r_{1:t})$. However, in case of a convex $r_{1:t}$, this weaker condition would again translate into having a linear $r'_{1:t}$, because a convex $r_{1:t}$ implies a convex $r'_{1:t}$ [45, Proposition 17.2]. While we do not *require* that $r_{1:t}$ be convex, all of our subsequent examples in the paper use convex $r_{1:t}$. Thus, in the interest of readability, we have made the stronger assumption of linear directional derivatives here.

Remark 2. Note that $r'_{1:t}$ needs to be linear only in the directions inside the domain of $r_{1:t}$. As such, we avoid the extra technical conditions required in previous work, e.g., that $r_{1:t}$ be a Legendre function to ensure x_t remains in the interior of $\operatorname{dom}(r_{1:t})$ and $\nabla r_{1:t}(x_t)$ is well-defined.

3.3. Analysis of ADA-FTRL and ADA-MD

Next we present bounds on the forward regret of ADA-FTRL and ADA-MD, and discuss their implications; the proof is provided in Appendix F.

Theorem 3 (Forward regret of ADA-FTRL and ADA-MD). For any $x^* \in \mathcal{X}$ and any sequence of linear losses $\langle g_t, \cdot \rangle$, $t = 1, 2, \dots, T$, the forward regret of ADA-FTRL under Assumption 1 satisfies

$$R_T^+(x^*) \leq \sum_{t=0}^T (q_t(x^*) - q_t(x_{t+1})) + \sum_{t=1}^T (p_t(x^*) - p_t(x_t)) - \sum_{t=1}^T \mathcal{B}_{r_{1:t}}(x_{t+1}, x_t), \quad (8)$$

whereas the forward regret of ADA-MD under Assumption 2 satisfies

$$R_T^+(x^*) \leq \sum_{t=0}^T (q_t(x^*) - q_t(x_{t+1})) + \sum_{t=1}^T \mathcal{B}_{p_t}(x^*, x_t) - \sum_{t=1}^T \mathcal{B}_{r_{1:t}}(x_{t+1}, x_t). \quad (9)$$

¹¹ We use the convention $(+\infty) - (+\infty) = +\infty$ in defining p_t .

¹² Without such assumptions, a Bregman divergence term in $r'_{1:t}$ appears in the regret bound of ADA-MD. Concavity ensures that this term is not positive and can be dropped, greatly simplifying the bounds.

Remark 3. Theorem 3 does not require the regularizers to be non-negative or (even non-strongly) convex.¹³ This opens up the possibility to accommodate a wide range of non-standard algorithmic solutions, such as the application of non-monotonic learning rates; see also Section 10 for a discussion.

Remark 4. In practice, ADA-FTRL and ADA-MD need to pick a specific x_{t+1} from the possibly multiple optima of (5) and (7). The bounds of Theorem 3 apply irrespective of the tie-breaking scheme.

In subsequent sections, we show that the generality of ADA-FTRL and ADA-MD, together with the flexibility of Assumptions 1 and 2, considerably facilitates the handling of various algorithmic ideas and problem settings, and allows us to combine them without requiring a new analysis for each new combination.

4. Recoveries and extensions

Lemma 2 and Theorem 3 together immediately result in generic upper bounds on the regret, given in (A.1) and (A.2) in Appendix A. Under different assumptions on the losses and regularizers, these generic bounds directly translate into concrete bounds for specific learning settings. We explore these concrete bounds in the rest of this section.

First, we provide a list of the assumptions on the losses and the regularizers for different learning settings.¹⁴ We consider two special cases of the setting of Section 2: Online optimization and stochastic optimization. In online optimization, we make the following assumption:

Assumption 3 (Online optimization setting). For $t = 1, 2, \dots, T$, f_t is locally sub-differentiable, and g_t is a local sub-gradient of f_t at x_t .

Note that f_t may be non-convex, and g_t does not need to define a global lower-bound (i.e., be a sub-gradient) of f_t ; see Section 1.2 for the formal definition of local sub-gradients.

Recall that the stochastic optimization setting is concerned with minimizing a function f , defined by $f(x) := \mathbb{E}_{\xi \sim D} \{F(x, \xi)\}$. In this case our performance metric is redefined to be the expected stochastic regret, $\mathbb{E}\{R_T(x^*)\} = \mathbb{E}\left\{\sum_{t=1}^T (f(x_t) - f(x^*))\right\}$ (see Section 2). Typically, if F is differentiable in x , then $g_t = \nabla F(x_t, \xi_t)$, where ξ_t is a random variable, e.g., sampled independently from D . In parallel to Assumption 3, we summarize our assumptions for this setting is as follows:

Assumption 4 (Stochastic optimization setting). The function f (defined above) is locally sub-differentiable, $f_t = f$ for all $t = 1, 2, \dots, T$, and g_t is, in expectation, a local sub-gradient of f at x_t : $\mathbb{E}\{g_t | x_t\} \in \partial f(x_t)$.

Again, it is enough for g_t to be an unbiased estimate of a local sub-gradient (Section 1.2).

In both settings we will rely on the non-negativity of the loss divergences at x^* :

Assumption 5 (Non-negative loss-divergence). For the competitor point of interest x^* (with respect to which we aim to bound $R_T(x^*)$), and for all $t = 1, 2, \dots, T$, $\mathcal{B}_{f_t}(x^*, x_t) \geq 0$.

It is well known that this assumption is satisfied when each f_t is convex. However, as we shall see in Section 9, this condition also holds for certain classes of non-convex functions (e.g., star-convex functions and more). In the stochastic optimization setting, since $f_t = f$, this condition boils down to $\mathcal{B}_f(x^*, x_t) \geq 0$, $t = 1, 2, \dots, T$.

In both settings, the regret can be reduced when the losses are strongly convex. Furthermore, in the stochastic optimization setting, the smoothness of the loss is also helpful in decreasing the regret. The next two assumptions capture these conditions.

Assumption 6 (Loss smoothness). The function f is differentiable and 1-smooth w.r.t. some norm $\|\cdot\|$.

Assumption 7 (Loss strong convexity). The losses are 1-strongly convex w.r.t. the regularizers, that is, $\mathcal{B}_{f_t}(x^*, x_t) \geq \mathcal{B}_{p_t}(x^*, x_t)$ for $t = 1, 2, \dots, T$.

Note that if p_t, q_{t-1} are convex, then it suffices to have $\mathcal{B}_{r_t}$ in the condition (rather than $\mathcal{B}_{p_t}$). Typically, if f_t is strongly convex w.r.t. a norm $\|\cdot\|_{(t)}$, then p_t (or r_t) is set to $\eta \|\cdot\|_{(t)}^2$ for some $\eta > 0$. Again, in stochastic optimization, Assumption 7

¹³ Nevertheless, such assumptions are useful when combining the theorem with Lemma 2.

¹⁴ In fact, compared to previous work (e.g., the references listed in Section 1 and Section 3), these are typically relaxed versions of the usual assumptions.




Fig. 2. A summary of the proof techniques that incorporate each of the assumptions into regret bounds; see Corollaries 19–21 and Table 2. The identity in the middle is from Lemma 2, whereas the top and bottom bounds on R_T^+ are due to Theorem 3. Each arrow shows the transformation of one of the terms, using the stated assumption or technique. The matching terms cancel, and the terms shown in red appear in the final bounds. (For interpretation of the colors in the figures, the reader is referred to the web version of this article.)

Table 2

Recovered and generalized standard results for online optimization (OO) and stochastic optimization (SO); see Corollaries 19–21. A number in parentheses indicates that the assumption is not directly required, but is implied by the other assumptions. In the bounds below, $\sigma_t := g_t - \nabla f(x_t)$, $D := f(x_1) - \inf_{\mathcal{X}} f(x)$, and $8'$ refers to a slightly modified version of Assumption 8, as described in Corollary 21. Note that setting $q_T = 0$ recovers the *off-by-one* property [20] in FTRL-CENTERED vs. FTRL-PROX; ADA-MD exhibits a similar property. For composite-objective ADA-FTRL and ADA-MD, these same bounds apply with $\tilde{q}_t$ in place of q_t in the summation; see Table 1 and Section 5.

Setting/algorithms	Assumptions	Regret/expected stochastic regret bound
OO/SO ADA-FTRL	1, 3/4, 5, 8	$\sum_{t=0}^T (q_t(x^*) - q_t(x_{t+1}))$ $+ \sum_{t=1}^T (p_t(x^*) - p_t(x_t)) + \sum_{t=1}^T \frac{1}{2} \ g_t\ _{(t),*}^2$
OO/SO ADA-MD	2, 3/4, 5, 8	$\sum_{t=0}^T (q_t(x^*) - q_t(x_{t+1}))$ $+ \sum_{t=1}^T \mathcal{B}_{p_t}(x^*, x_t) + \frac{1}{2} \ g_t\ _{(t),*}^2$
Strongly-convex OO/SO ADA-MD	2, 3/4, (5), 8, 7	$\sum_{t=0}^T (q_t(x^*) - q_t(x_{t+1}))$ $+ \sum_{t=1}^T \frac{1}{2} \ g_t\ _{(t),*}^2$
Smooth SO ADA-FTRL	1, 4, 6, 5, 8'	$\sum_{t=0}^T (q_t(x^*) - q_t(x_{t+1}))$ $+ \sum_{t=1}^T (p_t(x^*) - p_t(x_t)) + \sum_{t=1}^T \frac{1}{2} \ \sigma_t\ _{(t),*}^2 + D$
Smooth SO ADA-MD	2, 4, 6, 5, 8'	$\sum_{t=0}^T (q_t(x^*) - q_t(x_{t+1}))$ $+ \sum_{t=1}^T \mathcal{B}_{p_t}(x^*, x_t) + \frac{1}{2} \ \sigma_t\ _{(t),*}^2 + D$
Smooth & strongly-convex SO ADA-MD	2, 4, 6, (5), 8', 7	$\frac{1}{2} \ x^* - x_1\ ^2 + D + \sum_{t=0}^T (q_t(x^*) - q_t(x_{t+1}))$ $+ \sum_{t=1}^T \frac{1}{2} \ \sigma_t\ _{(t),*}^2$

simplifies to $\mathcal{B}_f(x^*, x_t) \geq \mathcal{B}_{p_t}(x^*, x_t)$, $t = 1, 2, \dots, T$. Furthermore, if p_t is convex, then Assumption 7 implies that f_t is convex.

Finally, the results that we recover depend on the assumption that the total regularization, in both ADA-FTRL and ADA-MD, is strongly convex:

Assumption 8 (Strong convexity of regularizers). For all $t = 1, 2, \dots, T$, $r_{1:t}$ is 1-strongly convex w.r.t. some norm $\|\cdot\|_{(t)}$.

Table 2 provides a summary of the standard results, under different sub-sets of the assumptions above, that are recovered and generalized using our framework. The derivations of these results are provided in the form of three corollaries in Appendix A. Note that the analysis is absolutely modular: each assumption is simply plugged into (A.1) or (A.2) to obtain the final bounds, without the need for a separate analysis of ADA-FTRL and ADA-MD for each individual setting. A schematic

Table 3

Concrete bounds for online optimization (OO) and stochastic optimization (SO) derived from the generic bounds of Table 2; references are given to the original bounds we recover, as well as to results showing their optimality or suboptimality. In all bounds we assume $\mathcal{H} = \mathbb{R}^d$. For SO, $\hat{x}_T$ is the average iterate $x_{1:T}$ or picked uniformly at random; see the discussion before Section 2.1. For smooth optimization, D and σ are defined as in Table 2. Derivations are straightforward, involving plugging in the parameters and/or applying specific results. In particular, the second line uses the basic inequality $\sum_{t=1}^T (1/t) \leq (1 + \log(T))$, the third line uses $\sum_{t=1}^T (1/\sqrt{t}) \leq 2\sqrt{T}$, and the fourth and fifth lines use, respectively, Lemmas 4 and 10 of Duchi et al. [2].

Algorithm	Regularizer	Setting & assumptions	Regret/expected stochastic bound
Online Gradient Descent (OGD)	ADA-FTRL $q_0 = \frac{1}{2\eta} \ \cdot\ _2^2$ $q_t = p_t = 0$	Convex OO / SO $\ x^*\ _2 \leq R$ $\mathbb{E}\{\ g_t\ _2^2 x_t\} \leq G^2$	$\frac{1}{2\eta} R^2 + \frac{\eta}{2} T G^2$ OO: $R_T \leq R G \sqrt{T}$ using $\eta = \frac{R}{G\sqrt{T}}$ SO: $\mathbb{E}\{f(\hat{x}_T) - f(x^*)\} \leq R G / \sqrt{T}$ Original bounds: Nemirovsky and Yudin [41], Warmuth and Jagota [46], Zinkevich [47] Optimality: Agarwal et al. [48]
Online Gradient Descent (OGD)	ADA-MD $q_t = 0$ $p_t = \frac{\beta}{2} \ \cdot\ _2^2$	Strongly convex OO / SO f_t is β -strongly-convex $\mathbb{E}\{\ g_t\ _2^2 x_t\} \leq G^2$	OO: $R_T \leq \frac{G^2}{\beta} \sum_{t=1}^T 1/t \leq \frac{G^2}{\beta} (1 + \log(T))$ SO: $\mathbb{E}\{f(\hat{x}_T) - f(x^*)\} \leq \frac{G^2}{\beta} (1 + \log(T))$ Original bounds: Hazan et al. [22], Hazan and Kale [34] Optimality: Agarwal et al. [48]
Dual Averaging (DA)	ADA-FTRL $q_{0:t} = \frac{L+\alpha_t}{2} \ \cdot\ _2^2$ $p_t = 0$	Convex smooth SO f convex and L -smooth $\mathbb{E}\{\ \sigma_t\ _2^2 x_t\} \leq \sigma^2$ $\ x^*\ _2 \leq R$	$R_T \leq \frac{L+\alpha_{T-1}}{2} R^2 + D + \sigma^2 \sum_{t=1}^T \frac{1}{2\alpha_{t-1}}$ Using $\alpha_{t-1} = \frac{\sigma}{R} \sqrt{2t}$: $\mathbb{E}\{f(\hat{x}_T) - f(x^*)\} \leq \frac{LR^2+2D}{2T} + \frac{\sqrt{2}\sigma R}{\sqrt{T}}$ Original bound: Dekel et al. [18] Suboptimality: Nemirovsky and Yudin [41]
AdaGrad - MD (diagonal update)	$p_{1:t} = \frac{1}{2\eta} x^\top (Q_{1:t}^{1/2}) x$ $Q_{1:t}^{(j,i)} = \sum_{s=1}^t g_{s,j}^2 g_{s,i}^2$ $q_t = 0$	Convex OO / SO $\ x - y\ _\infty \leq R, \forall x, y \in \mathcal{X}$	$R_T \leq (\frac{1}{2\eta} R^2 + \eta) \sum_{i=1}^d \sqrt{\sum_{t=1}^T g_{t,i}^2}$ Using $\eta = R/\sqrt{2}$: $\mathbb{E}\{f(\hat{x}_T) - f(x^*)\} \leq \frac{\sqrt{2}R}{T} \mathbb{E}\left\{\sum_{i=1}^d \ g_{1:T,i}\ _2\right\}$ Original bounds: Duchi et al. [2], McMahan and Streeter [1] Optimality: McMahan and Streeter [1]
AdaGrad - DA (full updates)	$q_{0:t} = \frac{1}{2\eta} x^\top (Q_{0:t}^{1/2}) x$ $Q_{1:t} = \sum_{s=1}^t g_s g_s^\top$ $Q_0 = G I$ and $p_t = 0$	Convex OO / SO $\ x^*\ _2 \leq R$ $\ g_t\ _2 \leq G$	$R_T \leq \frac{G}{\eta} R^2 + (\frac{1}{\eta} R^2 + \eta) \text{tr}(Q_{1:T}^{1/2})$ Using $\eta = R$: $\mathbb{E}\{f(\hat{x}_T) - f(x^*)\} \leq \frac{GR}{T} + \frac{2R}{T} \mathbb{E}\left\{\text{tr}(Q_{1:T}^{1/2})\right\}$ Original bounds: Duchi et al. [2], McMahan and Streeter [1] Optimality: McMahan and Streeter [1]

view of the (standard) proof ideas is given in Fig. 2. Finally, in Table 3, we provide examples of concrete bounds for common special cases of ADA-FTRL and ADA-MD recovered by the generic bounds in Table 2; see also Table 1.

5. Composite-objective learning and optimization

Next, we consider the composite-objective online learning setting. In this setting, the functions f_t , from which the (local sub-)gradients g_t are generated and fed to the algorithm, comprise only part of the loss. Instead of $R_T(x^*)$, we are interested in minimizing the regret

$$R_T^{(\ell)}(x^*) := \sum_{t=1}^T f_t(x_t) + \psi_t(x_t) - f_t(x^*) - \psi_t(x^*) = R_T(x^*) + \sum_{t=1}^T \psi_t(x_t) - \psi_t(x^*),$$

using the feedback $g_t \in \delta f_t(x_t)$, where $\psi_t : \mathcal{H} \rightarrow \overline{\mathbb{R}}$ are proper functions. The functions ψ_t are not linearized, but are passed directly to the algorithm.

Naturally, one can use the q_t regularizers to pass the ψ_t functions to ADA-FTRL and ADA-MD. Then, we can obtain the exact same bounds as in Table 2 on the composite regret $R_T^{(\ell)}(x^*)$; this recovers and extends the corresponding bounds by Xiao [13], Duchi et al. [2,14], McMahan [20]. In particular, consider the following two scenarios:

Setting 1: ψ_t is known before predicting x_t . In this case, we run ADA-FTRL or ADA-MD with $q_t = \psi_{t+1} + \tilde{q}_t$, $t = 0, 1, 2, \dots, T$ (where $\psi_{T+1} := 0$). Thus, we have the update

$$x_{t+1} \in \operatorname{argmin}_{x \in \mathcal{X}} \langle g_{1:t}, x \rangle + \psi_{1:t+1}(x) + \tilde{q}_{0:t}(x) + p_{1:t}(x), \quad (10)$$

for ADA-FTRL, and

$$x_{t+1} \in \operatorname{argmin}_{x \in \mathcal{X}} \langle g_t, x \rangle + \psi_{t+1}(x) + \tilde{q}_t(x) + \mathcal{B}_{r_{1:t}}(x, x_t), \quad (11)$$

for ADA-MD. Then, we have the following result.

Corollary 4. Suppose that the iterates $x_1, x_2, \dots, x_{T+1}$ are given by the ADA-FTRL update (10) or the ADA-MD update (11), and q_t, p_t , and r_t satisfy Assumption 1 for ADA-FTRL, or Assumption 2 for ADA-MD. Then, under the conditions of each section of Corollaries 19–21, the composite regret $R_T^{(\ell)}(x^*)$ enjoys the same bound as $R_T(x^*)$, but with $\sum_{t=0}^T \tilde{q}_t(x^*) - \tilde{q}_t(x_{t+1})$ in place of $\sum_{t=0}^T q_t(x^*) - q_t(x_{t+1})$.

Proof. By definition, $\sum_{t=0}^T \tilde{q}_t(x^*) - \tilde{q}_t(x_{t+1}) - q_t(x^*) + q_t(x_{t+1}) = \sum_{t=1}^T \psi_t(x_t) - \psi_t(x^*)$. Thus, $R_T^{(\ell)}(x^*) = R_T(x^*) + \sum_{t=0}^T \tilde{q}_t(x^*) - \tilde{q}_t(x_{t+1}) - q_t(x^*) + q_t(x_{t+1})$. Upper-bounding $R_T(x^*)$ by the aforementioned corollaries completes the proof. $\square$

Setting 2: ψ_t is revealed after predicting x_t , together with g_t . In this case, we run ADA-FTRL and ADA-MD with functions $q_0 = \tilde{q}_0$, $q_t = \psi_t + \tilde{q}_t$, $t = 1, 2, \dots, T$, so that

$$x_{t+1} \in \operatorname{argmin}_{x \in \mathcal{X}} \langle g_{1:t}, x \rangle + \psi_{1:t}(x) + \tilde{q}_{0:t}(x) + p_{1:t}(x), \quad (12)$$

for ADA-FTRL, and

$$x_{t+1} \in \operatorname{argmin}_{x \in \mathcal{X}} \langle g_t, x \rangle + \psi_t(x) + \tilde{q}_t(x) + \mathcal{B}_{r_{1:t}}(x, x_t), \quad (13)$$

for ADA-MD. Then, we have the following result, proved in Appendix B.

Corollary 5. Suppose that the iterates $x_1, x_2, \dots, x_{T+1}$ are given by the ADA-FTRL update (12) or the ADA-MD update (13), and q_t, p_t , and r_t satisfy Assumption 1 for ADA-FTRL, or Assumption 2 for ADA-MD. Also, assume that $\psi_1(x_1) = 0$ and the ψ_t are non-negative. Finally, suppose that the ψ_t form a non-increasing sequence of functions, that is, $\psi_1 \geq \psi_2 \geq \dots \geq \psi_{T+1} := 0$.¹⁵ Then, under the conditions of each section of Corollaries 19–21, the composite regret $R_T^{(\ell)}(x^*)$ enjoys the same bound as $R_T(x^*)$, but with $\sum_{t=0}^T \tilde{q}_t(x^*) - \tilde{q}_t(x_{t+1})$ in place of $\sum_{t=0}^T q_t(x^*) - q_t(x_{t+1})$.

Remark 5. In both settings, the functions ψ_t are passed as part of the regularizers q_t . Thus, if the ψ_t are strongly convex, less additional regularization is needed in ADA-FTRL to ensure the strong convexity of $r_{1:t}$ because $q_{0:t-1}$ will already have some strongly convex components. In addition, in ADA-MD, when the ψ_t are convex, the $\mathcal{B}_{p_t}$ terms in (9) will be smaller than the $\mathcal{B}_{r_t}$ terms found in previous analyses of MD. This is especially useful for implicit updates, as shown in the next section. This also demonstrates another benefit of the generalized Bregman divergence: the ψ_t , and hence the p_t , may be non-smooth in general.

6. Implicit-update ADA-MD and non-linearized ADA-FTRL

Other learning settings can be captured using the idea of passing information to the algorithm using the q_t functions. This information could include, for example, the curvature of the loss. In particular, consider the composite-objective ADA-FTRL and ADA-MD, and for $t = 1, 2, \dots, T$, let ℓ_t be a differentiable loss, $f_t = \langle \nabla \ell_t(x_t), x - x_t \rangle$, and $\psi_t = \mathcal{B}_{\ell_t}(\cdot, x_t)$.¹⁶ Then, $\ell_t = f_t + \psi_t$, $g_t = \nabla f_t(x_t) = \nabla \ell_t(x_t)$, and the composite-objective ADA-FTRL update (12) is equivalent to

$$x_{t+1} \in \operatorname{argmin}_{x \in \mathcal{X}} \ell_{1:t}(x) + \tilde{q}_{0:t}(x) + p_{1:t}(x). \quad (14)$$

Thus, non-linearized FTRL, studied by McMahan [20], is a special case of ADA-FTRL. With the same f_t, ψ_t , the composite-objective ADA-MD update (13) is equivalent to

$$x_{t+1} \in \operatorname{argmin}_{x \in \mathcal{X}} \ell_t(x) + \tilde{q}_t(x) + \mathcal{B}_{r_{1:t}}(x, x_t), \quad (15)$$

so the implicit-update MD is also a special case of ADA-MD.

Again, combining Lemma 2 and Theorem 3 results in a compact analysis of these algorithmic ideas. In particular, for both updates (14) and (15), the bounds of (A.1) and (A.2) apply on the regret in f_t . Then, moving the terms $\psi_t(x^*)$ to the left turns each bound to a bound on the regret in ℓ_t . Furthermore, the terms $-\psi_t(x_{t+1}) = -\mathcal{B}_{\ell_t}(x_{t+1}, x_t)$ that remained on the right-hand side can be merged with the $-\mathcal{B}_{r_{1:t}}(x_{t+1}, x_t)$ terms. Thus, instead of $r_{0:t}$, it is enough for $r_{1:t} + \ell_t$ to be strongly convex w.r.t. the norm $\|\cdot\|_{(\ell)}$ (see the proofs of Corollaries 19–21). This means that if ℓ_t are strongly convex, then no further regularization is required: $\ell_{1:t}$ is strongly convex, and we get back the well-known logarithmic bounds for strongly-convex FTRL (Follow-The-Leader) and implicit-update MD [11,12,21,22,19,20]. In addition, as mentioned before, convexity of ℓ_t further reduces the term $\mathcal{B}_{p_t}$ in implicit-update MD.

Finally, note that multiple pieces of information can be passed to the algorithm through q_t . In particular, none of the above interfere with further use of another composite term ϕ_t and obtaining regret bounds on $\ell_t + \phi_t$, as discussed in Section 5.

¹⁵ This relaxes the assumption in the literature, e.g., by McMahan [20], that $\psi_t = \alpha_t \psi$ for some fixed, non-negative ψ minimized at x_1 , and a non-increasing sequence $\alpha_t > 0$ (e.g., $\alpha_t = 1$); see also Setting 1.

¹⁶ For non-differentiable ℓ_t , let $f_t = \langle g_t, \cdot \rangle$ and $\psi_t = \mathcal{B}_{\ell_t}(\cdot, x_t) + \ell'(x_t; \cdot - x_t) - \langle g_t, \cdot \rangle$ to get the same effect.

7. Adaptive optimistic learning and variational bounds

The goal of optimistic online learning algorithms [6,7] is to obtain improved regret bounds when playing against “easy” (i.e., predictable) sequences of losses. This includes algorithms with regret rates that grow with the total “variation”, that is, the sum of the norms of the differences between pairs of consecutive losses f_t and f_{t+1} observed in the loss sequence: the regret will be small if the loss sequence changes slowly [8].

Recently, Mohri and Yang [9] proposed an interesting comprehensive framework for analyzing adaptive FTRL algorithms for predictable sequences. The framework has also been extended to MD by Kamalaruban [10]. However, despite their generality, the regret analyses of Mohri and Yang [9] and Kamalaruban [10] can be strengthened. Specifically, the two analyses do not recover the variation-based results of Chiang et al. [8] for smooth losses. In addition, their treatment of composite objectives introduces complications, e.g., only applies to Setting 1 of Section 5 where ψ_t is known before selecting x_t .

The flexibility of the framework introduced in this paper allows us to alleviate these and other limitations. In particular, we cast the Adaptive Optimistic FTRL (AO-FTRL) algorithm of Mohri and Yang [9] as a special case of ADA-FTRL, and obtain a much simpler form of Adaptive Optimistic MD (AO-MD) as a special case of ADA-MD. Then, we strengthen and simplify the corresponding analyses, and recover and extend the results of Chiang et al. [8]. Finally, building on the modularity of our framework, we obtain an adaptive composite-objective algorithm with variational bounds that improves upon the results of Mohri and Yang [9], Kamalaruban [10], Chiang et al. [8], Rakhlin and Sridharan [6,7].

7.1. Adaptive optimistic FTRL

Consider the online optimization setting of Section 4 (Assumption 3). Suppose that the losses $f_1, f_2, \dots, f_T$ satisfy Assumption 5 (e.g., they are convex), and the sequence of points $x_{t+1}, t = 0, 1, 2, \dots, T$ is given by

$$x_{t+1} \in \operatorname{argmin}_{x \in \mathcal{X}} \langle g_{1:t} + \tilde{g}_{t+1}, x \rangle + p_{1:t}(x) + \tilde{q}_{0:t}(x),$$

where $\tilde{g}_t, t = 1, 2, \dots, T+1$, is any sequence of vectors in $\mathcal{H}$. That is, we run ADA-FTRL, but we also incorporate $\tilde{g}_{t+1}$ as a “guess” of the future loss g_{t+1} that the algorithm will suffer. Mohri and Yang [9] refer to this algorithm as AO-FTRL.

It is easy to see that AO-FTRL is a special case of ADA-FTRL: Define $\tilde{g}_0 := 0$ ¹⁷ and for $t = 0, 1, \dots, T$, let $q_t = \tilde{q}_t + \langle \tilde{g}_{t+1} - \tilde{g}_t, \cdot \rangle$. Then, $q_{0:t} = \tilde{q}_{0:t} + \langle \tilde{g}_{t+1}, \cdot \rangle$, so we have

$$x_{t+1} \in \operatorname{argmin}_{x \in \mathcal{X}} \langle g_{1:t}, x \rangle + p_{1:t}(x) + q_{0:t}(x),$$

which is the ADA-FTRL update with this specific choice of q_t . Thus, the exact same manipulations as in Corollary 19 give the following theorem, proved in Appendix C:

Theorem 6. *If the losses satisfy Assumption 5, and the regularizers q_0 and $p_t, q_t, t = 1, 2, \dots, T$, satisfy Assumptions 1 and 8, then the regret of AO-FTRL is bounded as*

$$R_T(x^*) \leq \sum_{t=0}^{T-1} (\tilde{q}_t(x^*) - \tilde{q}_t(x_{t+1})) + \sum_{t=1}^T (p_t(x^*) - p_t(x_t)) + \sum_{t=1}^T \frac{1}{2} \|g_t - \tilde{g}_t\|_{(t),*}^2. \quad (16)$$

This bound recovers Theorems 1 and 2 of Mohri and Yang [9]. Similarly, one could prove parallels of Corollaries 20 and 21 for AO-FTRL. Then, the modularity property allows us (as we do in Section 7.4) to apply the composite-objective technique of Section 5 and recover Theorems 3–7 of Mohri and Yang [9] (and hence their corollaries). Indeed, the resulting analysis simplifies and improves on the analysis of Mohri and Yang [9] in several aspects: we do not need to separate the cases for FTRL-Prox and FTRL-General, we naturally handle the composite objective case for Settings 1 and 2 while avoiding any complications with proximal regularizers, and do not lose the constant 1/2 factor. Finally, Theorem 3 allows us to improve on the results of Chiang et al. [8], as we show in Section 7.3.

7.2. Adaptive optimistic MD

Interestingly, we can use the exact same assignment $q_t = \tilde{q}_t + \langle \tilde{g}_{t+1} - \tilde{g}_t, \cdot \rangle$ in ADA-MD. This results in the update

$$x_{t+1} \in \operatorname{argmin}_{x \in \mathcal{X}} \langle g_t + \tilde{g}_{t+1} - \tilde{g}_t, x \rangle + \tilde{q}_t(x) + \mathcal{B}_{p_t}(x, x_t).$$

Applying the same argument as in Theorem 6, one can show that this optimistic MD algorithm enjoys the regret bound of (16) with the $p_t(x^*) - p_t(x_t)$ terms replaced by $\mathcal{B}_{p_t}(x^*, x_t)$. This gives an optimistic MD algorithm with only one projection

¹⁷ This is different from the restriction in Mohri and Yang [9] that $\tilde{g}_1$ be 0; we do not require that restriction. In particular, we allow x_1 to depend on $\tilde{g}_1$, which can be arbitrary.

in each round; all other formulations [10,7,6,8] require two MD steps in each round. This new formulation has the potential to greatly simplify the previous analyses of variants of optimistic MD. In particular, handling implicit updates or composite terms is a matter of including them in $\tilde{q}_t$. Especially, unlike Kamalaruban [10], we can handle Setting 2 in the exact same way as we do in the AO-FTRL case (see Section 7.4). Further exploration of the properties of this new class of algorithms is left for future work.

7.3. Variation-based bounds for online learning

Suppose that the losses f_t are differentiable and convex, and define $f_0 := 0$. For any norm $\|\cdot\|$, we define the total variation of the losses in $\|\cdot\|_*^2$ as

$$D_{\|\cdot\|} := \sum_{t=1}^T \sup_{x \in \mathcal{X}} \|\nabla f_t(x) - \nabla f_{t-1}(x)\|_*^2. \quad (17)$$

Chiang et al. [8] use an optimistic MD algorithm to obtain regret bounds of order $O(\sqrt{D_2})$, where $D_2 = D_{\|\cdot\|_2}$, for linear as well as smooth losses.

If the losses are linear, that is, $f_t = \langle g_t, \cdot \rangle$, then Theorem 6 immediately recovers the result of Chiang et al. [8, Theorem 8]. In particular, let $\tilde{q}_0 = (1/2\eta)\|\cdot\|_2^2$, and for $t = 1, 2, \dots, T$, let $p_t = \tilde{q}_t = 0$, $\|\cdot\|_{(t)}^2 = \eta\|\cdot\|_2^2$, and $\tilde{g}_t = g_{t-1}$. Then (16) gives the regret bound $(\eta/2)\|x^*\|_2^2 + (1/(2\eta))D_2$. If $\|x\|_2 \leq 1$ and we set η based on D_2 (as Chiang et al. assume), we obtain their $O(\sqrt{D_2})$ bound.

If the losses are not linear but are L -smooth, then by the combination of Lemma 2 and Theorem 3, we still obtain $\sqrt{D_{\|\cdot\|}}$ -bounds, as Chiang et al. [8, Theorem 10] also obtain for D_2 . This is because, unlike the analysis of Mohri and Yang [9], we retain the negative terms $-B_{r_{1:t}}(x_{t+1}, x_t)$ (essentially having the same role as the B_t terms of [8]) in the regret bound. Combined with ideas from Lemma 13 of [8], this gives the desired bounds in terms of $D_{\|\cdot\|}$, proved in Appendix C:

Theorem 7. Consider the conditions of Theorem 6, and further suppose that the losses f_t are convex and L -smooth w.r.t. a norm $\|\cdot\|$. For $t = 1, 2, \dots, T+1$, let $\eta_t > 0$, and suppose that Assumption 8 holds with $\|\cdot\|_{(t)}^2 = \eta_t\|\cdot\|^2$. Further assume that $q_0 \geq 0$, $p_t, q_t \geq 0, t \geq 1$, and $\eta_t\eta_{t+1} \geq 8L^2, t = 1, 2, \dots, T$. Then, AO-FTRL with $\tilde{g}_t = g_{t-1}$ satisfies

$$R_T(x^*) \leq \tilde{q}_{0:T}(x^*) + p_{1:T}(x^*) + 2 \sum_{t=1}^T \frac{1}{\eta_t} \max_{x \in \mathcal{X}} \|\nabla f_t(x) - \nabla f_{t-1}(x)\|_*^2. \quad (18)$$

Letting $\eta_t = \eta = \sqrt{D_{\|\cdot\|}}$, and $\tilde{q}_0 = \eta\|\cdot\|^2, \tilde{q}_t, p_t = 0, t \geq 1$, generalizes the $O(\sqrt{D_2})$ bound of Chiang et al. [8] to any norm (under the same assumption they make, that $D_{\|\cdot\|} \geq 8L^2$). In the next section, we provide an algorithm that does not need prior knowledge of $D_{\|\cdot\|}$.

7.4. Adaptive optimistic composite-objective learning with variational bounds

Next, we provide a simple analysis of the composite-objective version of AO-FTRL, and obtain variational bounds in terms of $D_{\|\cdot\|}$ for composite objectives with smooth f_t . We focus on Setting 2; similar results are immediate for Setting 1. Consider the update

$$x_{t+1} \in \operatorname{argmin}_{x \in \mathcal{X}} \langle g_{1:t} + \tilde{g}_{t+1}, x \rangle + \psi_{1:t}(x) + p_{1:t}(x) + \tilde{q}_{0:t}(x), \quad (19)$$

that is, the composite-objective AO-FTRL algorithm. Then we have the following corollary of Theorem 6.

Corollary 8. Suppose that $\psi_t, t = 1, 2, \dots, T$, satisfy the conditions of Corollary 5, and $\tilde{q}_0$ and $p_t, \tilde{q}_t, t \geq 1$, are non-negative. Let $q_0 = \tilde{q}_0 + \langle \tilde{g}_1, \cdot \rangle$ and $q_t = \tilde{q}_t + \psi_t + \langle \tilde{g}_{t+1} - \tilde{g}_t, \cdot \rangle$. Suppose that $q_0, p_t, q_t, t \geq 1$ satisfy Assumptions 1 and 8, and $f_t, t \geq 1$ satisfy Assumption 5. Then, composite-objective AO-FTRL (update (19)) satisfies

$$R_T^{(\ell)}(x^*) \leq \tilde{q}_{0:T-1}(x^*) + p_{1:T}(x^*) + \sum_{t=1}^T \frac{1}{2} \|g_t - \tilde{g}_t\|_{(t),*}^2.$$

Proof. Starting as in Corollary 5, defining $\tilde{g}_0 = 0$, and noting that $0 = q_0 - \tilde{q}_0 - \langle \tilde{g}_1, \cdot \rangle$,

$$R_T^{(\ell)}(x^*) \leq R_T(x^*) + \sum_{t=0}^T q_t(x_{t+1}) - \tilde{q}_t(x_{t+1}) + \tilde{q}_t(x^*) - q_t(x^*) - \langle \tilde{g}_{t+1} - \tilde{g}_t, x_{t+1} - x^* \rangle.$$

Proceeding as in Theorem 6 completes the proof. $\square$

Table 4

Comparison of the adaptive, optimistic and variational bounds obtained in this paper vs previous work. “Non-linear variations” refers to whether the bounds are given in terms of gradient variations only (linear) or in terms of $D_{\|\cdot\|}$.

	Base algorithm	Composite objective	Non-linear variations	Needs doubling trick or knowledge of $D_{\ \cdot\ }$	Scale-free
Chiang et al. [8]	MD	No	Yes	Uses a bound on $D_{\ \cdot\ }$	Yes
Rakhlin and Sridharan [6]	FTRL	No	No	Uses doubling trick	No
Rakhlin and Sridharan [7]	MD	No	No	Not needed	No
Mohri and Yang [9]	FTRL	Yes	No	Not needed	No
Kamalaruban [10]	MD	Yes	No	Not needed	Yes
Corollary 9	FTRL	Yes	Yes	Not needed	Yes

The bounds of Mohri and Yang [9] for Setting 2 correspond to the non-proximal FTRL case. As such, one has to set the step-size sequences according to the Dual-Averaging AdaGrad recipe (cf. Table 1), which requires an additional regularization of $\tilde{q}_0 = \sqrt{\delta} \|\cdot\|_2^2$. In contrast, in FTRL-PROX, $\tilde{q}_0 = 0$. This δ value makes Dual-Averaging AdaGrad non-scale-free, while FTRL-PROX is scale-free (i.e., the x_t are independent of the scaling of f_t). Our analysis avoids this problem by the early separation of the proximal (p_t) and non-proximal regularizers (q_t) in ADA-FTRL. In particular, $p_t, \tilde{q}_t$ in Corollary 8 can be set as $\tilde{q}_t = 0$ and $p_t = \frac{\eta_t - \eta_{t-1}}{2} \|x - x_t\|^2$ with $\eta_t = \eta \sqrt{\sum_{s=1}^t \|g_s - \tilde{g}_s\|_*^2}$, $\eta > 0$ for $t = 1, 2, \dots, T$. This gives composite-objective AO-FTRL-PROX, a scale-free adaptive optimistic algorithm for Setting 2.

In addition, using Theorem 7, we can obtain a variational bound for composite-objective optimistic FTRL-PROX (proved in Appendix C), which was not available through the analysis of Mohri and Yang [9] even under Setting 1:

Corollary 9. Let $\psi_t, t = 1, 2, \dots, T$, be convex and satisfy the conditions of Corollary 5. Further assume that f_t are convex and L -smooth w.r.t. some norm $\|\cdot\|$. Suppose that $\mathcal{X}$ is closed, and let $R^2 = \sup_{x, y \in \mathcal{X}} \|x - y\|^2 < +\infty$ be the diameter of $\mathcal{X}$ measured in $\|\cdot\|$. Define $\eta = 2/R$. Suppose we run composite-objective AO-FTRL (update (19)) with the following parameters: $\tilde{q}_0 = 0$, and for $t = 1, 2, \dots, T$, $\tilde{g}_t = g_{t-1}$, $\tilde{q}_t = 0$, and $p_t = \frac{\eta_t - \eta_{t-1}}{2} \|x - x_t\|^2$, where $\eta_0 = 0$ and $\eta_t = 4RL^2 + \eta \sqrt{\sum_{s=1}^t \|g_s - \tilde{g}_s\|_*^2}$ for $t \geq 1$. Then,

$$R_T^{(\ell)}(x^*) \leq 2R^3L^2 + R + 2R\sqrt{2D_{\|\cdot\|}} = O\left(R\sqrt{D_{\|\cdot\|}}\right). \quad (20)$$

Note that the learning rate η_t is bounded from below (by $4RL^2$), which is essential in the algorithm to achieve a combination of the best properties of Mohri and Yang [9], Chiang et al. [8], and Rakhlin and Sridharan [6,7]: First, like Mohri and Yang [9], we allow the use of composite-objectives. Second, similarly to Chiang et al. [8] (but unlike [9,6,7]), our bound applies to the variation of general convex smooth functions, and is still optimal when $L = 0$ (e.g., Corollary 2 of [7]). Third, we do not need the knowledge of $D_{\|\cdot\|}$ (required by [8]) to set the step-sizes, and avoid the regret penalty of using a doubling trick (as done by Rakhlin and Sridharan [6]). Fourth, in the practically interesting case of a composite L1 penalty ($\psi_t = \alpha_t \|\cdot\|_1$), FTRL-PROX, which is the basis of our algorithm, gives sparser solutions [20] than MD, which is the basis of the algorithms of Chiang et al. [8] and Rakhlin and Sridharan [7]. Fifth, when $L = 0$, the algorithm is scale-free (unlike [9] and [6]). Finally, the results apply to the variation measured in any norm. Table 4 provides a summary of this comparison.

8. Variance reduction with composite objectives

In this section, we show that our methods are also directly applicable to stochastic variance reduction methods with composite objectives. We provide new variance-reduced algorithms for stochastic optimization, based on ADA-FTRL and ADA-MD. In contrast, to our knowledge, the algorithms in the literature have been limited to different variants of (typically unconstrained) MD. In Section 2.1, however, we separated the effect of the algorithm (ADA-FTRL vs. ADA-MD) from the effect of the gradient estimation technique (variance-reduced estimates vs. the standard stochastic gradient). Hence, our results directly provide ADA-MD and ADA-FTRL versions of SVRG and its variants, also allowing constrained optimization (with projections). This is important in practice because, as mentioned in Section 7.4, FTRL with a composite L1 penalty results in sparser models than MD. Finally we present an error bound for our methods that covers stochastic variance-reduced gradient methods such as SVRG [49] and SVRG++ [50].

Setting: We focus on the following composite-objective optimization problem:

$$\text{find } x^* = \operatorname{argmin}_{x \in \mathcal{X}} \ell(x) := f(x) + \phi(x), \quad (21)$$

where $\phi: \mathcal{X} \rightarrow \mathbb{R}$ is a convex function, $f = \mathbb{E}_{\xi \sim \mathbb{P}_\Xi} \{F(\cdot, \xi)\}$, and for all $\xi \in \Xi$, $F(\cdot, \xi): \mathcal{H} \rightarrow \mathbb{R}$ is convex, differentiable and L -smooth w.r.t. the Hilbert-space norm, which we denote by $\|\cdot\|$. In addition, we assume that we have access to two types of gradient oracles: an exact oracle which returns $f'(x)$ at any point $x \in \mathcal{X}$ and a stochastic oracle which, given a random element $\xi \sim \mathbb{P}_\Xi$ independent of x , returns $F'(x, \xi)$ (where the derivative is taken w.r.t. the first variable, x). An important special case, typically the focus of previous work, is when $f = \frac{1}{n} \sum_{i=1}^n f_i$, where each $f_i(n)$ is differentiable and the exact and stochastic oracles work, respectively, by calculating the gradient at x on all f_i 's or on a randomly selected f_i .

Algorithm 1: Generic SVRG.

Input: BASE: An instance of ADA-FTRL or ADA-MD; $m_1, m_2, \dots$: Epoch lengths

```

1  $x_1 \leftarrow$  the first iterate of BASE
2  $\tilde{x} \leftarrow x_1$ .
3 for  $s \leftarrow 1, 2, \dots$  do
4   for  $t \leftarrow m_{1:s-1} + 1 \dots m_{1:s}$  do
5     Sample  $\xi_t \sim \mathbb{P}_\xi$  independently of  $x_t, \tilde{x}$ 
6      $g_t \leftarrow F'(x_t, \xi_t) - F'(\tilde{x}, \xi_t) + f'(\tilde{x})$ 
7     Feed BASE with the gradient estimate  $g_t$ 
8      $x_{t+1} \leftarrow$  the next iterate of BASE
9   end
10  Option I:  $\tilde{x} \leftarrow \sum_{j=1}^{m_s} x_{m_{1:s-1}+j}$ 
11  Option II: Select  $\tilde{x}$  from  $x_{m_{1:s-1}+j}, j = 1, 2, \dots, m_s$ , with equal probability
12 end

```

In Algorithm 1, we provide an SVRG meta-algorithm, which receives an instance of ADA-FTRL and ADA-MD as the base algorithm, and feeds it with variance-reduced gradient estimates. That is, we run ADA-FTRL and ADA-MD, but feed them with the g_t given by Algorithm 1. Then, we can use the results of Section 4 to obtain a risk bound for Algorithm 1.

In particular, suppose that the regularizers used by BASE satisfy Assumption 8' with the norms $\|\cdot\|_{(t)} = \sqrt{c}\|\cdot\|$ for some $c > 0$ (recall that $\|\cdot\|$ is the Hilbert-space norm). Given the setting of this section, Assumption 6 also holds with the norm $\sqrt{L}\|\cdot\|$. Finally, by construction and the tower rule for expectation, $\mathbb{E}\{g_t|x_t\} = f'(x_t)$. Thus, we are in the ‘‘Stochastic Optimization’’ setting with smooth, convex objectives, and Corollary 21 applies. However, thanks to the special construction of g_t , we can further bound the $\|\sigma_t\|_*^2$ terms in the corollary, where $\sigma_t = g_t - f'(x_t)$ is the error of the gradient estimate. This is shown by the next lemma, which is standard in the variance-reduction literature (see, e.g, Lemma A.2 of [50]).

Lemma 10. Consider Algorithm 1 in the optimization setting above. Let x^* be defined as in (21), and let $\tilde{x}_s$ denote the value of $\tilde{x}$ used in computing g_t . Then,

$$\mathbb{E}\left\{\|\sigma_t\|^2\right\} \leq 4L \mathbb{E}\left\{\ell(x_t) - \ell(x^*) + \ell(\tilde{x}_s) - \ell(x^*)\right\}.$$

Using this lemma, we can prove the following performance guarantee for Algorithm 1 (the proof is relegated to Appendix D).

Theorem 11 (Composite-objective variance reduction). Let Algorithm 1 be used with epoch lengths $m_1, \dots, m_S$, and let $b > 0$ be such that $m_{s+1} \leq bm_s$ for all $s = 1, 2, \dots, S$. Let $c > 2b + 2$ be arbitrary, and suppose that BASE is either ADA-FTRL with regularizers $q_0 = (1+c)\frac{L}{2}\|\cdot\|^2$ and $q_t = \phi$, $p_t = 0$ for $t \geq 1$, or ADA-MD with regularizers $q_0 = 0$, $p_1 = \frac{(1+c)L}{2}\|\cdot\|^2$, and $q_t = \phi$, $p_{t+1} = 0$ for $t \geq 1$. Furthermore, let R be an upper bound on $\frac{1}{2}\|x_1 - x^*\|^2$ for ADA-MD and on $\frac{1}{2}\|x^*\|^2$ for ADA-FTRL. Then, after $T = m_{1:S}$ iterations, the estimate $\tilde{x}_T = \frac{1}{T} \sum_{t=2}^{T+1} x_t$ satisfies

$$\mathbb{E}\left\{\ell(\tilde{x}_T) - \ell(x^*)\right\} \leq \frac{2(1+b+m_1)(\ell(x_1) - \ell(x^*)) + c(c+1)LR}{T(c-2b-2)}.$$

Furthermore, letting $c = 2b + 2$, after S epochs, the estimate $\tilde{x}'_S = \frac{1}{m_S} \sum_{j=2}^{m_S+1} x_{m_{1:S}+j}$ satisfies

$$\mathbb{E}\left\{\ell(\tilde{x}'_S) - \ell(x^*)\right\} \leq \frac{(1+b+m_1)(\ell(x_1) - \ell(x^*)) + (2b^2 + 5b + 3)LR}{b m_S}.$$

Remark 6. This theorem recovers the convergence rates for vanilla SVRG (first part, with $m_s = m$), as well as SVRG++ (second part, with $b = 2$).

Remark 7. Instead of Lemma 10 for convex functions, one could use Lemma 5.2 of Allen-Zhu and Yuan [50] to extend our results to strongly-convex sums of non-convex smooth functions, and also extend the results of Allen-Zhu and Yuan [50] for this family of problems to FTRL(-Proximal) algorithms. The details are left for future work.

9. Application to non-convex optimization

In this section we examine how far our approach can lead in solving non-convex optimization problem, and we identify some interesting classes of non-convex loss functions for which we can obtain performance guarantees without any extra effort.




Fig. 3. Star-convex functions: (a) z defined on the unit circle and (b) its star-convex extension to two dimensions. (c) The linear approximation of a star-convex function fitted at any point is a lower bound at the minimum.

Central to this extension is the decomposition of assumptions in our analysis: we are not using the convexity of f_t in Lemma 2 or Theorem 3, but only at the very last stage of the analysis, where convexity can ensure that Assumption 5 holds. Thus, the analysis easily extends to non-convex optimization problems where Assumption 5 either holds or could be replaced by another technique at the final stage of the analysis. In the rest of this section, we explore such classes of non-convex problems, which are also related to the Polyak–Łojasiewicz (PL) condition used in the non-convex optimization and learning community. For background and a summary of related work, consult Karimi et al. [51].

9.1. Stochastic optimization of star-convex functions

First, we explore the class of non-convex functions for which Assumption 5 directly holds. As it turns out, this is a much larger class of functions than convex functions. In particular, consider the so-called “star-convex” functions [26]¹⁸:

Definition 12 (Star-convex function). A function f is *star-convex* at a point x^* if and only if x^* is a global minimizer of f , and for all $\alpha \in [0, 1]$ and all $x \in \text{dom}(f)$:

$$f(\alpha x^* + (1 - \alpha)x) \leq \alpha f(x^*) + (1 - \alpha)f(x). \quad (22)$$

A function is said to be star-convex when it is star-convex at some of its global minimizers.

The name “star-convex” comes from the fact that the sub-level sets $L_\beta = \{x : f(x) \leq \beta\}$ of a function f that is star-convex at x^* are star-shaped with center x^* (recall that a set U is star-convex with center x if for any $y \in U$, the segment between x and y is included in U). However, note that there are functions whose sub-level sets are star-convex that are themselves not star-convex. In particular, functions that are increasing along the rays (IAR) started from their global minima have star-shaped sub-level sets and vice versa, but some of these functions (e.g., $f(x) = \sqrt{|x|}$, $x \in \mathbb{R}$) is clearly not star-convex. Recall that quasi-convex functions are those whose sub-level sets are convex. In one dimension a star-convex function is thus also necessarily quasi-convex. However, clearly, there are star-convex functions (such as $x \mapsto |x|\mathbb{I}\{|x| \leq 1\} + 2|x|\mathbb{I}\{|x| > 1\}$, $x \in \mathbb{R}$) that are not convex and in multiple dimensions there are star-convex functions that are not quasi-convex (e.g., $x \mapsto \|x\|^2 g(\frac{x}{\|x\|_2})$ where $g(u)$ is, say, the sine of the angle of u with the unit vector e_1).

Star-convex functions arise in various optimization settings, often related to sums of squares [26,27]. It is easy to see from the definitions that the set of star-convex functions is closed under non-singular affine domain transformations, addition (of functions having the same center) and multiplication by non-negative constants. Further, for $x \in \mathbb{R}^d$, $x \mapsto \prod_i |x_i|^{p_i}$ is star-convex (at zero) whenever $\sum_i p_i \geq 1$. It is also easy to see that for any positive function z defined on the unit sphere, $\|x\|z(x/\|x\|)$ is star-convex at 0 (see Fig. 3a, b). For further properties and examples see Lee and Valiant [27].

We can immediately see that Assumption 5 holds for star-convex functions:

Lemma 13 (Non-negative Bregman divergence for star-convex functions). Let f be a directionally differentiable function with global optimum x^* . Then, f is star-convex at x^* if and only if for all $x \in \mathcal{H}$,

$$B_f(x^*, x) \geq 0.$$

¹⁸ We modify the definition so that it is relative to a given fixed global minimizer as this way we capture a larger class of functions and this is all we need.

The lemma essentially states that the linear approximation of a star-convex function f at any point x is a lower bound of the function at x^* . This is illustrated in Fig. 3c.

Proof. Both directions are routine. For illustration we provide the proof of the forward direction. Assume without loss of generality that $x^* = 0$ and $f(x^*) = 0$. Then star-convexity at x^* is equivalent to having $f(\alpha x) \leq \alpha f(x)$ for any x and $\alpha \in [0, 1]$. Further, $B_f(x^*, x) \geq 0$ is equivalent to $-f(x) - f'(x; -x) \geq 0$. Now, $f'(x; -x) = \lim_{\alpha \downarrow 0} \frac{f(x+\alpha(-x)) - f(x)}{\alpha}$. Under star-convexity, $f(x + \alpha(-x)) = f((1 - \alpha)x) \leq (1 - \alpha)f(x)$. Hence, $f'(x; -x) \leq \lim_{\alpha \downarrow 0} \frac{(1 - \alpha)f(x) - f(x)}{\alpha} = -f(x)$. $\square$

Thus, Corollaries 20 and 21 apply to star-convex functions. In other words:

- For stochastic optimization of directionally-differentiable star-convex functions in Hilbert spaces, ADA-FTRL and ADA-MD converge to the global optimum *with the same rate as they converge for convex functions* (including fast rates due to other assumptions, e.g., smoothness).

Of course, a similar result holds for the online setting, too, but in this case the assumption that each f_t is star-convex w.r.t. the same center x^* becomes restrictive.

Remark 8. Since the rate of regret depends on the norm of the gradients g_t , to get fast rates one needs to control these norms. This is trivial if f is Lipschitz-continuous. However, some star-convex functions are not Lipschitz, even arbitrarily close to the optima (e.g., $f(x, y) = (\sqrt{|x|} + \sqrt{|y|})^2$). For such functions, Lee and Valiant [27] propose alternative methods to gradient descent. However, it seems possible to control the norms in these settings using additional regularization (as in the normalized gradient descent method); see, e.g., the work of Hazan et al. [52], and the recent work of Levy [53]. Exploring this idea is left for future work.

9.2. Beyond star-convex functions

Inspecting our proofs we may notice that Assumption 5 is unnecessarily restrictive: to maintain the same rate of growth for regret, it suffices for the sum of Bregman divergences to grow with the same rate as the rest of the bound, rather than being negative and hence dropped. This extends all of our results to another interesting class of non-convex functions which generalize star-convexity:

Definition 14 (τ -star-convexity, Hardt et al. [28]). Let f be a directionally differentiable function f with global optimum x^* . Then f is τ -star-convex¹⁹ on a set $\mathcal{X}$ at $x^* \in \mathcal{H}$ if there is $\tau > 0$ such that for all $x \in \mathcal{X} \cap \text{dom}(f)$,

$$\tau(f(x) - f(x^*)) \leq -f'(x; x^* - x). \quad (23)$$

Note that by Lemma 13, star-convexity corresponds to the case when $\tau = 1$. Hardt et al. [28] demonstrated that an objective function that arises naturally in the identification of certain class of linear systems is τ -star-convex with some $\tau > 0$. For differentiable functions, (23) is equivalent to $f(x) - f(x^*) \leq \frac{1}{\tau} \langle \nabla f(x), x - x^* \rangle$, so it is a simple generalization of the linear upper bound one typically uses to reduce online convex optimization to online linear optimization. Therefore, any regret bound that is proved via upper bounding linearized losses automatically extends to τ -star-convex functions. However, in general, it may require substantial work to identify what assumptions are used exactly in proving an upper bound on the linearized loss (e.g., Hardt et al. [28] reproved the convergence guarantees for smooth SGD). The next lemma shows that our techniques can automatically separate the effects of different assumptions and provide fast regret rates under appropriate circumstances.

Lemma 15 (Basic regret bound under τ -star-convexity). Let f be locally directionally differentiable and τ -star-convex on a set $\mathcal{X}$ at x^* , $f_1 \cdots = f_T = f$. Then, for all $x_t \in \mathcal{X} \cap \text{dom}(f_t)$ and $g_t \in \mathcal{H}$ ($t = 1, 2, \dots, T$),

$$R_T(x^*) \leq \frac{1}{\tau} \left(R_T^+(x^*) + \sum_{t=1}^T \langle g_t, x_t - x_{t+1} \rangle + \delta_t \right).$$

Proof. The proof can be derived from the right-hand side of (3), but a shorter direct proof is also available: Add and subtract $\langle g_t, x^* - x_t \rangle$ to the right-hand side of (23). Noticing that $-f'(x_t; x^* - x_t) + \langle g_t, x^* - x_t \rangle = \delta_t$, summing up and using the definition $R_T^+(x^*) = \sum_t \langle g_t, x_{t+1} - x^* \rangle$ gives the result. $\square$

¹⁹ Hardt et al. [28] define the same concept under τ -weakly-quasi-convexity. However, per our previous discussion, it appears more appropriate to call this property τ -star-convexity. Especially since when $\tau = 1$ we get back star-convexity, which, as we have seen is not a weakening of quasi-convexity.

Now since the regret was bounded through the expression in the parentheses of the previous display, Corollaries 20 and 21 apply. In other words:

- For stochastic optimization of directionally-differentiable τ -star-convex functions in Hilbert spaces, ADA-FTRL and ADA-MD enjoy $1/\tau$ -times the same regret as when they are applied to linearized loss functions, including fast rates due to other assumptions, e.g., smoothness.

In the convex case the strong convexity of the losses (Assumption 7) implied that their Bregman divergences are non-negative (Assumption 5). The natural generalization of this leads to the following definition:

Definition 16 (τ -star-strong-convexity). Let f, r be directionally differentiable and let x^* be a global minimum of f . Then, f is τ -star-strongly-convex w.r.t. r if $S := \text{dom}(f) \cap \text{dom}(r)$ is non-empty and there exists $\tau > 0$ such that for all $x \in S$ and some minimizer x^* of f ,

$$\tau(f(x) - f(x^*)) \leq -f'(x; x^* - x) - B_r(x^*, x). \quad (24)$$

Replacing τ -star-convexity with τ -star-strong-convexity gives the following analogue of Lemma 15:

Lemma 17 (Basic regret bound under τ -star-strong-convexity). Let f, r be locally directionally differentiable. Assume that f is τ -star-strongly-convex w.r.t. r at x^* on a set $\mathcal{X}$. Then, for all $x_t \in \mathcal{X} \cap \text{dom}(f_t) \cap \text{dom}(r)$ and $g_t \in \mathcal{H}$ ($t = 1, 2, \dots, T$),

$$R_T(x^*) \leq \frac{1}{\tau} \left(R_T^+(x^*) + \sum_{t=1}^T \langle g_t, x_t - x_{t+1} \rangle + \delta_t - B_r(x^*, x_t) \right).$$

Proof. The proof follows the same step as that of Lemma 15, except that we need to use (24) instead of (23). $\square$

It follows that the same manipulations as in Corollaries 20 and 21 imply:

- For stochastic optimization of directionally-differentiable τ -star-strongly-convex functions in Hilbert spaces, ADA-FTRL and ADA-MD converge to the global optimum with $1/\tau$ -times the same rate as they converge for strongly convex functions.

It appears that τ -star-strong-convexity is related to the Polyak-Łojasiewicz (PL) inequality. Recall that a differentiable function f satisfies the PL inequality with constant $\mu > 0$ if

$$\mu(f(x) - f(x^*)) \leq \frac{1}{2} \|\nabla f(x)\|_2^2,$$

where x^* is a global minimizer of f . Proposed independently and simultaneously by Polyak [54] and Łojasiewicz [55], the PL inequality appears to play a fundamental role in the study of incremental gradient algorithms (see [51] and the references therein). As star-convexity, the PL inequality can also be satisfied by non-convex functions, partly explaining the prominent role it plays in the analysis of gradient methods. We can see that τ -star-strong-convexity implies the PL inequality when r is the squared Euclidean norm:

Lemma 18 (PL is implied by star-strong-convexity). Let $r(x) = \frac{1}{2} \|x\|_2^2$ and let f be differentiable. If f is τ -star-strongly-convex w.r.t. r , then f also satisfies the PL inequality with $\mu = \tau$.

Proof. Assume that f satisfies (24). We have

$$-f'(x; x - x^*) = \langle \nabla f(x), x^* - x \rangle \leq \frac{1}{2} (\|\nabla f(x)\|^2 + \|x^* - x\|^2),$$

where the second step follows from the Fenchel-Young inequality. As it is well known, $B_r(x, y) = \frac{1}{2} \|x - y\|^2$. Thus, (24) implies that $\tau(f(x) - f(x^*)) \leq \frac{1}{2} \|\nabla f(x)\|_2^2$. $\square$

Finally, note that the results can be combined with other algorithmic ideas, such as implicit-update and non-linearized learning; the same extensions of Corollaries 20 and 21, as discussed in Section 6, apply here as well. We can also do the same manipulations to obtain results similar to Section 5 for composite-objective non-convex learning. Note, however, that in this case, the star-convexity assumptions must hold with x^* being the minimizer of ℓ , not f , which makes them more restrictive than the non-composite case. We leave it for future work to handle composite-objective learning of star-convex functions in full generality.

We end this section by noting that there are interesting classes of non-convex problems other than the PL class; see, e.g., Karimi et al. [51]. A direction for future work is to explore whether these classes relate to specific conditions on Bregman divergences, and whether similar convergence results for general adaptive optimization are also possible under these function classes.

10. Discussion

In this section, we compare the results obtained in this paper to the previous attempts at unified analysis of adaptive FTRL and MD. A starting point of our work was the unifying treatment of online learning algorithms by McMahan [20], as well as the generalized adaptive FTRL analysis of Orabona et al. [19].

10.1. Comparison to the analysis of McMahan [20]

McMahan [20] also studied a unified, modular analysis of MD and FTRL algorithms (albeit with different modules), assuming that the regularizers p_t, q_t, r_t are convex, non-negative, and satisfy Assumption 8. ADA-FTRL and ADA-MD encompass all of the algorithms they considered. In particular, their Theorems 1 and 2 are special cases of Corollary 19 (recall that non-linearized FTRL, and in particular strongly-convex FTRL, are also special cases of ADA-FTRL; see Section 6). In addition, our analysis applies more generally to infinite-dimensional Hilbert spaces, our presentation of ADA-FTRL encompasses a larger set of algorithms, the relaxed assumptions under which we analyzed ADA-FTRL and ADA-MD remove certain practical limitations that existed in the work of McMahan [20], and our analysis captures a wider range of learning settings. We discuss these improvements below.

Importantly, McMahan [20] also provides a reduction from MD to a version of FTRL-Prox. This, in particular, illuminates important differences between MD and FTRL in composite-objective learning. We refer the reader to Section 6 of their paper. We decided to keep the presentation of the two algorithms separate to facilitate the relaxation of the assumptions on the regularizers; see Assumptions 1 and 2 and the discussion below.

10.1.1. Relaxing the assumptions on the regularizers

A central part of the modularity of our analysis comes from the flexibility of Assumptions 1 and 2 on the regularizers of ADA-FTRL and ADA-MD. In particular, unlike McMahan [20], we do not assume that the individual regularizers p_t, q_t, r_t are non-negative or convex.

This relaxation provides two benefits. First, with the non-negativity restriction removed, we can add arbitrary, possibly linear, components to the regularizers. As we showed above, this resulted in a simple recovery and analysis of optimistic FTRL and a new class of optimistic MD algorithms (Section 7), as well as a straightforward recovery of implicit and non-linearized updates, even for non-convex functions (Section 6).

Second, with the convexity assumption removed, ADA-FTRL and ADA-MD can accommodate algorithmic ideas such as non-decreasing regularization. For example, AdaDelay [15], an instance of ADA-MD for distributed delayed stochastic optimization, uses $r_{1:t} = \eta_t \|\cdot\|^2$, but η_t is not guaranteed to be non-decreasing, that is, r_t could be negative and non-convex (while $r_{0:t}$ still remains convex for all t). Now, note that MD and FTRL-Prox are closely related. Particularly, if the p_t are themselves Bregman divergences (as in proximal ADAGRAD), then FTRL-Prox and MD have identical regret bounds. Therefore, the techniques of Sra et al. [15] for controlling the regularizer terms in the bound could be naturally applied, almost with no modification, to an FTRL-Prox version of AdaDelay. This extension to FTRL-Prox is interesting since, as mentioned before, composite FTRL-Prox with an L1 penalty tends to produce sparser results compared to ADA-MD ([56], [20], Section 6.2). Thus, while this variant of FTRL-Prox is a special case of ADA-FTRL (e.g., Corollary 19 applies), it was not clear how to analyze this algorithm under the assumptions made by McMahan [20]. We leave the detailed study of this extension for future work.

Finally, the choice to separate the proximal and non-proximal regularizers in ADA-FTRL provides certain conveniences. In particular, the q_t terms can take the role of incorporating information (such as composite terms) into ADA-FTRL, while the proximal part p_t remains intact. This precludes the need to provide a separate analysis for FTRL-Prox every time the structure of information changes (e.g., when implicit updates are added). Thus, unlike Section 5 of McMahan [20], we did not need to provide a separate analysis (their Theorem 10) for composite-objective FTRL-Prox. We also avoided the complications with composite optimistic FTRL-Prox as in Mohri and Yang [9]; see Section 7.

10.1.2. The regret decomposition and analysis of new learning settings

In comparison to McMahan [20], the analysis we provided exhibits much flexibility across learning settings. In particular, the regret decomposition given by Lemma 2 enabled us to accommodate a wide range of learning settings, and separate the effect of the learning setting from the forward regret of the algorithm. Building on this, for example, we provided a clean analysis of variational and variance-dependent bounds for smooth losses (and generalized them to adaptive algorithms). In addition, by encapsulating the effect of loss convexity into Assumption 5, we could generalize the analysis to certain non-convex classes.

10.2. Comparison to the analysis of Orabona et al. [19]

Orabona et al. [19] study a special case of ADA-FTRL where Assumption 8 holds. The main result of Orabona et al. [19], that is, their Lemma 1, can be thought of as playing the same role as (A.1). We emphasize, however, that their Lemma 1 is a quite general result. For example, with a few algebraic operations we could recover a special case of Theorem 3 (including also the case of $p_t \neq 0$) from their Lemma 1, by setting $z_t = 0$ and moving the linear components to their f_t functions. Nevertheless, our analysis extends the work of Orabona et al. [19] to infinite-dimensional Hilbert-spaces and, more importantly, to ADA-MD. Furthermore, we demonstrated a principled way of mixing algorithmic ideas and incorporating information from the learning setting into FTRL and MD using the q_t functions, which in turn led to the discovery of the single-projection optimistic MD family of algorithms. Finally, the comments of Section 10.1.2 apply.

Importantly, the authors also provide a compact analysis of the Vovk-Azoury-Warmuth algorithm, as well as online binary classification algorithms. These results are essentially obtained from combining their Lemma 1 with interesting regret decompositions other than the one we presented in Lemma 2. It seems possible to combine their regret decompositions with our analysis to extend their result to ADA-MD algorithms, and to obtain refined bounds for smooth losses. We leave this direction for future work.

11. Conclusion and future work

We provided a generalized, unified and modular framework for analyzing online and stochastic optimization algorithms, and demonstrated its flexibility on several existing, as well as new, algorithms and learning settings. Our framework can be used together with other algorithmic ideas and learning settings, for example, adaptive delayed-feedback algorithms like AdaDelay [15], but exploring these problems are out of the scope of this work. There are many interesting questions related to non-convex optimization; while we showed that our results extend to the so-called τ -star(-strongly)-convex functions, which have already found some applications, it remains to be seen whether they also extend to other settings, such as optimization of quasi-convex functions, or functions that satisfy the Polyak-Łojasiewicz inequality. Exploring these and other applications of this framework is left for future work.

Declaration of competing interest

The authors declare that they have no known competing financial interests or personal relationships that could have appeared to influence the work reported in this paper.

Acknowledgements

We would like to thank the anonymous reviewers for several insightful comments that helped improve the quality of the paper. This work was supported by the Alberta Machine Intelligence Institute (Amii) (formerly AICML) and the Natural Sciences and Engineering Research Council of Canada (NSERC). P. Joulani was also supported by a Queen Elizabeth II Doctoral Scholarship at the University of Alberta.

Appendix A. Formal statements and proofs for the standard results described in Table 2

Putting Lemma 2 and Theorem 3 together, for ADA-FTRL we obtain

$$\begin{aligned} R_T(x^*) \leq & - \sum_{t=1}^T \mathcal{B}_{f_t}(x^*, x_t) + \sum_{t=0}^T (q_t(x^*) - q_t(x_{t+1})) + \sum_{t=1}^T (p_t(x^*) - p_t(x_t)) \\ & - \sum_{t=1}^T \mathcal{B}_{r_{1:t}}(x_{t+1}, x_t) + \sum_{t=1}^T \langle g_t, x_t - x_{t+1} \rangle + \sum_{t=1}^T \delta_t, \end{aligned} \quad (\text{A.1})$$

whereas for ADA-MD,

$$\begin{aligned} R_T(x^*) \leq & - \sum_{t=1}^T \mathcal{B}_{f_t}(x^*, x_t) + \sum_{t=0}^T (q_t(x^*) - q_t(x_{t+1})) + \sum_{t=1}^T \mathcal{B}_{p_t}(x^*, x_t) \\ & - \sum_{t=1}^T \mathcal{B}_{r_{1:t}}(x_{t+1}, x_t) + \sum_{t=1}^T \langle g_t, x_t - x_{t+1} \rangle + \sum_{t=1}^T \delta_t. \end{aligned} \quad (\text{A.2})$$

Next we prove the concrete regret bounds, given in Table 2, based on the above. A schematic view of the proof ideas is given in Fig. 2.

Corollary 19. Consider the “Online Optimization” setting (Assumption 3), using ADA-FTRL (under Assumption 1) or ADA-MD (under Assumption 2). Suppose that Assumptions 5 and 8 hold. Then,

(i) the regret of ADA-MD is bounded as

$$R_T(x^*) \leq \sum_{t=0}^T (q_t(x^*) - q_t(x_{t+1})) + \sum_{t=1}^T \mathcal{B}_{p_t}(x^*, x_t) + \sum_{t=1}^T \frac{1}{2} \|g_t\|_{(t),*}^2;$$

(ii) the regret of ADA-FTRL is bounded as

$$R_T(x^*) \leq \sum_{t=0}^T (q_t(x^*) - q_t(x_{t+1})) + \sum_{t=1}^T (p_t(x^*) - p_t(x_t)) + \sum_{t=1}^T \frac{1}{2} \|g_t\|_{(t),*}^2;$$

(iii) under Assumption 7, the regret of ADA-MD is bounded as

$$R_T(x^*) \leq \sum_{t=0}^T (q_t(x^*) - q_t(x_{t+1})) + \sum_{t=1}^T \frac{1}{2} \|g_t\|_{(t),*}^2.$$

Proof. Note that by Assumption 3, we have

$$\delta_t \leq 0, \quad (\text{A.3})$$

for all $t = 1, 2, \dots, T$. In addition, by the Fenchel-Young inequality and Assumption 8,

$$\begin{aligned} \langle g_t, x_t - x_{t+1} \rangle &\leq \frac{1}{2} \|x_t - x_{t+1}\|_{(t)}^2 + \frac{1}{2} \|g_t\|_{(t),*}^2 \\ &\leq \mathcal{B}_{r_{1,t}}(x_{t+1}, x_t) + \frac{1}{2} \|g_t\|_{(t),*}^2. \end{aligned} \quad (\text{A.4})$$

Putting (A.3), (A.4), and Assumption 5 into (A.1) and (A.2) and canceling out the matching terms proves (19) and (19). Finally, to prove (21), we use Assumption 7 to cancel the $\mathcal{B}_{f_t}(x^*, x_t)$ terms with the $\mathcal{B}_{p_t}(x^*, x_t)$ terms (rather than dropping them by Assumption 5). $\square$

Corollary 20. Consider the “Stochastic Optimization” setting (Assumption 4), using ADA-FTRL (under Assumption 1) or ADA-MD (under Assumption 2). Suppose that Assumptions 5 and 8 hold. Then,

(i) the regret of ADA-MD is bounded as

$$\mathbb{E}\{R_T(x^*)\} \leq \mathbb{E}\left\{\sum_{t=0}^T (q_t(x^*) - q_t(x_{t+1})) + \sum_{t=1}^T \mathcal{B}_{p_t}(x^*, x_t) + \sum_{t=1}^T \frac{1}{2} \|g_t\|_{(t),*}^2\right\};$$

(ii) the regret of ADA-FTRL is bounded as

$$\mathbb{E}\{R_T(x^*)\} \leq \mathbb{E}\left\{\sum_{t=0}^T (q_t(x^*) - q_t(x_{t+1})) + \sum_{t=1}^T (p_t(x^*) - p_t(x_t)) + \sum_{t=1}^T \frac{1}{2} \|g_t\|_{(t),*}^2\right\};$$

(iii) under Assumption 7, the regret of ADA-MD is bounded as

$$\mathbb{E}\{R_T(x^*)\} \leq \mathbb{E}\left\{\sum_{t=0}^T (q_t(x^*) - q_t(x_{t+1})) + \sum_{t=1}^T \frac{1}{2} \|g_t\|_{(t),*}^2\right\}.$$

Proof. Let $f_t = f$ in Lemma 2 (hence in (A.1) and (A.2)), and note that by Assumption 4, we have

$$\mathbb{E}\{\delta_t\} = \mathbb{E}\{f'(x_t; x^* - x_t) - \langle \mathbb{E}\{g_t|x_t\}, x_t - x^* \rangle\} \leq 0, \quad (\text{A.5})$$

for all $t = 1, 2, \dots, T$. Similarly to the proof of Corollary 19, putting (A.5), (A.4), and Assumption 5 into (A.1) and (A.2) proves (20) and (20). Finally, to prove (21), one can use Assumption 7 to cancel the $\mathcal{B}_f(x^*, x_t)$ terms with the $\mathcal{B}_{p_t}(x^*, x_t)$ terms (rather than dropping them by Assumption 5). $\square$

Corollary 21. Consider the “Stochastic Optimization” setting (Assumption 4), using ADA-FTRL (under Assumption 1) or ADA-MD (under Assumption 2). Suppose that Assumptions 5, 6 hold, and Assumption 8 holds with $r_{1:t} - \|\cdot\|^2/2$ in place of $r_{1:t}$.²⁰ Let $f^* := \inf_{x \in \mathcal{X}} f(x)$, and define $D := f(x_1) - f^*$ and $\sigma_t := g_t - \nabla f(x_t)$. Then,

(i) the regret of ADA-MD is bounded as

$$\mathbb{E}\{R_T(x^*)\} \leq \mathbb{E}\left\{\sum_{t=0}^T (q_t(x^*) - q_t(x_{t+1})) + \sum_{t=1}^T \mathcal{B}_{p_t}(x^*, x_t) + \sum_{t=1}^T \frac{1}{2} \|\sigma_t\|_{(t),*}^2 + D\right\};$$

(ii) the regret of ADA-FTRL is bounded as

$$\mathbb{E}\{R_T(x^*)\} \leq \mathbb{E}\left\{\sum_{t=0}^T (q_t(x^*) - q_t(x_{t+1})) + \sum_{t=1}^T (p_t(x^*) - p_t(x_t)) + \sum_{t=1}^T \frac{1}{2} \|\sigma_t\|_{(t),*}^2 + D\right\};$$

(iii) under Assumption 7, the regret of ADA-MD is bounded as

$$\mathbb{E}\{R_T(x^*)\} \leq \mathbb{E}\left\{\frac{1}{2} \|x^* - x_1\|^2 + \sum_{t=0}^T (q_t(x^*) - q_t(x_{t+1})) + \sum_{t=1}^T \frac{1}{2} \|\sigma_t\|_{(t),*}^2 + D\right\}.$$

Proof. Note that for all $t = 1, 2, \dots, T$, by Assumption 6 and the Fenchel-Young inequality,

$$\begin{aligned} \langle g_t, x_t - x_{t+1} \rangle &= f(x_t) - f(x_{t+1}) + \mathcal{B}_f(x_{t+1}, x_t) + \langle g_t - \nabla f(x_t), x_t - x_{t+1} \rangle \\ &\leq f(x_t) - f(x_{t+1}) + \frac{1}{2} \|x_t - x_{t+1}\|^2 + \langle \sigma_t, x_t - x_{t+1} \rangle \\ &\leq f(x_t) - f(x_{t+1}) + \frac{1}{2} \|x_t - x_{t+1}\|^2 + \frac{1}{2} \|\sigma_t\|_{(t),*}^2 + \frac{1}{2} \|x_t - x_{t+1}\|_{(t)}^2 \end{aligned} \quad (\text{A.6})$$

Putting (A.5), (A.6), and Assumption 5 into (A.1) and (A.2), telescoping the f terms, using $f(x_{T+1}) \geq f^*$, and canceling out the matching terms gives (21) and (21). Finally, to prove (21), one can use Assumption 7 to cancel the $\mathcal{B}_f(x^*, x_t)$ terms with the $\mathcal{B}_{p_t}(x^*, x_t)$ terms (rather than dropping them by Assumption 5). $\square$

Appendix B. Proofs for Section 5

Proof of Corollary 5. Define $\psi_0 := \psi_1$. Then, using our assumptions on ψ_t , we have

$$\begin{aligned} R_T^{(\ell)}(x^*) &= R_T(x^*) + \sum_{t=1}^T (\psi_t(x_t) - \psi_t(x^*)) \\ &= R_T(x^*) + \sum_{t=1}^T (\psi_t(x_{t+1}) - \psi_t(x^*)) + \sum_{t=1}^T (\psi_t(x_t) - \psi_{t-1}(x_t)) + \psi_1(x_1) - \psi_T(x_{T+1}) \\ &\leq R_T(x^*) + \sum_{t=1}^T (\psi_t(x_{t+1}) - \psi_t(x^*)) \\ &= R_T(x^*) + \sum_{t=1}^T (q_t(x_{t+1}) - \tilde{q}_t(x_{t+1}) + \tilde{q}_t(x^*) - q_t(x^*)). \end{aligned}$$

The rest of the proof is as in Corollary 4, noting that $\tilde{q}_0 = q_0$. $\square$

Appendix C. Proofs for Section 7

Proof of Theorem 6. Starting from inequality (A.1), by the exact same manipulations as in Corollary 19:

$$R_T(x^*) \leq \sum_{t=0}^T (q_t(x^*) - q_t(x_{t+1})) + \sum_{t=1}^T \langle g_t, x_t - x_{t+1} \rangle$$

²⁰ The modification to Assumption 8 is equivalent to adding an extra $\|x\|^2/2$ regularizer to ADA-FTRL and ADA-MD.

$$\begin{aligned}
& + \sum_{t=1}^T (p_t(x^*) - p_t(x_t)) + \sum_{t=1}^T -\mathcal{B}_{r_{1:t}}(x_{t+1}, x_t) \\
& = \sum_{t=0}^T \langle \tilde{g}_{t+1} - \tilde{g}_t, x^* - x_{t+1} \rangle + \sum_{t=1}^T \langle g_t, x_t - x_{t+1} \rangle \\
& \quad + \sum_{t=0}^T (\tilde{q}_t(x^*) - \tilde{q}_t(x_{t+1})) + \sum_{t=1}^T (p_t(x^*) - p_t(x_t)) + \sum_{t=1}^T -\mathcal{B}_{r_{1:t}}(x_{t+1}, x_t) \\
& = \langle \tilde{g}_{T+1}, x^* \rangle + \sum_{t=0}^T \langle \tilde{g}_{t+1}, -x_{t+1} \rangle + \sum_{t=1}^T \langle \tilde{g}_t, x_{t+1} \rangle + \sum_{t=1}^T \langle g_t, x_t - x_{t+1} \rangle \\
& \quad + \sum_{t=0}^T (\tilde{q}_t(x^*) - \tilde{q}_t(x_{t+1})) + \sum_{t=1}^T (p_t(x^*) - p_t(x_t)) + \sum_{t=1}^T -\mathcal{B}_{r_{1:t}}(x_{t+1}, x_t) \\
& = \langle \tilde{g}_{T+1}, x^* - x_{T+1} \rangle + \sum_{t=1}^T \langle g_t - \tilde{g}_t, x_t - x_{t+1} \rangle \\
& \quad + \sum_{t=0}^T (\tilde{q}_t(x^*) - \tilde{q}_t(x_{t+1})) + \sum_{t=1}^T (p_t(x^*) - p_t(x_t)) + \sum_{t=1}^T -\mathcal{B}_{r_{1:t}}(x_{t+1}, x_t) \tag{C.1} \\
& \leq \langle \tilde{g}_{T+1}, x^* - x_{T+1} \rangle + \tilde{q}_T(x^*) - \tilde{q}_T(x_{T+1}) \\
& \quad + \sum_{t=0}^{T-1} (\tilde{q}_t(x^*) - \tilde{q}_t(x_{t+1})) + \sum_{t=1}^T (p_t(x^*) - p_t(x_t)) + \sum_{t=1}^T \frac{1}{2} \|g_t - \tilde{g}_t\|_{(t),*}^2,
\end{aligned}$$

using the Fenchel-Young for the second term, and Assumption 8 for the last term, in the final step. Finally, note that the left-hand side is independent of $\tilde{q}_T$ and $\tilde{g}_{T+1}$, and without loss of generality, we can set them to zero, which makes the first two terms of the right-hand side zero, hence finishing the proof. $\square$

Proof of Theorem 7. Define $G_t = \|g_t - \tilde{g}_t\|_*^2$, and let $\lambda_t := \eta_t/2$. Starting from (C.1), and using the fact that setting $\tilde{g}_{T+1} = 0$ does not affect the value of $R_T(x^*)$, we get

$$\begin{aligned}
R_T(x^*) & \leq \sum_{t=1}^T \langle g_t - \tilde{g}_t, x_t - x_{t+1} \rangle + \sum_{t=1}^T -\mathcal{B}_{r_{1:t}}(x_{t+1}, x_t) \\
& \quad + \sum_{t=0}^T (\tilde{q}_t(x^*) - \tilde{q}_t(x_{t+1})) + \sum_{t=1}^T (p_t(x^*) - p_t(x_t)) \\
& \leq \sum_{t=0}^T (\tilde{q}_t(x^*) - \tilde{q}_t(x_{t+1})) + \sum_{t=1}^T (p_t(x^*) - p_t(x_t)) \\
& \quad + \sum_{t=1}^T -\frac{\eta_t}{2} \|x_t - x_{t+1}\|^2 + \sum_{t=1}^T \frac{\lambda_t}{2} \|x_t - x_{t+1}\|^2 + \sum_{t=1}^T \frac{1}{2\lambda_t} \|g_t - \tilde{g}_t\|_*^2, \\
& \leq \sum_{t=0}^T (\tilde{q}_t(x^*) - \tilde{q}_t(x_{t+1})) + \sum_{t=1}^T (p_t(x^*) - p_t(x_t)) + \sum_{t=1}^T \frac{-\eta_t}{4} \|x_t - x_{t+1}\|^2 + \sum_{t=1}^T \frac{1}{\eta_t} G_t \\
& \leq \sum_{t=0}^T (\tilde{q}_t(x^*) - \tilde{q}_t(x_{t+1})) + \sum_{t=1}^T (p_t(x^*) - p_t(x_t)) \\
& \quad + \sum_{t=1}^T \frac{-2L^2}{\eta_{t+1}} \|x_t - x_{t+1}\|^2 + \sum_{t=1}^T \frac{1}{\eta_t} G_t.
\end{aligned}$$

In the second inequality, we used Assumption 8 and the Fenchel-Young inequality. In the last inequality, we used the assumption $\eta_t \eta_{t+1} \geq 8L^2$. Now, let $x_0 := x_1$ and $f_0 := 0$, so that $\tilde{g}_1 = g_0 = \nabla f_0(x_0)$. Then, using ideas from Lemma 12 of Chiang et al. [8],

$$\begin{aligned}
\sum_{t=1}^T \frac{1}{\eta_t} G_t &= \sum_{t=1}^T \frac{1}{\eta_t} \|\nabla f_t(x_t) - \nabla f_{t-1}(x_{t-1})\|_*^2 \\
&\leq \sum_{t=1}^T 2 \frac{1}{\eta_t} \|\nabla f_t(x_t) - \nabla f_{t-1}(x_t)\|_*^2 + \sum_{t=1}^T 2 \frac{1}{\eta_t} \|\nabla f_{t-1}(x_t) - \nabla f_{t-1}(x_{t-1})\|_*^2 \\
&\leq 2 \sum_{t=1}^T \frac{1}{\eta_t} \|\nabla f_t(x_t) - \nabla f_{t-1}(x_t)\|_*^2 + \sum_{t=2}^T \frac{2L^2}{\eta_t} \|x_t - x_{t-1}\|^2 \\
&\leq 2 \sum_{t=1}^T \frac{1}{\eta_t} \|\nabla f_t(x_t) - \nabla f_{t-1}(x_t)\|_*^2 + \sum_{t=1}^T \frac{2L^2}{\eta_{t+1}} \|x_{t+1} - x_t\|^2.
\end{aligned}$$

Note that to get the first inequality, we used the fact that $\|\cdot\|^2$ is convex for any norm, together with Jensen's inequality, so that $\|x + y\|^2 = 4\|x/2 + y/2\|^2 \leq 4(\|x\|^2/2 + \|y\|^2/2) = 2\|x\|^2 + 2\|y\|^2$. This completes the proof. $\square$

Proof of Corollary 9. First, note that since ψ_t is convex, by definition, $r_{0:t} = \sum_{t=1}^t \frac{\eta_t - \eta_{t-1}}{2} \|\cdot - x_t\|^2$ is η_t -strongly-convex w.r.t. the norm $\|\cdot\|$, satisfying Assumption 8. Furthermore, Assumption 5 is satisfied by the convexity of f_t . Also, by assumption, $\mathcal{X}$ is closed and $R < +\infty$, so the objectives are always bounded below and Assumption 1 holds.

Let $G_t = \|\tilde{g}_t - \tilde{g}_t^*\|_*^2$, and define $C := \eta_t - \eta\sqrt{G_{1:t}} = 4RL^2$. Starting as in Corollary 8, and following the same steps as in the proof of Theorem 7, we have

$$\begin{aligned}
R_T^{(\ell)}(x^*) &\leq \sum_{t=0}^T (\tilde{q}_t(x^*) - \tilde{q}_t(x_{t+1})) + \sum_{t=1}^T (p_t(x^*) - p_t(x_t)) - \sum_{t=1}^T \frac{\eta_t}{4} \|x_t - x_{t+1}\|^2 + \sum_{t=1}^T \frac{1}{\eta_t} G_t \\
&\leq \sum_{t=1}^T \frac{1}{2} (\eta_t - \eta_{t-1}) R^2 + \sum_{t=1}^T \frac{-\eta_t}{4} \|x_t - x_{t+1}\|^2 + \sum_{t=1}^T \frac{1}{\eta\sqrt{G_{1:t}}} G_t \\
&\leq \frac{1}{2} \eta_T R^2 + \sum_{t=1}^T \frac{-C}{4} \|x_t - x_{t+1}\|^2 + \frac{2}{\eta} \sqrt{G_{1:T}} \\
&\leq \frac{1}{2} C R^2 + \left(\frac{2}{\eta} + \frac{\eta}{2} R^2 \right) \sqrt{G_{1:T}} + \sum_{t=1}^T \frac{-C}{4} \|x_t - x_{t+1}\|^2
\end{aligned}$$

In the first line, we used, as in Theorem 7, the Fenchel-Young inequality with $\lambda_t = \eta_t/2$. In the second line, we dropped the $\tilde{q}_t = 0$ terms and used the definition of p_t and R , and the fact that $\eta_t \geq \eta_{t-1}$, to get the first term, and obtained the last term using the fact that $\eta_t \geq \eta\sqrt{G_{1:t}}$ by definition. In the third inequality, we let the η_t in the first term telescope, used the fact that $\eta_t > C$ in the second term, and Lemma 22 to get the last term. In the last line, we used the definition of η_T and grouped the $\sqrt{G_{1:T}}$ terms together.

Next, we use the inequalities $\sqrt{a+b} \leq \sqrt{a} + \sqrt{b}$ and $\sqrt{a} \leq \frac{1}{2} + a$ (for $a, b \geq 0$), as well as Jensen's inequality on $\|\cdot\|^2$ (as in the proof of Theorem 7) to bound $\sqrt{G_{1:T}}$ with $\sqrt{D_{\|\cdot\|}}$:

$$\begin{aligned}
\sqrt{\sum_{t=1}^T G_t} &= \sqrt{\sum_{t=1}^T \|\nabla f_t(x_t) - \nabla f_{t-1}(x_{t-1})\|_*^2} \\
&\leq \sqrt{\sum_{t=1}^T 2 \|\nabla f_t(x_t) - \nabla f_{t-1}(x_t)\|_*^2 + \sum_{t=1}^T 2 \|\nabla f_{t-1}(x_t) - \nabla f_{t-1}(x_{t-1})\|_*^2} \\
&\leq \sqrt{2 \sum_{t=1}^T \|\nabla f_t(x_t) - \nabla f_{t-1}(x_t)\|_*^2 + \sum_{t=1}^T 2L^2 \|x_t - x_{t-1}\|^2} \\
&\leq \sqrt{2 \sum_{t=1}^T \|\nabla f_t(x_t) - \nabla f_{t-1}(x_t)\|_*^2} + \sqrt{\sum_{t=1}^T 2L^2 \|x_t - x_{t-1}\|^2} \\
&\leq \sqrt{2 \sum_{t=1}^T \|\nabla f_t(x_t) - \nabla f_{t-1}(x_t)\|_*^2} + \frac{1}{2} + \sum_{t=1}^T 2L^2 \|x_t - x_{t-1}\|^2.
\end{aligned}$$

$$= \sqrt{2 \sum_{t=1}^T \|\nabla f_t(x_t) - \nabla f_{t-1}(x_t)\|_*^2} + \frac{1}{2} + \sum_{t=1}^T 2L^2 \|x_{t+1} - x_t\|^2,$$

where the last line follows, as in the proof of Theorem 7, by defining $x_0 = x_1$ and adding the extra positive term $2L^2 \|x_{T+1} - x_T\|^2$. Putting back into the previous inequality,

$$\begin{aligned} R_T^{(\ell)}(x^*) &\leq \frac{1}{2}CR^2 + \sum_{t=1}^T \frac{-C}{4} \|x_t - x_{t+1}\|^2 \\ &\quad + \left(\frac{2}{\eta} + \frac{\eta}{2}R^2\right) \left(\sqrt{2D_{\|\cdot\|}} + \frac{1}{2} + \sum_{t=1}^T 2L^2 \|x_{t+1} - x_t\|^2\right) \\ &= \frac{1}{2}CR^2 + \sum_{t=1}^T \frac{-C}{4} \|x_t - x_{t+1}\|^2 \\ &\quad + 2R\sqrt{2D_{\|\cdot\|}} + R + \sum_{t=1}^T 4RL^2 \|x_{t+1} - x_t\|^2 \\ &= \frac{1}{2}4R^3L^2 + R + 2R\sqrt{2D_{\|\cdot\|}}. \end{aligned}$$

In the first equality, we used $\eta = 2/R$ while in the last one we used that $C = 4RL^2$ by definition. This completes the proof. $\square$

Lemma 22 (Lemma 4 of [20]). For any non-negative numbers $a_1, a_2, \dots, a_T$ with $a_1 > 0$,

$$\sum_{t=1}^T \frac{a_t}{\sqrt{\sum_{s=1}^t a_s}} \leq 2 \sqrt{\sum_{t=1}^T a_t}.$$

Appendix D. Proofs for Section 8

Proof of Theorem 11. As promised, we will use Corollary 21 to prove the theorem. First notice that defining the norm in the corollary as $\sqrt{L}\|\cdot\|$ and the norms $\|\cdot\|_{(t)} = \sqrt{cL}\|\cdot\|$ (with dual norm $\|\cdot\|_{(t),*} = \frac{1}{\sqrt{cL}}\|\cdot\|$), the conditions of the corollary are satisfied. Therefore, by Corollary 21, after moving the composite ϕ terms and D to the left-hand side, we get

$$\mathbb{E} \left\{ \sum_{t=1}^T \ell(x_{t+1}) - \ell(x^*) \right\} \leq \mathbb{E} \left\{ \frac{(1+c)L}{2} \|x^* - x_1\|^2 + \sum_{t=1}^T \frac{1}{2cL} \|\sigma_t\|_{(t),*}^2 \right\},$$

for ADA-MD, and

$$\mathbb{E} \left\{ \sum_{t=1}^T \ell(x_{t+1}) - \ell(x^*) \right\} \leq \mathbb{E} \left\{ \frac{(1+c)L}{2} \|x^*\|^2 + \sum_{t=1}^T \frac{1}{2cL} \|\sigma_t\|_{(t),*}^2 \right\},$$

for ADA-FTRL. Let $R_T^\ell := \sum_{t=1}^T \ell(x_{t+1}) - \ell(x^*)$. Then, by Lemma 10 and the definition of the dual norm $\|\cdot\|_{(t),*}$,

$$\begin{aligned} \mathbb{E} \{R_T^\ell\} &\leq (1+c)LR + \sum_{t=1}^T \mathbb{E} \left\{ \frac{4L \mathbb{E} \{ \ell(x_t) - \ell(x^*) + \ell(\tilde{x}_s) - \ell(x^*) \}}{2cL} \right\} \\ &= (1+c)LR + \frac{2}{c} \mathbb{E} \{R_T^\ell\} + \sum_{t=1}^T \mathbb{E} \left\{ \frac{4L \mathbb{E} \{ \ell(x_t) - \ell(x_{t+1}) + \ell(\tilde{x}_s) - \ell(x^*) \}}{2cL} \right\} \\ &= (1+c)LR + \frac{2}{c} \mathbb{E} \{R_T^\ell\} + \frac{2}{c} \mathbb{E} \{ \ell(x_1) - \ell(x_{T+1}) \} + \frac{2}{c} \sum_{t=1}^T \mathbb{E} \{ \ell(\tilde{x}_s) - \ell(x^*) \}. \end{aligned}$$

Next, we bound the last summation above. Let T_s denote the set of indices t such that g_t is calculated in the s -th epoch, and let $\tilde{x}_s$ be the first iterate used (in g_t) in epoch s . We have:

$$\begin{aligned}
\sum_{t=1}^T \mathbb{E} \{ \ell(\tilde{x}_s) - \ell(x^*) \} &= \sum_{s=1}^S \sum_{t \in T_s} \mathbb{E} \{ \ell(\tilde{x}_s) - \ell(x^*) \} \\
&\leq m_1 (\ell(x_1) - \ell(x^*)) + \sum_{s=2}^S \mathbb{E} \left\{ m_s \left(\frac{1}{m_{s-1}} \sum_{t \in T_{s-1}} (\ell(x_t) - \ell(x^*)) \right) \right\} \\
&= m_1 (\ell(x_1) - \ell(x^*)) + \sum_{s=1}^{S-1} \mathbb{E} \left\{ \frac{m_{s+1}}{m_s} \sum_{t \in T_s} (\ell(x_t) - \ell(x^*)) \right\} \\
&\leq m_1 (\ell(x_1) - \ell(x^*)) + \sum_{s=1}^{S-1} \mathbb{E} \left\{ b \sum_{t \in T_s} (\ell(x_t) - \ell(x^*)) \right\} \tag{D.1} \\
&\leq m_1 (\ell(x_1) - \ell(x^*)) + b \sum_{s=1}^S \sum_{t \in T_s} \mathbb{E} \{ \ell(x_t) - \ell(x^*) \} \\
&= m_1 (\ell(x_1) - \ell(x^*)) + b \sum_{t=1}^T \mathbb{E} \{ \ell(x_t) - \ell(x^*) \} \\
&= m_1 (\ell(x_1) - \ell(x^*)) + b \mathbb{E} \{ \ell(x_1) - \ell(x_{T+1}) \} + b \mathbb{E} \{ R_T^\ell \} \\
&\leq (b + m_1) (\ell(x_1) - \ell(x^*)) + b \mathbb{E} \{ R_T^\ell \}.
\end{aligned}$$

Putting together,

$$\mathbb{E} \{ R_T^\ell \} \leq (1 + c)LR + \frac{2 + 2b}{c} \mathbb{E} \{ R_T^\ell \} + \frac{2 + 2b + 2m_1}{c} (\ell(x_1) - \ell(x^*)).$$

Rearranging and applying Jensen's inequality proves the first part of the theorem. To prove the second part, we start from (D.1):

$$\begin{aligned}
&m_1 (\ell(x_1) - \ell(x^*)) + \sum_{s=1}^{S-1} \mathbb{E} \left\{ b \sum_{t \in T_s} \ell(x_t) - \ell(x^*) \right\} \\
&\leq (m_1 + b) (\ell(x_1) - \ell(x^*)) + b \sum_{s=1}^{S-1} \sum_{t \in T_s} \mathbb{E} \{ \ell(x_{t+1}) - \ell(x^*) \}.
\end{aligned}$$

Now, if $c = 2 + 2b$, we have $1 - 2/c = 2b/c$. Thus,

$$\begin{aligned}
&\left(1 - \frac{2}{c}\right) \sum_{s=1}^S \sum_{t \in T_s} \mathbb{E} \{ \ell(x_{t+1}) - \ell(x^*) \} \\
&= \left(1 - \frac{2}{c}\right) \mathbb{E} \{ R_T^\ell \} \\
&\leq (1 + c)LR + \frac{2 + 2b + 2m_1}{c} (\ell(x_1) - \ell(x^*)) + \frac{2b}{c} \sum_{s=1}^{S-1} \sum_{t \in T_s} \mathbb{E} \{ \ell(x_{t+1}) - \ell(x^*) \}.
\end{aligned}$$

Rearranging gives

$$\left(1 - \frac{2}{c}\right) \sum_{t \in T_S} \mathbb{E} \{ \ell(x_{t+1}) - \ell(x^*) \} \leq (1 + c)LR + \frac{2 + 2b + 2m_1}{c} (\ell(x_1) - \ell(x^*)).$$

Dividing both sides by $(1 - 2/c)m_S$ and using Jensen's inequality on ℓ completes the proof. $\square$

Appendix E. Technical results

In this appendix, we have gathered some technical results required in our proofs. The first lemma states that the Bregman divergence is invariant under addition of affine functions.

Lemma 23. Let $f : \mathcal{H} \rightarrow \overline{\mathbb{R}}$ be proper, and let $x, y \in \text{dom}(f)$. Suppose that $v \in \mathcal{H}$, and $w \in \mathbb{R}$, and let $g : \mathcal{H} \rightarrow \overline{\mathbb{R}}$ be given by $g(\cdot) = f(\cdot) + \langle v, \cdot \rangle + w$. Then,

- (i) g is proper, with $\text{dom}(g) = \text{dom}(f)$.
- (ii) For any $z \in \mathcal{H}$, the derivative $g'(x; z)$ exists in $[-\infty, +\infty]$ if and only if $f'(x; z)$ exists in $[-\infty, +\infty]$, in which case

$$g'(x; z) = f'(x; z) + \langle v, z \rangle.$$

- (iii) If $f'(x; y - x)$ or $g'(x; y - x)$ exist, then $\mathcal{B}_g(y, x) = \mathcal{B}_f(y, x)$.

Proof. That g is proper and $\text{dom}(f) = \text{dom}(g)$ is immediate since $\text{dom}(\langle v, \cdot \rangle) = \mathcal{H}$ and $w \in \mathbb{R}$. Then $x, y \in \text{dom}(g)$, and for any $z \in \mathcal{H}$, if either of $f'(x; z)$ or $g'(x; z)$ exist in $[-\infty, +\infty]$,

$$f'(x; z) + \langle v, z \rangle = \lim_{\alpha \downarrow 0} \frac{f(x + \alpha z) + \langle v, x + \alpha z \rangle + w - f(x) - \langle v, x \rangle - w}{\alpha} = g'(x; v),$$

which proves the second part of the lemma. Letting $z = y - x$ and using the definition of $\mathcal{B}_g$ gives $\mathcal{B}_f(y, x) = \mathcal{B}_g(y, x)$. $\square$

The next proposition gathers useful results based on Proposition 17.2 of Bauschke and Combettes [45].

Proposition 24. Let f be proper and convex, and let $x, y \in \text{dom}(f)$ and $z \in \mathcal{H}$. Then,

- (i) $f'(x; z)$ exists in $[-\infty, +\infty]$ and

$$f'(x; z) = \inf_{\alpha \in (0, +\infty)} \frac{f(x + \alpha z) - f(x)}{\alpha}.$$

- (ii) $f'(x; y - x) < +\infty$.
- (iii) $\mathcal{B}_f(y, x) \geq 0$.

Proof. Part (24) is proved in Proposition 17.2(ii) of Bauschke and Combettes [45]. Also, by their Proposition 17.2(iii),

$$f'(x; y - x) + f(x) \leq f(y),$$

proving part (24) since $f(y)$ and $f(x)$ are both real numbers. Part (24) then simply follows from the same equation, with the Bregman divergence being real and non-negative when $f'(x; y - x)$ is real-valued, and $+\infty$ when $f'(x; y - x) = -\infty$. $\square$

The next lemma is useful for decomposing Bregman divergences.

Lemma 25. Let $r : \mathcal{H} \rightarrow \overline{\mathbb{R}}$ and $q : \mathcal{H} \rightarrow \overline{\mathbb{R}}$ be directionally differentiable (hence also proper). Let $S := \text{dom}(r) \cap \text{dom}(q)$, suppose $S \neq \emptyset$, and let $x, y \in S$. Suppose that at least one of the two limits $q'(x; y - x)$ and $r'(x; y - x)$ is finite. Then,

$$\mathcal{B}_r(y, x) - \mathcal{B}_q(y, x) = \mathcal{B}_p(y, x),$$

where $p : \mathcal{H} \rightarrow \overline{\mathbb{R}}$ is given by

$$p(z) := \begin{cases} r(z) - q(z) & z \in \text{dom}(r) \cup \text{dom}(q), \\ +\infty & \text{otherwise.} \end{cases}$$

Proof. By the assumption that at least one of them is finite, we can take the difference of the two limits $q'(x; y - x)$ and $r'(x; y - x)$ to obtain

$$\begin{aligned} -r'(x; y - x) + q'(x; y - x) &= \lim_{\alpha \downarrow 0} \frac{-r(x + \alpha(y - x)) + r(x)}{\alpha} + \lim_{\alpha \downarrow 0} \frac{q(x + \alpha(y - x)) - q(x)}{\alpha} \\ &= \lim_{\alpha \downarrow 0} \frac{-p(x + \alpha(y - x)) + p(x)}{\alpha} = -p'(x; y - x). \end{aligned} \quad (\text{E.1})$$

In the derivation above, we have used that at most one of $r(x + \alpha(y - x))$ and $q(x + \alpha(y - x))$ can remain infinite as $\alpha \downarrow 0$. Formally, there exists an $\epsilon > 0$ such that for all $\alpha < \epsilon$, the summation $-r(x + \alpha(y - x)) + q(x + \alpha(y - x))$ is well defined, and is equal, by definition, to $-p(x + \alpha(y - x))$. Adding the real-valued equation $r(y) - r(x) - q(y) + q(x) = p(y) - p(x)$ to (E.1) completes the proof. $\square$

Note that, in light of Proposition 24, the above lemma holds if both q and r are proper convex functions, given the condition that at least one of the directional derivatives is finite (i.e., not equal to $-\infty$). The latter condition is needed even if p is also a proper convex function (note that this requires that $\text{dom}(r) \subset \text{dom}(q)$). In this case, Proposition 24 (ii) shows that the sum of the limits $p'(x; y - x)$ and $q'(x; y - x)$ is well-defined for any $x, y \in \text{dom}(p) \cap \text{dom}(q)$, and is equal to $r'(x; y - x)$. Therefore, since the function values are finite, we get $\mathcal{B}_p(y, x) + \mathcal{B}_q(y, x) = \mathcal{B}_r(y, x)$. On the other hand, $\mathcal{B}_r(y, x) - \mathcal{B}_q(y, x) = \mathcal{B}_p(y, x)$ only holds if the left-hand side is well-defined, which happens exactly if at least one of the Bregman divergences (necessarily $\mathcal{B}_q(y, x)$) is finite (equivalently, at least one of the directional derivatives is finite). An example illustrating this situation is when all functions are constant multiples of $-\sqrt{x} \mathbb{I}\{x \geq 0\} + \infty \mathbb{I}\{x < 0\}$, defined over the reals, in which case the Bregman divergences become $+\infty$ for $x = 0$ and $y > 0$.

Appendix F. Proof of Theorem 3

In this section, we provide a detailed proof of Theorem 3. First, we prove generalized versions of two lemmas that have appeared in several previous work; see, e.g., the work of Dekel et al. [18] and the references therein.

The first lemma is used for ADA-FTRL.

Lemma 26. Let $g \in \mathcal{H}$ and consider a proper, directionally differentiable function $r : \mathcal{H} \rightarrow \overline{\mathbb{R}}$. Define $S = \text{dom}(r)$, and let $\mathcal{X} \subset \mathcal{H}$ be a convex set such that $\mathcal{X} \cap S \neq \emptyset$. Further assume that $\text{argmin}_{x \in \mathcal{X}} \langle g, x \rangle + r(x)$ is non-empty. Then, for any $x^+ \in \text{argmin}_{x \in \mathcal{X}} \langle g, x \rangle + r(x)$ and any $x \in \mathcal{X} \cap S$,

$$+\infty > \langle g, x - x^+ \rangle + r(x) - r(x^+) \geq \mathcal{B}_r(x, x^+). \quad (\text{F.1})$$

Proof. Let $h : \mathcal{H} \rightarrow \overline{\mathbb{R}}$ be given by $h(\cdot) = \langle g, \cdot \rangle + r(\cdot)$, so that $x^+ \in \text{argmin}_{x \in \mathcal{X}} h(x)$. Note that by Lemma 23, $\text{dom}(h) = S$ and h is directionally differentiable with $h'(x; z) = \langle g, z \rangle + r'(x; z)$ for all $x \in S$ and $z \in \mathcal{H}$. Also note that $x^+ \in \mathcal{X} \cap S$ by definition. Since $x, x^+ \in \mathcal{X}$, and $\mathcal{X}$ is convex, for all $\alpha \in [0, 1]$, we have $x^+ + \alpha(x - x^+) \in \mathcal{X}$. Therefore, the optimality of x^+ over $\mathcal{X}$ implies that for all $\alpha \in (0, 1)$,

$$\frac{h(x^+ + \alpha(x - x^+)) - h(x^+)}{\alpha} \geq 0.$$

Thus, $0 \leq h'(x^+; x - x^+) = \langle g, x - x^+ \rangle + r'(x^+; x - x^+)$, and therefore $+\infty > \langle g, x - x^+ \rangle \geq -r'(x^+; x - x^+)$. Adding the real number $r(x) - r(x^+)$ to the sides completes the proof. $\square$

The second lemma is used for ADA-MD.

Lemma 27. Let $\mathcal{X}, S, g$ and r be as in Lemma 26. Let $y \in S \cap \mathcal{X}$ be such that $r'(y; \cdot - y)$ is real-valued and concave on S , that is, for all $x_1, x_2 \in S$ and all $\alpha \in [0, 1]$ for which $x_\alpha := x_1 + \alpha(x_2 - x_1) \in S$,

$$+\infty > r'(y; x_\alpha - y) \geq \alpha r'(y; x_2 - y) + (1 - \alpha) r'(y; x_1 - y) > -\infty.$$

Let $q : \mathcal{H} \rightarrow \overline{\mathbb{R}}$ be proper and directionally differentiable, with $S_q := S \cap \mathcal{X} \cap \text{dom}(q) \neq \emptyset$. Assume that $\mathcal{X}^+ := \text{argmin}_{x \in \mathcal{X}} \langle g, x \rangle + q(x) + \mathcal{B}_r(x, y)$ is non-empty, and the associated optimal value is finite. Then, for any $x^+ \in \mathcal{X}^+$ and any $x \in S_q$,

$$+\infty > \langle g, x - x^+ \rangle + q(x) - q(x^+) + \mathcal{B}_r(x, y) - \mathcal{B}_r(x^+, y) \geq \mathcal{B}_{r+q}(x, x^+). \quad (\text{F.2})$$

Proof. Let $h : \mathcal{H} \rightarrow \overline{\mathbb{R}}$ be given by $h(\cdot) = \langle g, \cdot \rangle + q + \mathcal{B}_r(\cdot, y)$, so that $x^+ \in \text{argmin}_{x \in \mathcal{X}} h(x)$. Note that by assumption, $h(x^+) < +\infty$. In addition, $\text{dom}(h) \subset S \cap \text{dom}(q)$. Thus, $x^+ \in S_q$.

Now, fix $\alpha \in (0, 1)$, and let $x_\alpha = x^+ + \alpha(x - x^+)$. If $x_\alpha \in S_q$, then $q(x_\alpha)$ and $r(x_\alpha)$ are real-valued, and by the optimality of x^+ over $\mathcal{X}$ and the concavity of $r'(y; \cdot - y)$ over S ,

$$\begin{aligned} 0 &\leq h(x_\alpha) - h(x^+) = q(x_\alpha) - q(x^+) + \langle g, x^+ + \alpha(x - x^+) - x^+ \rangle + \mathcal{B}_r(x_\alpha, y) - \mathcal{B}_r(x^+, y) \\ &= q(x_\alpha) - q(x^+) + \alpha \langle g, x - x^+ \rangle + r(x_\alpha) - r(x^+) \\ &\quad - r'(y; x_\alpha - y) + r'(y; x^+ - y) \\ &\leq q(x_\alpha) - q(x^+) + \alpha \langle g, x - x^+ \rangle + r(x_\alpha) - r(x^+) \\ &\quad - ((1 - \alpha)r'(y; x^+ - y) + \alpha r'(y; x - y)) + r'(y; x^+ - y) \\ &= q(x_\alpha) - q(x^+) + \alpha \langle g, x - x^+ \rangle + r(x_\alpha) - r(x^+) + \alpha (r'(y; x^+ - y) - r'(y; x - y)), \end{aligned}$$

Suppose, on the other hand, that $x_\alpha \notin S_q$. Then, given that by the assumption of convexity of $\mathcal{X}$, $x_\alpha \in \mathcal{X}$, we must have $x_\alpha \notin S \cap \text{dom}(q)$, so that $(r+q)(x_\alpha) = +\infty$. In addition, $r'(y; \cdot - y)$ is real-valued over S and $x, x^+ \in S$, so $r'(y; x^+ - y) - r'(y; x - y)$ is real-valued. Putting this together, we will again have that for $x_\alpha \notin S_q$,

$$0 \leq q(x_\alpha) - q(x^+) + \alpha \langle g, x - x^+ \rangle + r(x_\alpha) - r(x^+) + \alpha (r'(y; x^+ - y) - r'(y; x - y)),$$

Thus, dividing by the positive α , for all $\alpha \in (0, 1)$, we have

$$0 \leq \langle g, x - x^+ \rangle + \frac{q(x_\alpha) - q(x^+) + r(x_\alpha) - r(x^+)}{\alpha} - r'(y; x - y) + r'(y; x^+ - y).$$

Taking infimum over α , we obtain

$$\begin{aligned} 0 &\leq \langle g, x - x^+ \rangle - r'(y; x - y) + r'(y; x^+ - y) + \inf_{\alpha \in (0, 1)} \frac{q(x_\alpha) - q(x^+) + r(x_\alpha) - r(x^+)}{\alpha} \\ &\leq \langle g, x - x^+ \rangle - r'(y; x - y) + r'(y; x^+ - y) + (r+q)'(x^+; x - x^+), \end{aligned}$$

using directional differentiability of $q+r$ in the final step. Adding the real-valued equation $0 = q(x) - q(x^+) + r(x) - r(y) + r(y) - r(x^+) + (r+q)(x^+) - (r+q)(x)$, using the definition of Bregman divergence, and rearranging terms completes the proof. $\square$

We can now prove Theorem 3.

Proof of Theorem 3. First consider ADA-FTRL. For $t = 0, 1, \dots, T$, let $h_t^{\text{ftrl}} : \mathcal{H} \rightarrow \overline{\mathbb{R}}$ be given by $h_t^{\text{ftrl}}(\cdot) := \langle g_{1:t}, \cdot \rangle + q_t(\cdot) + r_{1:t}(\cdot)$, recalling that $c_{i:j} \equiv 0$ whenever $i > j$. Let $S_t = \text{dom}(r_{1:t})$. By Assumption 1, for $t = 1, 2, \dots, T$,

$$-\infty < h_{t-1}^{\text{ftrl}}(x_t) = \langle g_{1:t-1}, x_t \rangle + q_{t-1}(x_t) + r_{1:t-1}(x_t) < +\infty.$$

Therefore, $x_t \in \text{dom}(r_{1:t-1} + q_{t-1})$. In addition, by (4), $x_t \in \text{dom}(p_t)$. Thus, $x_t \in \text{dom}(r_{1:t-1} + q_{t-1} + p_t)$, that is, $x_t \in \text{dom}(r_{1:t}) = S_t$. Furthermore, $h_t^{\text{ftrl}}(x_{t+1}) < +\infty$, so $x_{t+1} \in \text{dom}(q_t) \cap \text{dom}(r_{1:t})$. Thus, $x_t, x_{t+1} \in \mathcal{X} \cap S_t \subset \bigcap_{s=1}^t (\text{dom}(q_{s-1}) \cap \text{dom}(p_s))$.

Now, for any $t = 1, 2, \dots, T$, since x_t minimizes p_t over $\mathcal{X}$, if we add p_t to the objective of the optimization above, we will still have

$$x_t \in \arg\min_{x \in \mathcal{X}} h_{t-1}^{\text{ftrl}}(x) + p_t(x) = \arg\min_{x \in \mathcal{X}} \langle g_{1:t-1}, x \rangle + r_{1:t}(x).$$

By Assumption 1, $r_{1:t}$ is directionally differentiable. Therefore, for any $t = 1, 2, \dots, T$, we can apply Lemma 26 with $g \leftarrow g_{1:t-1}$, $r \leftarrow r_{1:t}$, $x^+ \leftarrow x_t$, and $x \leftarrow x_{t+1}$, to obtain

$$-\infty < \langle g_{1:t-1}, x_t - x_{t+1} \rangle + p_{1:t}(x_t) - p_{1:t}(x_{t+1}) + q_{0:t-1}(x_t) - q_{0:t-1}(x_{t+1}) \leq -\mathcal{B}_{r_{1:t}}(x_{t+1}, x_t).$$

In the inequality above, the right-hand side cannot be equal to $-\infty$ (by Lemma 26), and all other terms are real-valued. Thus, we can sum up this inequality over $t = 1, 2, \dots, T$, to obtain

$$\begin{aligned} & - \sum_{t=1}^T \mathcal{B}_{r_{1:t}}(x_{t+1}, x_t) \\ & \geq \sum_{t=1}^T \langle g_{1:t-1}, x_t \rangle - \sum_{t=1}^T \langle g_{1:t-1}, x_{t+1} \rangle + \sum_{t=1}^T p_{1:t}(x_t) - \sum_{t=1}^T p_{1:t}(x_{t+1}) \\ & \quad + \sum_{t=1}^T q_{0:t-1}(x_t) - \sum_{t=1}^T q_{0:t-1}(x_{t+1}) \\ & = \sum_{t=0}^{T-1} \langle g_{1:t}, x_{t+1} \rangle - \sum_{t=1}^T \langle g_{1:t-1}, x_{t+1} \rangle + \sum_{t=1}^T p_{1:t}(x_t) - \sum_{t=2}^{T+1} p_{1:t-1}(x_t) \\ & \quad + \sum_{t=0}^{T-1} q_{0:t}(x_{t+1}) - \sum_{t=0}^T q_{0:t-1}(x_{t+1}) \\ & = \sum_{t=1}^{T-1} \langle g_{1:t}, x_{t+1} \rangle - \sum_{t=1}^T \langle g_{1:t-1}, x_{t+1} \rangle + \sum_{t=1}^T p_{1:t}(x_t) - \sum_{t=1}^{T+1} p_{1:t-1}(x_t) \end{aligned}$$

$$\begin{aligned}
& + \sum_{t=0}^{T-1} q_{0:t}(x_{t+1}) - \sum_{t=0}^T q_{0:t-1}(x_{t+1}) \\
& = -\langle g_{1:T}, x_{T+1} \rangle + \sum_{t=1}^T \langle g_t, x_{t+1} \rangle + \sum_{t=1}^T p_t(x_t) - p_{1:T}(x_{T+1}) - q_{0:T}(x_{T+1}) + \sum_{t=0}^T q_t(x_{t+1}) \\
& = \sum_{t=1}^T \langle g_t, x_{t+1} \rangle + \sum_{t=1}^T p_t(x_t) + \sum_{t=0}^T q_t(x_{t+1}) - \left(\langle g_{1:T}, x_{T+1} \rangle + p_{1:T}(x_{T+1}) + q_{0:T}(x_{T+1}) \right) \\
& \geq \sum_{t=1}^T \langle g_t, x_{t+1} \rangle + \sum_{t=1}^T p_t(x_t) + \sum_{t=0}^T q_t(x_{t+1}) - \left(\langle g_{1:T}, x^* \rangle + p_{1:T}(x^*) + q_{0:T}(x^*) \right) \\
& = R_T^+(x^*) + \sum_{t=1}^T p_t(x_t) + \sum_{t=0}^T q_t(x_{t+1}) - p_{1:T}(x^*) - q_{0:T}(x^*),
\end{aligned}$$

using, in the last inequality, the optimality of x_{T+1} over $\mathcal{X}$, as well as the fact that p_t, q_t are proper and all terms on the right-hand side not involving x^* are real-valued (hence the term in the parentheses involving x^* is well-defined and can be added to the rest of the expression). Now, if $x^* \notin \text{dom}(p_{1:T} + q_{0:T})$, the bound of Theorem 3 holds trivially (recalling that the Bregman divergences cannot be $+\infty$). Otherwise, $(p_{1:T} + q_{0:T})(x^*)$ is real-valued, and rearranging completes the proof for ADA-FTRL.

For ADA-MD, we start by presenting the implications of Assumption 2.

To simplify notation, let $x_0 := g_0 := 0$, and define $h_t^{\text{md}} := \langle g_t, x \rangle + q_t(x) + \mathcal{B}_{r_{1:t}}(x, x_t)$ and $S_t = \text{dom}(r_{1:t})$ for $t = 0, 1, \dots, T$ (so that $S_0 = \text{dom}(r_{1:0}) = \mathcal{H}$). Then, by Assumption 2, $h_t^{\text{md}}(x_{t+1}) < +\infty$ for all $t = 0, 1, \dots, T$, so $x_{t+1} \in \mathcal{X} \cap \text{dom}(q_t) \cap S_t$. Thus, given that $r'_{1:t}(x_t; \cdot - x_t)$ is real-valued on S_t , and $x_t \in S_t$ by assumption, $\mathcal{B}_{r_{1:t}}(x_{t+1}, x_t)$ is also real-valued.

Now, note that by the optimality of x_{T+1} , and because $h_T^{\text{md}}(x_{T+1})$ is finite, for all $x^* \in \mathcal{X}$,

$$\langle g_T, x_{T+1} - x^* \rangle \leq q_T(x^*) - q_T(x_{T+1}) + \mathcal{B}_{r_{1:T}}(x^*, x_T) - \mathcal{B}_{r_{1:T}}(x_{T+1}, x_T). \quad (\text{F.3})$$

Next, fix $t \in \{0, 1, 2, \dots, T-1\}$ and suppose that $p_{t+1}(x^*)$ is finite-valued. Then, by the definition of p_{t+1} , we have $x^* \in \mathcal{X} \cap \text{dom}(q_t)$ and $x^* \in \text{dom}(r_{t+1}) = \text{dom}(r_{1:t+1}) \subset S_t$. Furthermore, by the argument above, $x_{t+1} \in \mathcal{X} \cap S_t \cap \text{dom}(q_t)$ and $x_t \in S_t$. Thus, for all $t = 0, 1, \dots, T-1$, we can apply Lemma 27 with $g \leftarrow g_t$, $r \leftarrow r_{1:t}$, $q \leftarrow q_t$, $y \leftarrow x_t$, $x^+ \leftarrow x_{t+1}$, and $x \leftarrow x^*$, to obtain

$$\begin{aligned}
\langle g_t, x_{t+1} - x^* \rangle & \leq q_t(x^*) - q_t(x_{t+1}) + \mathcal{B}_{r_{1:t}}(x^*, x_t) - \mathcal{B}_{r_{1:t}}(x_{t+1}, x_t) \\
& \quad - \mathcal{B}_{r_{1:t}+q_t}(x^*, x_{t+1}).
\end{aligned} \quad (\text{F.4})$$

Note that this also implies that the right-hand side above cannot be $-\infty$, and only the last Bregman divergence term could be infinite. Now, since $r'_{1:t+1}(x_{t+1}; \cdot - x_{t+1})$ is real-valued on S_{t+1} , $r_{1:t} + q_t$ is directionally differentiable, and $x^* \in S_{t+1}$, by Lemma 25 we have

$$\mathcal{B}_{r_{1:t+1}}(x^*, x_{t+1}) - \mathcal{B}_{r_{1:t}+q_t}(x^*, x_{t+1}) = \mathcal{B}_{p_{t+1}}(x^*, x_{t+1}).$$

In particular, this implies that $\mathcal{B}_{p_{t+1}}(x^*, x_{t+1})$ cannot be $-\infty$. Moving the (real-valued) first term to the right-hand side, and substituting into (F.4), we have

$$\begin{aligned}
\langle g_t, x_{t+1} - x^* \rangle & \leq q_t(x^*) - q_t(x_{t+1}) + \mathcal{B}_{r_{1:t}}(x^*, x_t) - \mathcal{B}_{r_{1:t}}(x_{t+1}, x_t) + \mathcal{B}_{p_{t+1}}(x^*, x_{t+1}) \\
& \quad - \mathcal{B}_{r_{1:t+1}}(x^*, x_{t+1}).
\end{aligned}$$

In light of the above, if $p_{t+1}(x^*)$ is finite-valued for all $t = 0, 1, 2, \dots, T-1$, then summing up the above inequality for all these t s and adding (F.3), we obtain

$$\begin{aligned}
\sum_{t=0}^T \langle g_t, x_{t+1} - x^* \rangle & \leq \sum_{t=0}^{T-1} (q_t(x^*) - q_t(x_{t+1})) + \sum_{t=0}^{T-1} \mathcal{B}_{p_{t+1}}(x^*, x_{t+1}) \\
& \quad + \sum_{t=0}^{T-1} \mathcal{B}_{r_{1:t}}(x^*, x_t) - \sum_{t=0}^{T-1} \mathcal{B}_{r_{1:t+1}}(x^*, x_{t+1}) - \sum_{t=0}^{T-1} \mathcal{B}_{r_{1:t}}(x_{t+1}, x_t) \\
& \quad + q_T(x^*) - q_T(x_{T+1}) + \mathcal{B}_{r_{1:T}}(x^*, x_T) - \mathcal{B}_{r_{1:T}}(x_{T+1}, x_T)
\end{aligned}$$

$$\begin{aligned}
&= \sum_{t=0}^T (q_t(x^*) - q_t(x_{t+1})) + \sum_{t=0}^{T-1} \mathcal{B}_{p_{t+1}}(x^*, x_{t+1}) \\
&\quad + \sum_{t=1}^T \mathcal{B}_{r_{1:t}}(x^*, x_t) - \sum_{t=0}^{T-1} \mathcal{B}_{r_{1:t+1}}(x^*, x_{t+1}) - \sum_{t=1}^T \mathcal{B}_{r_{1:t}}(x_{t+1}, x_t) \\
&= \sum_{t=0}^T (q_t(x^*) - q_t(x_{t+1})) + \sum_{t=1}^T \mathcal{B}_{p_t}(x^*, x_t) - \sum_{t=1}^T \mathcal{B}_{r_{1:t}}(x_{t+1}, x_t),
\end{aligned}$$

and (9) holds.

On the other hand, if $p_{t+1}(x^*)$ is infinite for at least one t in $\{0, 1, 2, \dots, T-1\}$, then $\mathcal{B}_{p_{t+1}}(x^*, x_{t+1}) = +\infty$ by definition. Therefore, the right-hand side of (9) will be $+\infty$, given that by the argument above, $\mathcal{B}_{p_{t+1}}(x^*, x_{t+1})$ cannot be equal to $-\infty$ if $p_{t+1}(x^*)$ is finite-valued. Thus, in this case as well, the bound of (9) holds trivially, completing the proof. $\square$

References

- [1] H.B. McMahan, M. Streeter, Adaptive bound optimization for online convex optimization, in: Proceedings of the 23rd Conference on Learning Theory, 2010.
- [2] J. Duchi, E. Hazan, Y. Singer, Adaptive subgradient methods for online learning and stochastic optimization, *J. Mach. Learn. Res.* 12 (2011) 2121–2159.
- [3] N. Cesa-Bianchi, G. Lugosi, Prediction, Learning, and Games, Cambridge University Press, New York, NY, USA, 2006.
- [4] S. Shalev-Shwartz, Online learning and online convex optimization, *Found. Trends Mach. Learn.* 4 (2) (2011) 107–194.
- [5] E. Hazan, Introduction to online convex optimization, *Found. Trends Optim.* 2 (3–4) (2016) 157–325.
- [6] A. Rakhlin, K. Sridharan, Online learning with predictable sequences, in: Conference on Learning Theory, 2013, pp. 993–1019.
- [7] S. Rakhlin, K. Sridharan, Optimization, learning, and games with predictable sequences, in: Advances in Neural Information Processing Systems, 2013, pp. 3066–3074.
- [8] C.-K. Chiang, T. Yang, C.-J. Lee, M. Mahdavi, C.-J. Lu, R. Jin, S. Zhu, Online optimization with gradual variations, in: Conference on Learning Theory, 2012.
- [9] M. Mohri, S. Yang, Accelerating online convex optimization via adaptive prediction, in: Proceedings of the 19th International Conference on Artificial Intelligence and Statistics, 2016, pp. 848–856.
- [10] P. Kamalaruban, Improved optimistic mirror descent for sparsity and curvature, preprint, arXiv:1609.02383.
- [11] J. Kivinen, M.K. Warmuth, Exponentiated gradient versus gradient descent for linear predictors, *Inf. Comput.* 132 (1) (1997) 1–63.
- [12] B. Kulis, P.L. Bartlett, Implicit online learning, in: Proceedings of the 27th International Conference on Machine Learning, ICML-10, 2010, pp. 575–582.
- [13] L. Xiao, Dual averaging method for regularized stochastic learning and online optimization, in: Advances in Neural Information Processing Systems, 2009, pp. 2116–2124.
- [14] J.C. Duchi, S. Shalev-Shwartz, Y. Singer, A. Tewari, Composite objective mirror descent, in: Conference on Learning Theory, 2010, pp. 14–26.
- [15] S. Sra, A.W. Yu, M. Li, A.J. Smola, AdaDelay: delay adaptive distributed stochastic optimization, in: Proceedings of the 19th International Conference on Artificial Intelligence and Statistics, AISTATS, 2016, pp. 957–965.
- [16] A. Juditsky, A. Nemirovski, C. Tauvel, et al., Solving variational inequalities with stochastic mirror-prox algorithm, *Stoch. Syst.* 1 (1) (2011) 17–58.
- [17] G. Lan, An optimal method for stochastic composite optimization, *Math. Program.* 133 (1) (2012) 365–397.
- [18] O. Dekel, R. Gilad-Bachrach, O. Shamir, L. Xiao, Optimal distributed online prediction using mini-batches, *J. Mach. Learn. Res.* 13 (Jan) (2012) 165–202.
- [19] F. Orabona, K. Crammer, N. Cesa-Bianchi, A generalized online mirror descent with applications to classification and regression, *Mach. Learn.* 99 (3) (2015) 411–435.
- [20] H.B. McMahan, A survey of algorithms and analysis for adaptive online learning, *J. Mach. Learn. Res.* 18 (90) (2017) 1–50.
- [21] S. Shalev-Shwartz, S.M. Kakade, Mind the duality gap: logarithmic regret algorithms for online optimization, in: Advances in Neural Information Processing Systems, 2009, pp. 1457–1464.
- [22] E. Hazan, A. Agarwal, S. Kale, Logarithmic regret algorithms for online convex optimization, *Mach. Learn.* 69 (2) (2007) 169–192.
- [23] I. Goodfellow, Y. Bengio, A. Courville, Deep Learning, MIT Press, 2016.
- [24] R. Ge, J.D. Lee, T. Ma, Matrix completion has no spurious local minimum, in: Advances in Neural Information Processing Systems, vol. 29, 2016, pp. 2973–2981.
- [25] S. Bhojanapalli, B. Neyshabur, N. Srebro, Global optimality of local search for low rank matrix recovery, in: Advances in Neural Information Processing Systems, vol. 29, 2016, pp. 3873–3881.
- [26] Y. Nesterov, B.T. Polyak, Cubic regularization of Newton method and its global performance, *Math. Program.* 108 (1) (2006) 177–205.
- [27] J.C.H. Lee, P. Valiant, Optimizing star-convex functions, in: IEEE 57th Annual Symposium on Foundations of Computer Science, FOCS, 2016, pp. 603–614.
- [28] M. Hardt, T. Ma, B. Recht, Gradient descent learns linear dynamical systems, preprint, arXiv:1609.05191.
- [29] M. Telgarsky, S. Dasgupta, Agglomerative Bregman clustering, in: Proceedings of the 29th International Conference on Machine Learning – ICML, 2012.
- [30] K.C. Kiwiel, Proximal minimization methods with generalized Bregman functions, *SIAM J. Control Optim.* 35 (4) (1997) 1142–1168.
- [31] D.P. Bertsekas, S.E. Shreve, Stochastic Optimal Control: The Discrete Time Case, vol. 23, Academic Press, New York, 1978.
- [32] A. Rakhlin, O. Shamir, K. Sridharan, Making gradient descent optimal for strongly convex stochastic optimization, in: Proceedings of the 29th International Conference on Machine Learning, ICML’12, 2012, pp. 1571–1578.
- [33] O. Shamir, T. Zhang, Stochastic gradient descent for non-smooth optimization: convergence results and optimal averaging schemes, in: Proceedings of the 30th International Conference on Machine Learning, ICML’13, in: JMLR Workshop and Conference Proceedings, vol. 28, 2013, pp. 71–79.
- [34] E. Hazan, S. Kale, Beyond the regret minimization barrier: optimal algorithms for stochastic strongly-convex optimization, *J. Mach. Learn. Res.* 15 (1) (2014) 2489–2512.
- [35] Z. Zhou, P. Mertikopoulos, N. Bambos, S. Boyd, P.W. Glynn, Stochastic mirror descent in variationally coherent optimization problems, in: Advances in Neural Information Processing Systems, vol. 30, 2017, pp. 7040–7049.
- [36] J.-K. Wang, J.D. Abernethy, Acceleration through optimistic no-regret dynamics, in: Advances in Neural Information Processing Systems, vol. 31, 2018, pp. 3824–3834.
- [37] Z. Allen-Zhu, Natasha 2: faster non-convex optimization than SGD, in: Advances in Neural Information Processing Systems, vol. 31, 2018, pp. 2675–2686.

- [38] P. Joulani, A. György, Cs. Szepesvári, Delay-tolerant online convex optimization: unified analysis and adaptive-gradient algorithms, in: Proceedings of the 30th Conference on Artificial Intelligence, AAAI-16, 2016.
- [39] A. Saha, P. Jain, A. Tewari, The interplay between stability and regret in online learning, preprint, arXiv:1211.6158.
- [40] A. Agarwal, J. Duchi, Distributed delayed stochastic optimization, in: J. Shawe-Taylor, R. Zemel, P. Bartlett, F. Pereira, K. Weinberger (Eds.), Advances in Neural Information Processing Systems (NIPS), vol. 24, 2011, pp. 873–881.
- [41] A. Nemirovsky, D. Yudin, Problem Complexity and Method Efficiency in Optimization, Wiley, Chichester, New York, 1983.
- [42] A. Beck, M. Teboulle, Mirror descent and nonlinear projected subgradient methods for convex optimization, Oper. Res. Lett. 31 (3) (2003) 167–175.
- [43] M.K. Warmuth, A.K. Jagota, Continuous and discrete-time nonlinear gradient descent: relative loss bounds and convergence, in: Electronic Proceedings of the 5th International Symposium on Artificial Intelligence and Mathematics, 1997.
- [44] Y. Nesterov, Primal-dual subgradient methods for convex problems, Math. Program. 120 (1) (2009) 221–259.
- [45] H.H. Bauschke, P.L. Combettes, Convex Analysis and Monotone Operator Theory in Hilbert Spaces, Springer Science & Business Media, 2011.
- [46] M.K. Warmuth, A. Jagota, Continuous and discrete-time nonlinear gradient descent: relative loss bounds and convergence, in: Fifth International Symposium on Artificial Intelligence and Mathematics, Florida, 1998.
- [47] M. Zinkevich, Online convex programming and generalized infinitesimal gradient ascent, in: Proceedings of the Twentieth International Conference on Machine Learning, 2003, pp. 928–936.
- [48] A. Agarwal, M.J. Wainwright, P.L. Bartlett, P.K. Ravikumar, Information-theoretic lower bounds on the oracle complexity of convex optimization, in: Advances in Neural Information Processing Systems, 2009, pp. 1–9.
- [49] R. Johnson, T. Zhang, Accelerating stochastic gradient descent using predictive variance reduction, in: Advances in Neural Information Processing Systems, 2013, pp. 315–323.
- [50] Z. Allen-Zhu, Y. Yuan, Improved SVRG for non-strongly-convex or sum-of-non-convex objectives, in: Proceedings of the 33rd International Conference on Machine Learning, ICML '16, 2016, full version available at <http://arxiv.org/abs/1506.01972>, 2016.
- [51] H. Karimi, J. Nutini, M.W. Schmidt, Linear convergence of gradient and proximal-gradient methods under the Polyak-Łojasiewicz condition, in: Proceedings of ECML/PKDD, 2016, pp. 795–811.
- [52] E. Hazan, K.Y. Levy, S. Shalev-Shwartz, Beyond convexity: stochastic quasi-convex optimization, in: Advances in Neural Information Processing Systems (NIPS 28), vol. 28, 2015, pp. 1594–1602.
- [53] K.Y. Levy, Online to offline conversions and adaptive minibatch sizes, preprint, arXiv:1705.10499.
- [54] B.T. Polyak, Gradient methods for minimizing functionals, Zh. Vychisl. Mat. Mat. Fiz. 3 (1963) 643–653.
- [55] S. Łojasiewicz, A topological property of real analytic subsets, in: Coll. du CNRS, Les équations aux dérivées partielles, vol. 117, 1963, pp. 87–89.
- [56] H.B. McMahan, Follow-the-regularized-leader and mirror descent: equivalence theorems and L1 regularization, in: Proceedings of the Fourteenth International Conference on Artificial Intelligence and Statistics, AISTATS 2011, Fort Lauderdale, USA, April 11–13, 2011, 2011, pp. 525–533.