

# TRANSIENT DYNAMICS OF ABSORBED SINGULAR DIFFUSIONS

MIN JI, WEIWEI QI, ZHONGWEI SHEN, AND YINGFEI YI

*Dedicated to the memory of Professor Geneviève Raugel*

ABSTRACT. We consider a class of one-dimensional absorbed diffusion processes with small singular noises that exhibit multi-scale dynamics in the sense that typical trajectories of the solution process first quickly approach to transient states, captured by quasi-stationary distributions (QSDs), then stay with transient states for a very long period, and finally deviate from transient states and slowly relax to the absorbing state. The main purpose of the present paper is to give qualitative characterizations of such multi-scale dynamics with particular interest in the intriguing transient dynamics governed by transient states. This is achieved by the establishment of (i) noise-vanishing asymptotics of the first eigenvalue and the gap between the first and second eigenvalues of the generator that are respectively the rates for solutions to get absorbed by the absorbing state and attracted by QSDs; (ii) sophisticated estimates quantifying the distance between solutions and convex combinations of QSDs and the absorbing state. Applications to stochastic models arising in chemical reactions and population dynamics are discussed.

## CONTENTS

1. <b>Introduction</b>	2
2. <b>Preliminaries</b>	8
2.1. The potential function	8
2.2. Semi-classical Schrödinger operators	9
2.3. Spectrum and semigroup	11
2.4. Uniform estimate of the first eigenfunctions	12
2.5. QSD: existence, uniqueness and convergence	13
3. <b>Exponential asymptotic of the first eigenvalue</b>	13
4. <b>Exponential asymptotics of other eigenvalues</b>	17
4.1. Upper bound	17
4.2. Auxiliary eigenvalue asymptotics	19
4.3. Lower bound	20
5. <b>Non-exponential asymptotics of eigenvalues</b>	25
5.1. Asymptotics of Schrödinger eigenvalues	25
5.2. Proof of Theorem B	25

---

2010 *Mathematics Subject Classification*. Primary 60J60, 60J70, 34F05; secondary 92D25, 60H10.

*Key words and phrases*. Diffusion process, degeneracy, singularity, absorbing state, quasi-stationary distribution, extinction rate, transient dynamics, transient state, eigenvalue asymptotic, chemical reaction, population dynamics.

M.J. was partially supported by NSFC grant 11571344. W.Q. was partially supported by a postdoctoral fellowship from the University of Alberta. Z.S. was partially supported by a start-up grant from the University of Alberta, NSERC RGPIN-2018-04371 and NSERC DGEER-2018-00353. Y.Y. was partially supported by NSERC RGPIN-2020-04451, PIMS CRG grant, a faculty development grant from the University of Alberta, and a Scholarship from Jilin University.

5.3. Upper bounds of eigenvalues	26
6. Qualitative characterizations of multi-scale dynamics	27
7. Applications	31
Acknowledgements	33
Appendix A. Asymptotic of the first eigenfunctions	33
References	35

## 1. Introduction

Transient dynamics, roughly understood as interesting and important dynamical behaviors of real or artificial systems that display only over finite time scales, arise naturally in chemical reactions (see e.g. [52, 57]), population dynamics (see e.g. [25, 26, 47]) and many other scientific areas (see e.g. [39]). In a closed chemical system, chemical oscillations could last hours and even longer, while the whole system must eventually relax to the thermal equilibrium due to the dissipation of heat and energy. In population dynamics, although the eventual loss of diversity or extinction is inevitable due to limited resources, mortality, etc., species of large numbers typically coexist or persist for a very long period in comparison to human timescales. In spite of the ubiquity of transient dynamics in multi-scale systems, there have not been many mathematically rigorous studies.

The present paper is devoted to the investigation of transient dynamics of a class of multi-scale diffusion processes arising in chemical reactions, population dynamics, etc. More precisely, we consider the following family of stochastic differential equations (SDEs):

$$dx = b(x)dt + \epsilon\sqrt{a(x)}dW_t, \quad x \in [0, \infty), \quad (1.1)$$

where  $0 < \epsilon \ll 1$  is a parameter,  $b : [0, \infty) \rightarrow \mathbb{R}$ ,  $a : [0, \infty) \rightarrow [0, \infty)$  and  $W_t$  is the standard one-dimensional Wiener process on some probability space. SDEs of the form (1.1) are often derived as diffusion approximations of re-scaled Markov jump processes that model the evolution of chemical substrates or species of large numbers [19, 1]. In Remark 7.1, we give some comments on the dynamics between re-scaled Markov jump processes and their diffusion approximations. Throughout this paper, we make the following standard assumptions on the coefficients  $a$  and  $b$ .

**(H)** The functions  $b : [0, \infty) \rightarrow \mathbb{R}$  and  $a : [0, \infty) \rightarrow [0, \infty)$  satisfy the following conditions:

- (1)  $b \in C^1([0, \infty))$ ,  $b(0) = 0$ ,  $b'(0) > 0$ , and  $\limsup_{x \rightarrow \infty} b(x) < 0$ ;
- (2)  $a \in C^2([0, \infty))$ ,  $a(0) = 0$ ,  $a'(0) > 0$ , and  $a > 0$  on  $(0, \infty)$ ;
- (3)  $\lim_{x \rightarrow \infty} \frac{b^2(x)}{a(x)} = \infty$ , and  $\limsup_{x \rightarrow \infty} \frac{\max\{a(x), |a'(x)|, |a''(x)|, |b'(x)|\}}{|b(x)|} < \infty$ .

Clearly,  $b$  is a generalized logistic growth rate function, and the noise coefficient  $\epsilon\sqrt{a}$  vanishes and is singular at 0. This latter fact causes substantial troubles in the analysis of (1.1). The reader is directed to Section 7 for examples satisfying these assumptions. For the time being, it is beneficial to keep in mind the typical one:  $dx = x(1-x)dt + \epsilon\sqrt{x}dW_t$ .

We see from **(H)** that the state 0, often referred to as the extinction state, is an absorbing state of the SDE (1.1). Denote by  $X_t^\epsilon$  the stochastic process generated by solutions of (1.1), and by  $T_0^\epsilon$  the first time that  $X_t^\epsilon$  hits 0, namely,  $T_0^\epsilon = \inf\{t \geq 0 : X_t^\epsilon = 0\}$ . Under **(H)**, it is a routine task to check (see e.g. [31]) that  $\mathbb{P}_x^\epsilon[T_0^\epsilon < \infty] = 1$  for all  $x \in [0, \infty)$ , where  $\mathbb{P}_x^\epsilon$  is the law of  $X_t^\epsilon$  with initial distribution  $\mu$  and  $\mathbb{P}_x^\epsilon := \mathbb{P}_{\delta_x}^\epsilon$  for the Dirac measure  $\delta_x$  at  $x$ . That is, trajectories or sample paths of  $X_t^\epsilon$  reach the extinction state 0 in finite time almost surely, implying the non-existence of a stationary

distribution with positive concentration on  $(0, \infty)$ . This is in sharp contrast to the eventual dynamics of the unperturbed mean-field ODE:

$$\dot{x} = b(x), \quad x \in (0, \infty), \quad (1.2)$$

whose solutions are attracted by the global attractor  $\mathcal{A}$ , which is a compact interval with its left and right endpoints being respectively the smallest positive zero and largest zero of  $b$ , and has the compact dissipativity property in the sense that for each compact  $K \subset (0, \infty)$ ,  $\sup_{x_0 \in K} \text{dist}(x_t, \mathcal{A}) \rightarrow 0$  as  $t \rightarrow \infty$ , where  $x_t$  is the solution of (1.2). On the contrary, the sample path large deviation principle (see e.g. [21, 15]) says that  $X_t^\epsilon$  stays close to the solutions of (1.2) during a finite time period, which is long enough to ensure that  $X_t^\epsilon$  spends considerable amount of time with the global attractor  $\mathcal{A}$ . Hence,  $X_t^\epsilon$  exhibits multi-scale dynamics. Our goal is to characterize the multi-scale dynamics of  $X_t^\epsilon$ , in which, the transient dynamics, living on the timescale that is longer than the large deviation timescale and shorter than the extinction timescale, is of particular interest and importance.

Because of the almost sure extinction in finite time of  $X_t^\epsilon$ , it is natural to examine  $X_t^\epsilon$  before it hits the extinction state 0 in order to understand in particular the transient dynamics of  $X_t^\epsilon$ . Quasi-stationary distributions (see e.g. [44, 13]) have been widely used in literature to achieve this goal.

**Definition 1.1** (Quasi-stationary distribution). *A Borel probability measure  $\mu_\epsilon$  on  $(0, \infty)$  is called a quasi-stationary distribution (QSD) of  $X_t^\epsilon$  if*

$$\mathbb{P}_{\mu_\epsilon}^\epsilon [X_t^\epsilon \in B | t < T_0^\epsilon] = \mu_\epsilon(B), \quad \forall t \geq 0, \quad B \in \mathcal{B}((0, \infty)),$$

where  $\mathcal{B}((0, \infty))$  is the Borel  $\sigma$ -algebra of  $(0, \infty)$ .

General theory of QSDs (see e.g. [44, 13]) tells that each QSD  $\mu_\epsilon$  of  $X_t^\epsilon$  is associated with a unique positive number  $\lambda_\epsilon$ , often called the *extinction rate*, such that if  $X_0^\epsilon$  is distributed according to  $\mu_\epsilon$ , then  $T_0^\epsilon$  is exponentially distributed with rate  $\lambda_\epsilon$ , namely,  $\mathbb{P}_{\mu_\epsilon}^\epsilon [T_0^\epsilon > t] = e^{-\lambda_\epsilon t}$  for all  $t \geq 0$ .

It is shown in [5] (also see Proposition 2.3) that  $X_t^\epsilon$  admits a unique QSD  $\mu_\epsilon$  with the associated extinction rate  $\lambda_{\epsilon,1}$  being the first eigenvalue of  $-\mathcal{L}_\epsilon$ , where  $\mathcal{L}_\epsilon$  is essentially the generator of  $X_t^\epsilon$ . Moreover,  $X_t^\epsilon$  conditioned on non-absorption, i.e., the event  $[t < T_0^\epsilon]$ , converges to the QSD  $\mu_\epsilon$  exponentially fast with rate  $\lambda_{\epsilon,2} - \lambda_{\epsilon,1}$ , where  $\lambda_{\epsilon,2}$  is the second eigenvalue of  $-\mathcal{L}_\epsilon$ . Thus,  $X_t^\epsilon$  is simultaneously exponentially attracted by the extinction state 0 and the QSD  $\mu_\epsilon$  with rates  $\lambda_{\epsilon,1}$  and  $\lambda_{\epsilon,2} - \lambda_{\epsilon,1}$ , respectively, suggesting the importance of comparing  $\lambda_{\epsilon,1}$  with  $\lambda_{\epsilon,2} - \lambda_{\epsilon,1}$ . By doing so, we roughly conclude:

- (i) if  $\lambda_{\epsilon,1} < \lambda_{\epsilon,2} - \lambda_{\epsilon,1}$ , then  $X_t^\epsilon$  first approaches to and then spend considerable amount of time with the QSD  $\mu_\epsilon$  before eventually relaxing to the extinction state; in this case,  $X_t$  exhibits multi-scale dynamics, and  $\mu_\epsilon$  is a true physical state governing the transient dynamics;
- (ii) if  $\lambda_{\epsilon,1} > \lambda_{\epsilon,2} - \lambda_{\epsilon,1}$ , then most trajectories of  $X_t^\epsilon$  have already reached the extinction state before the conditioned process approaches to  $\mu_\epsilon$ , saying that  $X_t^\epsilon$  exhibits simple extinction dynamics, and the QSD  $\mu_\epsilon$  does not play an interesting role.

The scenario described in situation (i) is expected. To obtain this piece of information, we study the asymptotics of  $\lambda_{\epsilon,1}$  and  $\lambda_{\epsilon,2}$  as  $\epsilon \rightarrow 0$ .

To put our study in a rigorous framework, we consider for each  $0 < \epsilon \ll 1$  the symmetric form  $\mathcal{E}_\epsilon : C_0^\infty((0, \infty)) \times C_0^\infty((0, \infty)) \rightarrow \mathbb{R}$  defined by

$$\mathcal{E}_\epsilon(\phi, \psi) = \frac{\epsilon^2}{2} \int_0^\infty a \phi' \psi' u_\epsilon^G dx, \quad \forall \phi, \psi \in C_0^\infty((0, \infty)),$$

where  $u_\epsilon^G = \frac{1}{a} e^{\frac{2}{\epsilon^2} \int_0^\bullet \frac{b(s)}{a(s)} ds}$  is the non-integrable Gibbs density. More precisely,  $u_\epsilon^G$  is the unique (up to constant multiplication) solution of the stationary Fokker-Planck equation  $\frac{\epsilon^2}{2}(au)'' - (bu)' = 0$  in  $(0, \infty)$ , and satisfies  $\int_0^\infty u_\epsilon^G dx = \infty$  due to its singularity near 0, that is,  $u_\epsilon^G$  behaves like  $\frac{1}{x}$  near  $x = 0$ . It is known (see e.g. [23]) that  $\mathcal{E}_\epsilon$  is Markovian and closable. Its smallest closed extension, still denoted by  $\mathcal{E}_\epsilon$ , is a Dirichlet form with domain  $D(\mathcal{E}_\epsilon)$  being the closure of  $C_0^\infty((0, \infty))$  under the norm  $\|u\|_{D(\mathcal{E}_\epsilon)}^2 := \|u\|_{L^2(u_\epsilon^G)}^2 + \mathcal{E}_\epsilon(u, u)$ . Denote by  $\mathcal{L}_\epsilon$  the non-positive self-adjoint operator in the weighted space  $L^2(u_\epsilon^G) := L^2((0, \infty), u_\epsilon^G(x)dx)$  associated to  $\mathcal{E}_\epsilon$  such that

$$\mathcal{E}_\epsilon(\phi, \psi) = \langle -\mathcal{L}_\epsilon \phi, \psi \rangle_{L^2(u_\epsilon^G)}, \quad \phi, \psi \in D(\mathcal{L}_\epsilon), \quad (1.3)$$

where

$$D(\mathcal{L}_\epsilon) := \{u \in D(\mathcal{E}_\epsilon) : \exists f \in L^2(u_\epsilon^G) \text{ s.t. } \mathcal{E}_\epsilon(u, \phi) + \langle u, \phi \rangle_{L^2(u_\epsilon^G)} = \langle f, \phi \rangle_{L^2(u_\epsilon^G)}, \forall \phi \in D(\mathcal{E}_\epsilon)\}$$

is the domain of  $\mathcal{L}_\epsilon$ . It is informative to mention

$$\mathcal{L}_\epsilon \phi = \frac{\epsilon^2}{2} a \phi'' + b \phi', \quad \forall \phi \in C_0^\infty((0, \infty)),$$

that is,  $\mathcal{L}_\epsilon$  is a self-adjoint extension in  $L^2(u_\epsilon^G)$  of the generator of  $X_t^\epsilon$  or (1.1). In particular,  $\mathcal{L}_\epsilon$  is degenerate at the extinction state 0.

We show in Proposition 2.1 that the spectrum of  $-\mathcal{L}_\epsilon$  consists of simple eigenvalues  $\lambda_{\epsilon, i}$ ,  $i \in \mathbb{N} := \{1, 2, 3, \dots\}$  listed as follows:  $0 < \lambda_{\epsilon, 1} < \lambda_{\epsilon, 2} < \lambda_{\epsilon, 3} < \dots \rightarrow \infty$ . To study their asymptotics, in particular, that of  $\lambda_{\epsilon, 1}$  and  $\lambda_{\epsilon, 2}$ , as  $\epsilon \rightarrow 0$ , we follow [43] to define *valleys* revealing the geometry of the potential function

$$V(x) = -2 \int_0^x \frac{b(s)}{a(s)} ds, \quad x \in (0, \infty). \quad (1.4)$$

Some basic properties of  $V$  are collected in Lemma 2.1. For  $I \subset (0, \infty)$ , set  $\gamma(I) := \sup_I V - \inf_I V$ .

**Definition 1.2.** *An open interval  $I \subset (0, \infty)$  is called a valley if it is one of the connected components of the sub-level set  $\{x \in (0, \infty) : V(x) < \rho\}$  and satisfies  $V(\partial I) = \rho$  for some  $\rho \in \mathbb{R}$ . We say  $I \subset (0, \infty)$  a  $r$ -valley if it is a valley and has depth  $r$ , namely,  $\gamma(I) = r$ .*

As  $V(0+) = 0$ , it is possible that a connected component of  $\{x \in (0, \infty) : V(x) < \rho\}$  for  $\rho > 0$  takes the form  $(0, \ell)$  with  $V(\ell) = \rho > 0 = V(0+)$ . The condition  $V(\partial I) = \rho$  in the definition of a valley is to exclude such possibilities. Hence,  $(0, \ell)$  is a valley if and only if  $\ell$  is the smallest positive zero of  $V$ .

For  $r > 0$ , let  $N(r)$  be the number of  $r$ -valleys. Clearly,  $r \mapsto N(r)$  is a non-negative, non-increasing, left-continuous and bounded step function on  $(0, \infty)$ . For each  $i \in \mathbb{N}$ , we define

$$r_i := \inf \{r > 0 : N(r) < i\}. \quad (1.5)$$

Since  $V(0+) = 0$  and  $V'(x) > 0$  for  $x \gg 1$ , the deepest valley must have finite depth. As a result, for each  $i \in \mathbb{N}$ , there always exists  $r \in (0, \infty)$  such that  $N(r) < i$ , and hence,  $r_i$  is well-defined. Intuitively,  $r_i$ ,  $i \in \mathbb{N}$  are the points where  $N(r)$  has jump discontinuities.

**Remark 1.1.** *Using the properties that  $V(0+) = 0$  and  $V'(x) > 0$  for  $x \gg 1$ , we make some remarks about  $r_i$ ,  $i \in \mathbb{N}$ .*

- (1) *There is at least one  $r_1$ -valley, but not a single  $r$ -valley with  $r > r_1$  exists. For  $r \in (0, r_1]$  satisfying  $N(r) \geq i$ , there holds  $r_i = \sup \{r > 0 : N(r) \geq i\}$ . Otherwise,  $r_i = 0$ .*

(2) Obviously,  $r_1 \geq r_2 \geq r_3 \geq \dots \geq 0$  and  $r_1 > 0$ . Moreover,  $r_1$  has the representation:

$$r_1 = \sup_{x \in (0, \infty)} \left[ \inf_{\xi \in C_x} \sup_{t \in [0, 1]} V(\xi(t)) - V(x) \right],$$

where for each  $x \in (0, \infty)$ ,

$$C_x = \{\xi : [0, 1] \rightarrow [0, \infty) : \xi \text{ is continuous and satisfies } \xi(0) = x \text{ and } \xi(1) = 0\}.$$

(3) Note that  $r_1 > r_2$  if and only if there is a unique  $r_1$ -valley, and  $r_2 = 0$  if and only if  $b$  changes its sign once in  $(0, \infty)$  (or equivalently,  $V$  has a single well).

(4) There is  $i_* \in \mathbb{N}$  such that  $r_{i_*} > 0 = r_{i_*+1}$ . This  $i_*$  is nothing but the number of wells that  $V$  has. Hence, if  $i_* \in \mathbb{N} \setminus \{1\}$ , the ODE (1.2) has multiple stable equilibria separated by unstable ones, and thus,  $X_t^\epsilon$  exhibits metastability (see e.g. [24, 48]).

For readers' convenience, we illustrate the definition of valleys and  $r_i$ ,  $i = 1, 2, \dots$  by means of Figure 1, where two typical double-well potential functions  $V$  are given. It is easy to check that for the potential function in Figure 1(a):

- if  $0 < r \leq V(x_3) - V(x_4)$ , there are exactly two  $r$ -valleys; one is contained in  $(x_1, x_3)$ , and the other is contained in  $(x_3, x_5)$ ;
- if  $V(x_3) - V(x_4) < r \leq -V(x_2)$ , there is a unique  $r$ -valley; more precisely, if  $V(x_3) - V(x_4) < r \leq V(x_3) - V(x_2)$ , the unique  $r$ -valley is contained in  $(x_1, x_3)$ , and if  $V(x_3) - V(x_2) < r \leq -V(x_2)$ , it contains  $(x_1, x_5)$  and is contained in  $(0, x_6)$ ;
- if  $r > -V(x_2)$ , no  $r$ -valley exists.

Then, according to (1.5), we find  $r_1 = -V(x_2)$  and  $(0, x_6)$  is the unique  $r_1$ -valley,  $r_2 = V(x_3) - V(x_4)$  and  $r_3 = r_4 = \dots = 0$ . Similarly, for the potential function  $V$  in Figure 1(b), we find  $r_1 = V(x_3) - V(x_4)$  and  $(x_3, x_5)$  is the unique  $r_1$ -valley,  $r_2 = -V(x_2)$  and  $r_3 = r_4 = \dots = 0$ .

Our first result studies exponential asymptotics of the eigenvalues  $\lambda_{\epsilon, i}$ ,  $i \in \mathbb{N}$  as  $\epsilon \rightarrow 0$ .

**Theorem A.** *Assume (H). Then,  $\lim_{\epsilon \rightarrow 0} \epsilon^2 \ln \lambda_{\epsilon, i} = -r_i$  for each  $i \in \mathbb{N}$ .*

We mention that exponential asymptotics of eigenvalues associated to non-degenerate, regular and reversible diffusion processes with small noises have been studied in [21] by large deviation methods, and in [43, 45] by variational formulas. When local minimal points of the potential function are non-degenerate, much refined results are established in [4] by a potential-theoretic approach developed in [3], and in [27, 28, 40] by a Witten complex approach. For the asymptotic of the first eigenvalue, special attention has been attracted due to its connections with the first exit time or first passage time (see e.g. [56, 22, 16, 35, 36, 17, 30, 42]). Because of the degeneracy and singularity of noises at the extinction state 0 resulting in in particular the degeneracy of  $\mathcal{L}_\epsilon$  and the non-integrable singularity of  $u_\epsilon^G$  at 0, known results do not directly apply in our case. Addressing difficulties caused by these unpleasant facts, we manage to adapt approaches in [30, 42, 43, 45] all based on variational formulas for eigenvalues.

Since  $r_1 > 0$  as mentioned in Remark 1.1 (2), Theorem A implies that  $\lambda_{\epsilon, 1}$  is exponentially small in  $\epsilon$ . The exponential smallness of  $\lambda_{\epsilon, 2}$  in  $\epsilon$  follows if  $r_2 > 0$ . But,  $r_2 = 0$  takes place in typical applications. For instance, it is the case when  $b$  is a logistic growth rate function, that is,  $b(x) = b_1x - b_2x^2$  for  $b_1, b_2 > 0$ . In this case, Theorem A only says that the leading asymptotic of  $\lambda_{\epsilon, 2}$  in  $\epsilon$  is non-exponential, and does not tell any further information about  $\lambda_{\epsilon, 2}$ . In Proposition 5.2, upper bounds of the eigenvalues  $\lambda_{\epsilon, i}$ ,  $i \in \mathbb{N}$  are established to complement Theorem A.




FIGURE 1. Two typical double-well potential functions

Our next result aims at investigating in particular the non-exponential asymptotic of  $\lambda_{\epsilon,2}$  as  $\epsilon \rightarrow 0$  when  $r_2 = 0$ . Note from Remark 1.1 (3) that  $r_2 = 0$  if and only if  $b$  only changes its sign once.

**Theorem B.** *Assume (H). Suppose, in addition, that*

- *there is  $x^* \in (0, \infty)$  such that  $b > 0$  on  $(0, x^*)$ ,  $b < 0$  on  $(x^*, \infty)$  and  $b'(x^*) < 0$ ;*
- *$b$  is twice continuously differentiable near  $x^*$ .*

*Then,  $\lim_{\epsilon \rightarrow 0} \lambda_{\epsilon,i} = b'(x^*)(1 - i)$  for each  $i \in \mathbb{N}$ .*

We remark that Theorem B has been essentially proven in [14, 54, 37]. Indeed, if we consider the Schrödinger operator  $-\mathcal{L}_\epsilon^S$  that is unitarily equivalent to  $-\mathcal{L}_\epsilon$  (see Subsection 2.2), then the operator  $-2\epsilon^2 \mathcal{L}_\epsilon^S$  is in the form used for studying semi-classical limits of eigenvalues, except that there is an extra term of negligible order  $\epsilon^4$ , which however blows up at the extinction state 0. Fortunately, this term can be absorbed into the main part of the operator, allowing us to apply classical results to Schrödinger operators that are comparable in the sense of quadratic forms to the original one. See Subsection 5.2 for more details.

Combining Theorem A and Theorem B, we obtain the following result.

**Corollary A.** *Assume (H). Suppose, in addition, that*

- *there is  $x^* \in (0, \infty)$  such that  $b > 0$  on  $(0, x^*)$ ,  $b < 0$  on  $(x^*, \infty)$  and  $b'(x^*) < 0$ ;*
- *$b$  is twice continuously differentiable near  $x^*$ .*

*Then,  $\lim_{\epsilon \rightarrow 0} \epsilon^2 \ln \lambda_{\epsilon,1} = -2 \int_0^{x^*} \frac{b(s)}{a(s)} ds$  and  $\lim_{\epsilon \rightarrow 0} \lambda_{\epsilon,i} = b'(x^*)(1 - i)$  for each  $i \in \mathbb{N} \setminus \{1\}$ .*

We discuss some implications of Theorem A and Theorem B.

- The exponential smallness of  $\lambda_{\epsilon,1}$  asserts the exponentially long lifetime of  $X_t^\epsilon$ . On average, it takes  $\frac{1}{\lambda_{\epsilon,1}}$  long for  $X_t^\epsilon$  to get absorbed by the extinction state 0.
- As long as  $r_1 > r_2$  (if and only if there is a unique  $r_1$ -valley), Theorem A yields that  $\lambda_{\epsilon,1} < \lambda_{2,\epsilon} - \lambda_{1,\epsilon} (\approx \lambda_{\epsilon,2})$  for all small  $\epsilon$ , implying the multi-scale nature of the dynamics of  $X_t^\epsilon$  as mentioned earlier. When  $r_2 > 0$ , the gap  $\lambda_{2,\epsilon} - \lambda_{1,\epsilon}$  is also exponentially small, saying that  $X_t^\epsilon$  needs to spend exponentially long time ( $\frac{1}{\lambda_{2,\epsilon} - \lambda_{1,\epsilon}}$  long on average) in order to get close to the QSD  $\mu_\epsilon$ . This happens because of the metastability as mentioned in Remark 1.1 (4) that trajectories of  $X_t^\epsilon$  are trapped in potential wells of  $V$  for exponentially long times before transitions can be made. Nonetheless,  $X_t^\epsilon$  begins to interact with  $\mu_\epsilon$  at a much earlier time.
- In the situation of Theorem B or Corollary A, the gap  $\lambda_{\epsilon,2} - \lambda_{\epsilon,1}$  is of order  $O(1)$ , and hence,  $X_t^\epsilon$  quickly approaches to the QSD  $\mu_\epsilon$ .

Our last result, establishing an upper bound on the distance between  $X_t^\epsilon$  and a convex combination of the unique QSD  $\mu_\epsilon$  and the extinction state, gives a qualitative characterization of the multi-scale dynamics of  $X_t^\epsilon$ . Denote by  $\mathbb{E}_\mu^\epsilon$  the expectation with respect to  $\mathbb{P}_\mu^\epsilon$ , by  $\|\cdot\|_{TV}$  the total variation distance, by  $\mathcal{P}([0, \infty))$  the set of Borel probability measures on  $[0, \infty)$ , and by  $\phi_{\epsilon,1}$  the positive eigenfunction of  $-\mathcal{L}_\epsilon$  associated to  $\lambda_{\epsilon,1}$  satisfying the normalization  $\|\phi_{\epsilon,1}\|_{L^2(u_\epsilon^G)} = 1$ . We point out that  $\phi_{\epsilon,1}$  also belongs to  $L^1(u_\epsilon^G)$  (see Proposition 2.3). Recall that  $\mathcal{A}$  is the global attractor of (1.2), and the uniqueness of  $r_1$ -valleys is equivalent to  $r_1 > r_2$ .

**Theorem C.** *Assume (H) and there exists a unique  $r_1$ -valley.*

(1) *For each compact  $K \subset [0, \infty)$ , there are  $C = C(K) > 0$  and small  $\epsilon_* = \epsilon_*(K) > 0$  such that*

$$\begin{aligned} & \sup_{\substack{\mu \in \mathcal{P}([0, \infty)) \\ \text{supp}(\mu) \subset K}} \left\| \mathbb{P}_\mu^\epsilon[X_t^\epsilon \in \bullet] - [\alpha_\epsilon(\mu)e^{-\lambda_{\epsilon,1}t}\mu_\epsilon + (1 - \alpha_\epsilon(\mu)e^{-\lambda_{\epsilon,1}t})\delta_0] \right\|_{TV} \\ & \leq \exp \left\{ \frac{C}{\epsilon^2} - \lambda_{\epsilon,2}t \right\}, \quad \forall t > 0 \text{ and } \epsilon \in (0, \epsilon_*), \end{aligned}$$

where  $\alpha_\epsilon(\mu) := \|\phi_{\epsilon,1}\|_{L^1(u_\epsilon^G)}\|\phi_{\epsilon,1}\|_{L^1(\mu)}$  satisfies

$$\sup_{\substack{\mu \in \mathcal{P}([0, \infty)) \\ \text{supp}(\mu) \subset K}} \alpha_\epsilon(\mu) \leq 1 + e^{-\frac{r_1 - r_2}{2\epsilon^2}}, \quad \forall \epsilon \in (0, \epsilon_*).$$

(2) *For each  $0 < \delta \ll 1$  and compact sets  $K_1, K_2 \subset (0, \infty)$ , there are  $T = T(K_1, \delta) > 0$ ,  $C_1 = C_1(K_2) > 0$ ,  $C_2 = C_2(K_1, \delta) > 0$ ,  $C_3 = C_3(K_1, \delta) > 0$ ,  $C_4 = C_4(K_1, \delta) > 0$  and  $\epsilon_* = \epsilon_*(K_1) > 0$  such that for each continuous function  $f : (0, \infty) \rightarrow \mathbb{R}$  with  $\text{supp}(f) \subset K_2$ , there holds*

$$\begin{aligned} & \sup_{\substack{\mu \in \mathcal{P}([0, \infty)) \\ \text{supp}(\mu) \subset K_1}} \left| \mathbb{E}_\mu^\epsilon[f(X_t^\epsilon)] - \hat{\alpha}_\epsilon(\mu, \delta)e^{-\lambda_{\epsilon,1}t} \int_0^\infty f d\mu_\epsilon \right| \\ & \leq \frac{C_1}{\epsilon} \exp \left\{ \frac{\sup_{\mathcal{A}_\delta} V - \inf_{K_2} V}{2\epsilon^2} - \lambda_{\epsilon,2}(t - T) \right\} \|f\|_\infty \\ & \quad + C_2 \exp \left\{ -\frac{C_3}{\epsilon^2} - \frac{\lambda_{\epsilon,1}(t - T)}{C_4} \right\} \|f\|_\infty, \quad \forall t > T + 2 \text{ and } \epsilon \in (0, \epsilon_*), \end{aligned}$$

where  $\mathcal{A}_\delta$  is the  $\delta$ -neighborhood of  $\mathcal{A}$  and  $\hat{\alpha}_\epsilon(\mu, \delta) := \|\phi_{\epsilon,1}\|_{L^1(u_\epsilon^G)} \int_{\mathcal{A}_\delta} \phi_{\epsilon,1} d\mu_T^\epsilon e^{\lambda_{\epsilon,1}T}$  with  $\mu_T^\epsilon := \mathbb{P}_\mu^\epsilon[X_T^\epsilon \in \bullet]$  satisfies

$$\sup_{\substack{\mu \in \mathcal{P}([0, \infty)) \\ \text{supp}(\mu) \subset K_1}} \sup_{0 < \delta \ll 1} \hat{\alpha}_\epsilon(\mu, \delta) \leq 1 + e^{-\frac{r_1 - r_2}{2\epsilon^2}}, \quad \forall \epsilon \in (0, \epsilon_*).$$

If, in addition,  $K_1$  is contained in the basin of attraction of a local attractor  $\mathcal{A}^{loc}$  of (1.2), then the same estimate holds with  $\mathcal{A}_\delta$  replaced by  $\mathcal{A}_\delta^{loc}$ , the  $\delta$ -neighborhood of  $\mathcal{A}^{loc}$ .

Results in Theorem C can be interpreted as follows.

- If  $t$  is such that  $t \gg \frac{1}{\lambda_{\epsilon,1}}$ , then  $\|\mathbb{P}_\mu^\epsilon[X_t^\epsilon \in \bullet] - \delta_0\|_{TV} \ll 1$ .
- If  $t$  is such that  $\frac{1}{\epsilon^2 \lambda_{\epsilon,2}} \ll t \ll \frac{1}{\lambda_{\epsilon,1}}$ , then

$$\|\mathbb{P}_\mu^\epsilon[X_t^\epsilon \in \bullet] - [\alpha_\epsilon(\mu)\mu_\epsilon + (1 - \alpha_\epsilon(\mu))\delta_0]\|_{TV} \ll 1.$$



More precise interpretation requires more detailed knowledge of the term  $\alpha_\epsilon(\mu)$ , which is expected to be very close to 1, and therefore,  $\|\mathbb{P}_\mu^\epsilon[X_t^\epsilon \in \bullet] - \mu_\epsilon\|_{TV} \ll 1$ . Since  $\mu_\epsilon$  tends to concentrate on the global attractor  $\mathcal{A}$  and approaches to the set of invariant measures of (1.2) as  $\epsilon \rightarrow 0$  (see [53, Theorem A and B]), transient dynamics of  $X_t^\epsilon$  has to have a close relation to the structure of  $\mathcal{A}$ , that we explore in future works.

Showing  $\alpha_\epsilon(\mu) \approx 1$  amounts to proving

$$\|\phi_{\epsilon,1}\|_{L^1(u_\epsilon^G)} \phi_{\epsilon,1} \rightarrow 1 \quad \text{locally uniformly in } (0, \infty) \quad \text{as } \epsilon \rightarrow 0, \quad (1.6)$$

which turns out to be far-reaching. In Appendix A, we justify (1.6) as well as the limits of  $\alpha_\epsilon(\mu)$  and  $\hat{\alpha}_\epsilon(\mu, \delta)$  in the case that  $b$  changes its sign only once appealing to results established in the forthcoming work [49].

- The estimate in (2) for the dynamics of compactly supported observables can be interpreted in the similar way, can be made much more concise if the constants are allowed to depend on all of  $K_1$ ,  $K_2$  and  $\delta$ , and has an advantage over that in (1) if  $\sup_{\mathcal{A}_\delta} V < \inf_{K_2} V$ , or  $\sup_{\mathcal{A}_\delta^{loc}} V < \inf_{K_2} V$ . In which case, the estimate in (2) begins to tell useful dynamical information of  $X_t^\epsilon$  when  $t \gg 1$  (independent of  $0 < \epsilon \ll 1$ ). In contrast, the estimate in (1) does not tell anything until  $t \gg \frac{1}{\epsilon^2 \lambda_{\epsilon,2}}$ .

We point out that when the global attractor  $\mathcal{A}$  is a singleton set  $\{x^*\}$ , the potential function  $V$  has a unique local minimal point at  $x^*$ , and hence, the condition  $\sup_{\mathcal{A}_\delta} V < \inf_{K_2} V$  is true as long as  $\mathcal{A}_\delta \cap K_2 = \emptyset$ . More generally, if the local attractor  $\mathcal{A}^{loc}$  is contained in  $E := \{x \in (0, \infty) : V(x) = \min V\}$ , then for each compact  $K_2 \subset (0, \infty)$  satisfying  $K_2 \cap E = \emptyset$ , there is  $\delta_* = \delta_*(K_2) > 0$  such that  $\sup_{\mathcal{A}_\delta^{loc}} V < \inf_{K_2} V$  for all  $\delta \in (0, \delta_*)$ .

Estimates as in Theorem C (1) have been established in [11, 12] for absorbed birth-and-death processes whose mean-field ODEs, restricted to the interior of the phase space, have simple dynamics in the sense that the global attractor is a non-degenerate fixed point. If we further explore certain uniform-in- $\epsilon$  ‘‘coming down from infinity’’ of  $X_t^\epsilon$  as did in [11, 12], it is possible to release Theorem C (1) from the dependence on the compact set  $K$ . Since no additional dynamical information can be obtained from doing so, we do not pursue here.

In Theorem C (2),  $\text{supp}(\mu)$  being contained in the compact set  $K_1$  plays an important role. In fact, in the proof, we first use the compact dissipativity property of the dynamical system generated by (1.2) to find  $T$ , which is the smallest time such that  $\bigcup_{x_0 \in K_1} \bigcup_{t > T} x_t \subset \mathcal{A}_{\frac{\delta}{2}}$  (resp.  $\mathcal{A}_{\frac{\delta}{2}}^{loc}$ ), where  $x_t$  is the solution of (1.2), and then apply the sample path large deviation principle to conclude that the distribution of  $X_T^\epsilon$  mainly concentrates in  $\mathcal{A}_\delta$  (resp.  $\mathcal{A}_\delta^{loc}$ ). From where, better estimates can be derived.

The rest of the paper is organized as follows. In Section 2, we collect some preliminary results for later use. In Section 3, we prove Theorem A in the case  $i = 1$ . In Section 4, we prove Theorem A in the case  $i \in \mathbb{N} \setminus \{1\}$ . Section 5 and Section 6 are respectively devoted to the proof of Theorem B and Theorem C. In Section 7, we discuss some applications. We justify (1.6) in Appendix A.

## 2. Preliminaries

This is a service section. We collect preliminary results for later purposes.

**2.1. The potential function.** Basic properties of the potential function  $V$  defined in (1.4) are summarized in the next lemma.



**Lemma 2.1.** *The following hold.*

- $V \in C^2((0, \infty)) \cap C((0, \infty))$ ,  $V(0+) := \lim_{x \rightarrow 0+} V(x) = 0$  and  $V(\infty) := \lim_{x \rightarrow \infty} V(x) = \infty$ .
- $V(x) < 0$  for all  $0 < x \ll 1$ . In particular, the minimum of  $V$  is negative and attainable.

If, in addition, that zeros of  $b$  in  $(0, \infty)$  are isolated, and there is a zero  $x^* \in (0, \infty)$  of  $b$  such that  $b \geq 0$  on  $(0, x^*)$ ,  $b \leq 0$  on  $(x^*, \infty)$  and  $b'(x^*) < 0$ , then  $V$  has a unique local minimal point at  $x^*$ , and satisfies  $V''(x^*) > 0$ .

**2.2. Semi-classical Schrödinger operators.** Consider the change of variable

$$y = \xi(x) = \int_0^x \frac{1}{\sqrt{a(z)}} dz, \quad x \in (0, \infty).$$

Assumptions on  $a$  ensure that  $\xi' > 0$  on  $(0, \infty)$ ,  $\lim_{x \rightarrow 0+} \xi(x) = 0$  and  $y_\infty := \lim_{x \rightarrow \infty} \xi(x) \in (0, \infty]$ . In particular,  $\xi : (0, \infty) \rightarrow (0, y_\infty)$  is invertible.

Let  $v_\epsilon^G(y) := \frac{u_\epsilon^G(x)}{\xi'(x)}$  and define

$$\mathcal{L}_\epsilon^Y = \frac{\epsilon^2}{2} \frac{d^2}{dy^2} - q_\epsilon(y) \frac{d}{dy} : L^2(v_\epsilon^G) \rightarrow L^2(v_\epsilon^G),$$

where  $q_\epsilon = -(\mathcal{L}_\epsilon \xi) \circ \xi^{-1}$  and  $L^2(v_\epsilon^G) := L^2((0, y_\infty), v_\epsilon^G(y) dy)$ . It is straightforward to check that  $U_\epsilon \mathcal{L}_\epsilon = \mathcal{L}_\epsilon^Y U_\epsilon$ , where  $U_\epsilon : L^2(u_\epsilon^G) \rightarrow L^2(v_\epsilon^G)$ ,  $f \mapsto f \circ \xi^{-1}$  is a unitary transform.

Consider the following semi-classical Schrödinger operator

$$\mathcal{L}_\epsilon^S = \frac{\epsilon^2}{2} \frac{d^2}{dy^2} - \frac{1}{2} \left[ \frac{q_\epsilon^2(y)}{\epsilon^2} - q'_\epsilon(y) \right] : L^2((0, y_\infty)) \rightarrow L^2((0, y_\infty)). \quad (2.1)$$

It is easy to verify that  $\tilde{U}_\epsilon \mathcal{L}_\epsilon^Y = \mathcal{L}_\epsilon^S \tilde{U}_\epsilon$ , where  $\tilde{U}_\epsilon : L^2(v_\epsilon^G) \rightarrow L^2((0, y_\infty))$ ,  $f \mapsto f \sqrt{v_\epsilon^G}$  is a unitary transform. We include the following commutative diagram for readers' convenience:

$$\begin{array}{ccccc} L^2(u_\epsilon^G) & \xrightarrow{U_\epsilon} & L^2(v_\epsilon^G) & \xrightarrow{\tilde{U}_\epsilon} & L^2((0, y_\infty)) \\ \downarrow \mathcal{L}_\epsilon & & \downarrow \mathcal{L}_\epsilon^Y & & \downarrow \mathcal{L}_\epsilon^S \\ L^2(u_\epsilon^G) & \xrightarrow{U_\epsilon} & L^2(v_\epsilon^G) & \xrightarrow{\tilde{U}_\epsilon} & L^2((0, y_\infty)) \end{array}$$

Denote by  $V_\epsilon$  the potential of the Schrödinger operator  $-\mathcal{L}_\epsilon^S$ , namely,  $V_\epsilon = \frac{1}{2} \left( \frac{q_\epsilon^2}{\epsilon^2} - q'_\epsilon \right)$ .

**Lemma 2.2.** *The following hold.*

- (1)  $2V_\epsilon(y) = \epsilon^2 \left\{ \frac{3}{16} \frac{[a'(x)]^2}{a(x)} - \frac{a''(x)}{4} \right\} + b'(x) - \frac{a'(x)b(x)}{a(x)} + \frac{1}{\epsilon^2} \frac{b^2(x)}{a(x)}$ .
- (2) *There exists  $y_1 \in (0, y_\infty)$  such that*

$$V_\epsilon(y) \gtrsim \frac{\epsilon^2}{\xi^{-1}(y)}, \quad \forall y \in (0, y_1], \quad 0 < \epsilon \ll 1 \quad \text{and} \quad \inf_{0 < \epsilon \ll 1} \inf_{(0, y_1]} V_\epsilon > 0.$$

- (3) *There exists  $y_2 \in (0, y_\infty)$  such that*

$$V_\epsilon(y) \gtrsim \frac{1}{\epsilon^2} \frac{b^2(\xi^{-1}(y))}{a(\xi^{-1}(y))}, \quad \forall y \in [y_2, y_\infty), \quad 0 < \epsilon \ll 1.$$

- (4) *The family  $\{V_\epsilon\}_{0 < \epsilon \ll 1}$  is uniformly lower bounded, that is,  $\inf_{0 < \epsilon \ll 1} \min_{(0, y_\infty)} V_\epsilon > -\infty$ .*

*Proof.* (1) Note that  $q_\epsilon(y) = -\mathcal{L}_\epsilon \xi(x)$  and  $q'_\epsilon(y) = -\frac{(\mathcal{L}_\epsilon \xi)'(x)}{\xi'(x)}$ . Straightforward calculations give

$$\begin{aligned}\mathcal{L}_\epsilon \xi(x) &= -\frac{\epsilon^2}{4} \frac{a'(x)}{\sqrt{a(x)}} + \frac{b(x)}{\sqrt{a(x)}}, \\ \frac{(\mathcal{L}_\epsilon \xi)^2(x)}{\epsilon^2} &= \frac{\epsilon^2}{16} \frac{[a'(x)]^2}{a(x)} - \frac{1}{2} \frac{a'(x)b(x)}{a(x)} + \frac{1}{\epsilon^2} \frac{b^2(x)}{a(x)}, \\ \frac{(\mathcal{L}_\epsilon \xi)'(x)}{\xi'(x)} &= -\frac{\epsilon^2}{4} a''(x) + \frac{\epsilon^2}{8} \frac{[a'(x)]^2}{a(x)} + b'(x) - \frac{1}{2} \frac{a'(x)b(x)}{a(x)}.\end{aligned}$$

The formula for  $V_\epsilon$  follows.

(2) Let  $x_1 \in (0, 1)$  and  $0 < \delta \ll 1$  be such that the following hold for  $x \in (0, x_1]$ :

$$\begin{aligned}0 \leq a(x) \leq (a'(0) + \delta)x, \quad a'(x) \in [a'(0) - \delta, a'(0) + \delta], \quad a''(x) \leq a''(0) + \delta, \\ b(x) \in [(b'(0) - \delta)x, (b'(0) + \delta)x], \quad b'(x) \geq b'(0) - \delta.\end{aligned}$$

Then, for each  $x \in (0, x_1]$  and  $0 < \epsilon \ll 1$ ,

$$\begin{aligned}2V_\epsilon(\xi(x)) &\geq \epsilon^2 \left\{ \frac{3}{16} \frac{(a'(0) - \delta)^2}{(a'(0) + \delta)x} - \frac{a''(0) + \delta}{4} \right\} + b'(0) - \delta \\ &\quad - \frac{(a'(0) + \delta)(b'(0) + \delta)x}{(a'(0) + \delta)x} + \frac{1}{\epsilon^2} \frac{(b'(0) - \delta)^2 x^2}{(a'(0) + \delta)x} \\ &= \epsilon^2 \left( \frac{c_1}{x} - \frac{a''(0) + \delta}{4} \right) - 2\delta + \frac{c_2 x}{\epsilon^2} \geq \frac{\epsilon^2 c_1}{x} - 3\delta + \frac{c_2 x}{\epsilon^2},\end{aligned}\tag{2.2}$$

where  $c_1 = \frac{3}{16} \frac{(a'(0) - \delta)^2}{a'(0) + \delta}$  and  $c_2 = \frac{(b'(0) - \delta)^2}{a'(0) + \delta}$ . The estimate (2.2) gives  $V_\epsilon(\xi(x)) \gtrsim \frac{\epsilon^2}{x}$  for all  $x \in (0, x_1]$  and  $0 < \epsilon \ll 1$ . Moreover,

$$\inf_{0 < \epsilon \ll 1} \inf_{(0, x_1]} V_\epsilon \circ \xi \geq \frac{1}{2} \inf_{0 < \epsilon \ll 1} \min_{x \in (0, x_1]} \left( \frac{\epsilon^2 c_1}{x} - 3\delta + \frac{c_2 x}{\epsilon^2} \right) = \sqrt{c_1 c_2} - \frac{3\delta}{2} > 0.$$

The results follow by setting  $y_1 = \xi(x_1)$ .

(3) It is a simple consequence of **(H)**(3).

(4) Let  $y_1$  and  $y_2$  be as in (2) and (3), respectively. We may assume, without loss of generality, that  $y_1 < y_2$ . Then,

$$\inf_{0 < \epsilon \ll 1} \min_{[y_1, y_2]} V_\epsilon \geq \frac{1}{2} \inf_{0 < \epsilon \ll 1} \min_{[\xi^{-1}(y_1), \xi^{-1}(y_2)]} \left( -\frac{\epsilon^2 a''}{4} + b' - \frac{a'b}{a} \right) > -\infty.$$

This together with (2) and (3) yields the result.  $\square$

**Remark 2.1.** As  $\xi^{-1}(y) \rightarrow 0$  as  $y \rightarrow 0^+$ , Lemma 2.2 (2) implies that for each  $0 < \epsilon \ll 1$ , there holds  $V_\epsilon(y) \rightarrow \infty$  as  $y \rightarrow 0^+$ . But, it is an unpleasant fact that this blowup is not uniform in  $0 < \epsilon \ll 1$ . In fact, for  $x \approx 0$ , there holds

$$\begin{aligned}2V_\epsilon(\xi(x)) &\approx \epsilon^2 \left\{ \frac{3}{16} \frac{(a'(0))^2}{a'(0)x} - \frac{a''(0)}{4} \right\} + b'(0) - \frac{(a'(0) + a''(0)x)b'(0)x}{a'(0)x} + \frac{1}{\epsilon^2} \frac{(b'(0))^2 x^2}{a'(0)x} \\ &= \epsilon^2 \left[ \frac{3}{16} \frac{a'(0)}{x} - \frac{a''(0)}{4} \right] - \frac{a''(0)b'(0)}{a'(0)} x + \frac{(b'(0))^2 x}{a'(0)\epsilon^2},\end{aligned}$$

leading to

$$2V_\epsilon(\xi(\epsilon^2)) \approx \frac{3a'(0)}{16} - \frac{\epsilon^2 a''(0)}{4} - \frac{\epsilon^2 a''(0)b'(0)}{a'(0)} + \frac{(b'(0))^2}{a'(0)} \approx \frac{3a'(0)}{16} + \frac{(b'(0))^2}{a'(0)}.$$

**2.3. Spectrum and semigroup.** We present the following result on the spectrum of  $-\mathcal{L}_\epsilon$  and the semigroup generated by  $\mathcal{L}_\epsilon$ .

**Proposition 2.1.** *For each  $0 < \epsilon \ll 1$ , the following hold.*

(1)  $-\mathcal{L}_\epsilon$  has purely discrete spectrum contained in  $(0, \infty)$  and listed as follows:

$$\lambda_{\epsilon,1} < \lambda_{\epsilon,2} < \lambda_{\epsilon,3} < \dots \rightarrow \infty.$$

(2) Each  $\lambda_{\epsilon,i}$  is associated with a unique eigenfunction  $\phi_{\epsilon,i} \in L^2(u_\epsilon^G) \cap L^1(u_\epsilon^G) \cap C^2((0, \infty))$  subject to the normalization  $\|\phi_{\epsilon,i}\|_{L^2(u_\epsilon^G)} = 1$ . Moreover,  $\phi_{\epsilon,1}$  is positive on  $(0, \infty)$ .

(3) The set  $\{\phi_{\epsilon,i}, i \in \mathbb{N}\}$  is an orthonormal basis of  $L^2(u_\epsilon^G)$ .

(4)  $\mathcal{L}_\epsilon$  generates a positive analytic semigroup  $(P_t^\epsilon)_{t \geq 0}$  of contractions on  $L^2(u_\epsilon^G)$  satisfying

$$P_t^\epsilon f = \mathbb{E}_\bullet^\epsilon[f(X_t^\epsilon) \mathbb{1}_{t < T_0^\epsilon}], \quad \forall f \in L^2(u_\epsilon^G) \cap C_b([0, \infty)), \quad t \geq 0.$$

(5) For each  $k \in \mathbb{N}$ ,

$$P_t^\epsilon f = \sum_{i=1}^{k-1} e^{-\lambda_{\epsilon,i} t} \langle f, \phi_{\epsilon,i} \rangle_{L^2(u_\epsilon^G)} \phi_{\epsilon,i} + P_t^\epsilon Q_k^\epsilon f, \quad \forall t \geq 0 \text{ and } f \in L^2(u_\epsilon^G), \quad (2.3)$$

where  $Q_k^\epsilon$  is the spectral projection of  $\mathcal{L}_\epsilon$  corresponding to the eigenvalues  $\{\lambda_{\epsilon,j}\}_{j \geq k}$ . Moreover,

$$\|P_t^\epsilon Q_k^\epsilon f\|_{L^2(u_\epsilon^G)} \leq e^{-\lambda_{\epsilon,k} t} \|f\|_{L^2(u_\epsilon^G)}, \quad \forall f \in L^2(u_\epsilon^G).$$

(6) For each  $f \in C_b([0, \infty))$ , (2.3) holds pointwise.

*Proof.* Fix  $0 < \epsilon \ll 1$ . By Lemma 2.2,  $V_\epsilon$  is lower bounded and satisfies  $V_\epsilon(y) \rightarrow \infty$  as  $y \rightarrow 0^+$  and  $y \rightarrow y_\infty^-$ . As a result (see e.g. [2]), the spectrum of  $-\mathcal{L}_\epsilon^S$  is purely discrete and consists of simple eigenvalues. The associated normalized eigenfunctions  $\{\Phi_{\epsilon,i}\}_{i \in \mathbb{N}}$  are twice continuously differentiable and form an orthonormal basis of  $L^2((0, y_\infty))$ . The eigenfunction  $\Phi_{\epsilon,1}$  associated to  $\lambda_{\epsilon,1}$  can be chosen to be positive on  $(0, y_\infty)$ . Moreover,  $\{\Phi_{\epsilon,i}\}_{i \in \mathbb{N}} \subset L^2((0, y_\infty), (V_\epsilon + M)dy)$  for some  $M > 0$  satisfying  $\inf V_\epsilon + M > 0$ .

As  $\mathcal{L}_\epsilon$  and  $\mathcal{L}_\epsilon^S$  are unitarily equivalent, all conclusions in (1)-(3) except  $\lambda_{\epsilon,1} > 0$  and  $\phi_{\epsilon,i} \in L^1(u_\epsilon^G)$  follow, where  $\phi_{\epsilon,i} := U_\epsilon^{-1} \tilde{U}_\epsilon^{-1} \Phi_{\epsilon,i}$ . Obviously,  $\lambda_{\epsilon,1} \geq 0$ . Suppose  $\lambda_{\epsilon,1} = 0$  for contradiction. Then,  $-\mathcal{L}_\epsilon \phi_{\epsilon,1} = 0$ . It follows from (1.3) that  $\frac{\epsilon^2}{2} \int_0^\infty a |\phi'_{\epsilon,1}|^2 u_\epsilon^G dx = 0$ , which implies that  $\phi_{\epsilon,1}$  is a positive constant function on  $(0, \infty)$ . As  $u_\epsilon^G$  is non-integrable on  $(0, \infty)$ , we find  $\int_0^\infty \phi_{\epsilon,1}^2 u_\epsilon^G dx = \infty$ , which is contradictory to  $\phi_{\epsilon,1} \in L^2(u_\epsilon^G)$ . Hence,  $\lambda_{\epsilon,1} > 0$ . The decay of  $u_\epsilon^G$  near  $\infty$  ensures  $\int_1^\infty \phi_{\epsilon,i} u_\epsilon^G dx < \infty$ , and the combination of  $\Phi_{\epsilon,i} \in L^2((0, y_\infty), (V_\epsilon + M)dy)$  and the behavior of  $V_\epsilon$  near 0 guarantee  $\int_0^1 \phi_{\epsilon,i} u_\epsilon^G dx < \infty$  (see [5, Proposition 4.3]). Thus,  $\phi_{\epsilon,i} \in L^1(u_\epsilon^G)$ .

Due to the self-adjointness of  $\mathcal{L}_\epsilon$ ,

$$\|(\lambda - \mathcal{L}_\epsilon)^{-1}\|_{L^2(u_\epsilon^G) \rightarrow L^2(u_\epsilon^G)} = \frac{1}{\text{dist}(\lambda, \sigma(\mathcal{L}_\epsilon))} \leq \frac{1}{|\lambda|}, \quad \forall \lambda \in \mathbb{C} \text{ with } \Re \lambda > 0.$$

Therefore (see e.g. [50, Theorem 2.5.2]),  $\mathcal{L}_\epsilon$  generates an analytic semigroup  $(P_t^\epsilon)_{t \geq 0}$  of contractions on  $L^2(u_\epsilon^G)$ . The positivity of  $(P_t^\epsilon)_{t \geq 0}$  follows from the maximum principle. The theory of symmetric Dirichlet forms (see e.g. [5]) gives the following stochastic representation of  $P_t^\epsilon$ :

$$P_t^\epsilon f(x) = \mathbb{E}_x^\epsilon[f(X_t^\epsilon) \mathbb{1}_{t < T_0^\epsilon}], \quad \forall x \in (0, \infty) \text{ and } f \in C_b([0, \infty)) \cap L^2(u_\epsilon^G).$$

This proves (4). The statements in (5) follow from general theory concerning the asymptotic behaviors of analytic semigroups (see e.g. [18, Theorem V.3.1]) and the simplicity of eigenvalues  $\lambda_{\epsilon,i}$ ,  $i \in \mathbb{N}$ .

Since  $\{\phi_{\epsilon,i}\}_{i \in \mathbb{N}} \subset L^1(u_\epsilon^G)$  and  $Q_k^\epsilon = id - \sum_{i=1}^{k-1} \langle \bullet, \phi_{\epsilon,i} \rangle_{L^2(u_\epsilon^G)} \phi_{\epsilon,i}$ , each term on the right-hand side of (2.3) is well-defined. Standard approximations then ensure that (2.3) holds in the pointwise sense for each  $f \in C_b([0, \infty))$ . This proves (6), and finishes the proof of the proposition.  $\square$

As a result of Proposition 2.1, the following variational formulas for  $\lambda_{\epsilon,i}$ ,  $i \in \mathbb{N}$  hold:

$$\lambda_{\epsilon,1} = \inf_{\phi \in C_0^\infty((0,\infty)) \setminus \{0\}} \frac{\langle -\mathcal{L}_\epsilon \phi, \phi \rangle_{L^2(u_\epsilon^G)}}{\|\phi\|_{L^2(u_\epsilon^G)}^2} = \inf_{\phi \in C_0^\infty((0,\infty)) \setminus \{0\}} \frac{\frac{\epsilon^2}{2} \int_0^\infty a |\phi'|^2 u_\epsilon^G dx}{\int_0^\infty \phi^2 u_\epsilon^G dx}, \quad (2.4)$$

$$\lambda_{\epsilon,i} = \inf_{\substack{\phi \in C_0^\infty((0,\infty)) \setminus \{0\} \\ \phi \perp \phi_{\epsilon,j}, j \in \{1, \dots, i-1\}}} \frac{\langle -\mathcal{L}_\epsilon \phi, \phi \rangle_{L^2(u_\epsilon^G)}}{\|\phi\|_{L^2(u_\epsilon^G)}^2}, \quad i \in \mathbb{N} \setminus \{1\}, \quad (2.5)$$

where the orthogonality  $\perp$  is taken in  $L^2(u_\epsilon^G)$ .

**2.4. Uniform estimate of the first eigenfunctions.** We prove the following result asserting that the normalized eigenfunction  $\phi_{\epsilon,1}$  given in Proposition 2.1, as an element of  $L^2(u_\epsilon^G)$ , has uniform-in- $\epsilon$  positive concentration away from 0 and  $\infty$ .

**Proposition 2.2.** *There are  $\kappa \in (0, 1)$  and  $K \gg 1$  such that  $\int_{(0,\infty) \setminus (\frac{1}{K}, K)} \phi_{\epsilon,1}^2 u_\epsilon^G dx \leq \kappa$  for all  $0 < \epsilon \ll 1$ .*

Set  $\Phi_{\epsilon,1} := \tilde{U}_\epsilon U_\epsilon \phi_{\epsilon,1}$ , where  $U_\epsilon$  and  $\tilde{U}_\epsilon$  are the unitary transforms defined in Subsection 2.2. The next result is a simple consequence of the definitions and Proposition 2.1.

**Lemma 2.3.** *The following hold:*

- (1)  $\Phi_{\epsilon,1} \in L^2((0, y_\infty)) \cap C^2((0, y_\infty))$ ,  $\|\Phi_{\epsilon,1}\|_{L^2((0, y_\infty))} = 1$ ;
- (2)  $-\mathcal{L}_\epsilon^S \Phi_{\epsilon,1} = \lambda_{\epsilon,1} \Phi_{\epsilon,1}$ ;
- (3) for each  $I \subset (0, \infty)$ , there holds  $\int_I \phi_{\epsilon,1}^2 u_\epsilon^G dx = \int_{\xi(I)} \Phi_{\epsilon,1}^2 dy$ .

**Lemma 2.4.** *There holds  $\lim_{\epsilon \rightarrow 0} \lambda_{\epsilon,1} = 0$ .*

*Proof.* Clearly,  $E := \{x \in (0, \infty) : V(x) = \min V\}$  is compact in  $(0, \infty)$ . Let  $I$  be an open interval satisfying  $E \subset I \subset \subset (0, \infty)$ . Let  $\phi \in C_0^\infty((0, \infty))$  satisfy  $\phi(x) = x$  for  $x \in I$ .

As  $\min V < 0$  and  $\inf_{(0,\infty) \setminus I} V > \min_I V$ , we find  $\int_I e^{-\frac{V(x)}{\epsilon^2}} dx \rightarrow \infty$  and  $\frac{\int_{(0,\infty) \setminus I} e^{-\frac{V(x)}{\epsilon^2}} dx}{\int_I e^{-\frac{V(x)}{\epsilon^2}} dx} \rightarrow 0$  as  $\epsilon \rightarrow 0$ . It follows from (2.4) that

$$\begin{aligned} \lambda_{\epsilon,1} &\leq \frac{\epsilon^2 \int_I a |\phi'|^2 u_\epsilon^G dx + \int_{(0,\infty) \setminus I} a |\phi'|^2 u_\epsilon^G dx}{\int_I \phi^2 u_\epsilon^G dx} \\ &\leq \frac{\epsilon^2 \int_I e^{-\frac{V(x)}{\epsilon^2}} dx + \left( \sup_{\text{supp}(\phi)} |\phi'|^2 \right) \int_{(0,\infty) \setminus I} e^{-\frac{V(x)}{\epsilon^2}} dx}{\left( \inf_{x \in I} \frac{x^2}{a(x)} \right) \int_I e^{-\frac{V(x)}{\epsilon^2}} dx} \rightarrow 0 \quad \text{as } \epsilon \rightarrow 0. \end{aligned}$$

This proves the lemma.  $\square$

**Proof of Proposition 2.2.** Let  $y_1$  and  $y_2$  be as in Lemma 2.2 (2) and (3), respectively. We may assume, without loss of generality, that  $y_1 < y_2$ . Then,

$$V_{\inf} := \inf_{0 < \epsilon \ll 1} \min V_\epsilon > -\infty \quad \text{and} \quad V_0 := \inf_{0 < \epsilon \ll 1} \inf_{(0, y_1] \cup [y_2, y_\infty)} V_\epsilon > 0.$$

Denote by  $D(\mathcal{L}_\epsilon^S)$  the domain of  $\mathcal{L}_\epsilon^S$ . By (1.3) and the unitary equivalence between  $\mathcal{L}_\epsilon$  and  $\mathcal{L}_\epsilon^S$ ,

$$\frac{\epsilon^2}{2} \int_0^{y_\infty} \phi' \psi' dy + \int_0^{y_\infty} V_\epsilon \phi \psi dy = -\langle \mathcal{L}_\epsilon^S \phi, \psi \rangle_{L^2((0, y_\infty))}, \quad \forall \phi, \psi \in D(\mathcal{L}_\epsilon^S),$$

which together with Lemma 2.3 (1)(2) ensures  $\frac{\epsilon^2}{2} \int_0^{y_\infty} |\Phi'_{\epsilon,1}|^2 dy + \int_0^{y_\infty} V_\epsilon \Phi_{\epsilon,1}^2 dy = \lambda_{\epsilon,1}$ , leading to  $\int_{(0, y_1] \cup [y_2, y_\infty)} V_\epsilon \Phi_{\epsilon,1}^2 dy \leq \lambda_{\epsilon,1} - \int_{y_1}^{y_2} V_\epsilon \Phi_{\epsilon,1}^2 dy$ . Therefore,

$$V_0 \int_{(0, y_1] \cup [y_2, y_\infty)} \Phi_{\epsilon,1}^2 dy \leq \lambda_{\epsilon,1} - \min\{0, V_{\inf}\} \int_{y_1}^{y_2} \Phi_{\epsilon,1}^2 dy.$$

Since  $\int_{(0, y_1] \cup [y_2, y_\infty)} \Phi_{\epsilon,1}^2 dy + \int_{y_1}^{y_2} \Phi_{\epsilon,1}^2 dy = 1$ , we deduce  $\int_{(0, y_1] \cup [y_2, y_\infty)} \Phi_{\epsilon,1}^2 dy \leq \kappa_\epsilon := \frac{\lambda_{\epsilon,1} - \min\{0, V_{\inf}\}}{V_0 - \min\{0, V_{\inf}\}}$ , which together with Lemma 2.3 (3) yields  $\int_{(0, \xi^{-1}(y_1)] \cup [\xi^{-1}(y_2), \infty)} \phi_{\epsilon,1}^2 u_\epsilon^G dx \leq \kappa_\epsilon$ .

Lemma 2.4 guarantees  $\kappa := \sup_{0 < \epsilon \ll 1} \kappa_\epsilon \in (0, 1)$ . Setting  $K := \max\left\{\frac{1}{\xi^{-1}(y_1)}, \xi^{-1}(y_2)\right\}$  gives  $\int_{(0, \infty) \setminus (\frac{1}{K}, K)} \phi_{\epsilon,1}^2 u_\epsilon^G dx \leq \kappa$  for all  $0 < \epsilon \ll 1$ . This completes the proof.  $\square$

**2.5. QSD: existence, uniqueness and convergence.** Recall that  $X_t^\epsilon$  is the stochastic process generated by solutions of (1.1), and  $T_0^\epsilon$  is the first time that  $X_t^\epsilon$  hits 0, namely,  $T_0^\epsilon = \inf\{t \geq 0 : X_t^\epsilon = 0\}$ . The next result is proven in [5]. We refer the reader to [6, 41, 46, 8, 29, 9, 10] and references therein for further developments of QSDs for singular diffusion processes.

**Proposition 2.3** ([5]). *For each  $0 < \epsilon \ll 1$ , the following hold.*

- (1)  $X_t^\epsilon$  admits a unique QSD  $\mu_\epsilon$  with density  $u_\epsilon = \frac{\phi_{\epsilon,1} u_\epsilon^G}{\|\phi_{\epsilon,1}\|_{L^1(u_\epsilon^G)}}$ . The extinction rate associated to  $\mu_\epsilon$  is given by  $\lambda_{\epsilon,1}$ .
- (2) For each Borel probability measure  $\mu$  on  $(0, \infty)$  with compact support,

$$\begin{aligned} & \lim_{t \rightarrow \infty} e^{(\lambda_{\epsilon,2} - \lambda_{\epsilon,1})t} \left( \mathbb{P}_\mu^\epsilon [X_t^\epsilon \in B | t < T_0^\epsilon] - \mu_\epsilon(B) \right) \\ &= \frac{\int_0^\infty \phi_{\epsilon,2} d\mu}{\int_0^\infty \phi_{\epsilon,1} d\mu} \left( \frac{\langle \mathbb{1}_B, \phi_{\epsilon,2} \rangle_{L^2(u_\epsilon^G)}}{\|\phi_{\epsilon,1}\|_{L^1(u_\epsilon^G)}} - \frac{\langle \mathbb{1}_B, \phi_{\epsilon,1} \rangle_{L^2(u_\epsilon^G)} \langle \mathbb{1}, \phi_{\epsilon,2} \rangle_{L^2(u_\epsilon^G)}}{\|\phi_{\epsilon,1}\|_{L^1(u_\epsilon^G)}^2} \right), \quad \forall B \in \mathcal{B}((0, \infty)), \end{aligned}$$

where  $\mathcal{B}((0, \infty))$  is the Borel  $\sigma$ -algebra of  $(0, \infty)$ .

*Proof.* Given Proposition 2.1, it is easy to see that  $\mu_\epsilon$  is a QSD with extinction rate  $\lambda_{\epsilon,1}$ . The uniqueness and convergence follow respectively from [5, Theorem 7.3] and [5, Proposition 5.5].  $\square$

### 3. Exponential asymptotic of the first eigenvalue

In this section, we study the asymptotic of the first eigenvalue  $\lambda_{\epsilon,1}$  as  $\epsilon \rightarrow 0$  and prove Theorem A in the case  $i = 1$ , which is restated in the following theorem.

**Theorem 3.1.** *Assume (H). Then,  $\lim_{\epsilon \rightarrow 0} \epsilon^2 \ln \lambda_{\epsilon,1} = -r_1$ .*

We first establish the upper bound.

**Proof of Theorem 3.1: Upper Bound.** Let  $I \subset (0, \infty)$  be an  $r_1$ -valley. Then, Definition 1.2 gives

$$V(\partial I) - \inf_I V = r_1. \quad (3.1)$$

For  $0 < \delta \ll 1$ , let  $0 < \eta \ll 1$  be such that  $|V(x) - V(y)| \leq \delta$  whenever  $x, y \in \bar{I}$  and  $|x - y| \leq 3\eta$ . Let  $\psi \in C_0^\infty((0, \infty))$  satisfy  $\psi = 1$  on  $[\inf I + 3\eta, \sup I - 3\eta]$  and  $\psi = 0$  on  $(0, \infty) \setminus [\inf I + \eta, \sup I - \eta]$ . According to Laplace's method for integrals or the large deviation principle, we find

$$\limsup_{\epsilon \rightarrow 0} \epsilon^2 \ln \int_0^\infty |\psi'|^2 e^{-\frac{V}{\epsilon^2}} dx \leq \limsup_{\epsilon \rightarrow 0} \epsilon^2 \ln \left[ \max_{J_1} |\psi'|^2 \int_{J_1} e^{-\frac{V}{\epsilon^2}} dx \right] \leq -\min_{J_1} V + \delta \leq -V(\partial I) + 2\delta,$$

where  $J_1 = [\inf I + \eta, \inf I + 3\eta] \cup [\sup I - 3\eta, \sup I - \eta]$ . Moreover,

$$\liminf_{\epsilon \rightarrow 0} \epsilon^2 \ln \int_0^\infty \frac{\psi^2}{a} e^{-\frac{V}{\epsilon^2}} dx \geq \liminf_{\epsilon \rightarrow 0} \epsilon^2 \ln \left[ \frac{1}{\max_{J_2} a} \int_{J_2} e^{-\frac{V}{\epsilon^2}} dx \right] \geq -\min_{J_2} V - \delta,$$

where  $J_2 = [\inf I + 3\eta, \sup I - 3\eta]$ . Without loss of generality, we assume that  $\delta$  is so small that  $\inf_{J_2} V = \inf_I V$ . It then follows from (2.4) and (3.1) that

$$\limsup_{\epsilon \rightarrow 0} \epsilon^2 \ln \lambda_{\epsilon,1} \leq -V(\partial I) + 2\delta + \inf_I V + \delta = -r_1 + 3\delta.$$

Since  $0 < \delta \ll 1$  is arbitrary, we conclude the upper bound  $\limsup_{\epsilon \rightarrow 0} \epsilon^2 \ln \lambda_{\epsilon,1} \leq -r_1$ .  $\square$

We proceed to prove the lower bound. In the next lemma, we construct special minimizing sequences that help to overcome the difficulties caused by the non-integrability of  $u_\epsilon^G$  near 0.

**Lemma 3.1.** *There are sequences  $\{\psi_{\epsilon,n}\}_{n \in \mathbb{N}, 0 < \epsilon \ll 1} \subset C_0^\infty((0, \infty))$  satisfying the following properties:*

- $\|\psi_{\epsilon,n}\|_{L^2(u_\epsilon^G)} = 1$  for each  $n \in \mathbb{N}$  and  $0 < \epsilon \ll 1$ ;
- $\lambda_{\epsilon,1} = \lim_{n \rightarrow \infty} \frac{\epsilon^2}{2} \int_0^\infty a |\psi'_{\epsilon,n}|^2 u_\epsilon^G dx$  for each  $0 < \epsilon \ll 1$ ;
- $\sup_{0 < \epsilon \ll 1} \|\psi_{\epsilon,n} - \phi_{\epsilon,1}\|_{L^2(u_\epsilon^G)} \rightarrow 0$  as  $n \rightarrow \infty$ .

In particular, there are  $\kappa \in (0, 1)$  and  $K \gg 1$  such that  $\int_{(0, \infty) \setminus (\frac{1}{K}, K)} \psi_{\epsilon,n}^2 u_\epsilon^G dx \leq \kappa$  for all  $0 < \epsilon \ll 1$  and  $n \gg 1$ .

*Proof.* For each  $0 < \epsilon \ll 1$ , let  $\{\tilde{\psi}_{\epsilon,n}\}_{n \in \mathbb{N}} \subset C_0^\infty((0, \infty))$  satisfy  $\|\tilde{\psi}_{\epsilon,n}\|_{L^2(u_\epsilon^G)} = 1$  for each  $n \in \mathbb{N}$  and  $\tilde{\psi}_{\epsilon,n} \rightarrow \phi_{\epsilon,1}$  in  $D(\mathcal{E}_\epsilon)$  as  $n \rightarrow \infty$ , where  $D(\mathcal{E}_\epsilon)$  is the domain of  $\mathcal{E}_\epsilon$  given in Section 1. The expected sequences  $\{\psi_{\epsilon,n}\}_{n \in \mathbb{N}, 0 < \epsilon \ll 1}$  can be obtained by extracting subsequences from  $\{\tilde{\psi}_{\epsilon,n}\}_{n \in \mathbb{N}, 0 < \epsilon \ll 1}$ . Indeed, for each  $m \in \mathbb{N}$ , let  $n(\epsilon, m) \gg 1$  be such that  $\|\tilde{\psi}_{\epsilon,n(\epsilon,m)} - \phi_{\epsilon,1}\|_{D(\mathcal{E}_\epsilon)} \leq \frac{1}{m}$ . We then only need to define  $\psi_{\epsilon,m} := \tilde{\psi}_{\epsilon,n(\epsilon,m)}$  for each  $m \in \mathbb{N}$  and  $0 < \epsilon \ll 1$ . The ‘‘In particular’’ part is a simple consequence of Proposition 2.2 and the construction.  $\square$

**Proof of Theorem 3.1: Lower Bound.** As  $V \in C([0, \infty))$ , it can be extended to be a function in  $C(\mathbb{R})$ , still denoted by  $V$  (note we do not need any specific property of  $V$  on  $(-\infty, 0)$ ). Let  $\{\psi_{\epsilon,n}\}_{n \in \mathbb{N}, 0 < \epsilon \ll 1}$ ,  $\kappa$  and  $K$  be as in Lemma 3.1. By zero extension on  $(-\infty, 0]$ , we consider  $\{\psi_{\epsilon,n}\}_{n \in \mathbb{N}, 0 < \epsilon \ll 1}$  as functions on  $\mathbb{R}$ .

Fix  $0 < \delta \ll 1$ . Let  $\eta : \mathbb{R} \rightarrow (0, \infty)$  be continuously differentiable and satisfy  $|V(y) - V(z)| \leq \delta$  whenever  $x, y, z \in \mathbb{R}$  satisfy  $|y - x| \leq \eta(x)$  and  $|z - x| \leq \eta(x)$ .

Let  $\{x_i\}_{i=1}^N \subset (\frac{1}{K}, K)$  be such that  $I_i := (x_i - \eta(x_i), x_i + \eta(x_i)) \subset (\frac{1}{K+1}, K+1)$  for all  $i \in \{1, \dots, N\}$  and  $[\frac{1}{K}, K] \subset \bigcup_{i=1}^N I_i$ . By Lemma 3.1, for  $0 < \epsilon \ll 1$  and  $n \gg 1$ ,

$$1 = \int_{(0, \infty) \setminus (\frac{1}{K}, K)} \frac{\psi_{\epsilon, n}^2}{a} e^{-\frac{V}{\epsilon^2}} dx + \int_{\frac{1}{K}}^K \frac{\psi_{\epsilon, n}^2}{a} e^{-\frac{V}{\epsilon^2}} dx \leq \kappa + \frac{1}{\min_{[\frac{1}{K}, K]} a} \sum_{i=1}^N e^{-\frac{V(x_i) + \delta}{\epsilon^2}} \int_{I_i} \psi_{\epsilon, n}^2 dx. \quad (3.2)$$

Fix  $i \in \{1, \dots, N\}$ . Let  $\xi_i \in C_{x_i}$  be  $C^1$ . For  $x \in I_i$ , we define

$$\ell_{i, x}(t) = \xi_i(t) + \frac{(1-t)(x - x_i) + t \frac{\eta(0)}{2\eta(x_i)}(x - x_i - \eta(x_i))}{(1-t)\eta(x_i) + t\eta(0)} \eta(\xi_i(t)), \quad t \in [0, 1],$$

It is straightforward to check that the family  $\{\ell_{i, x}\}_{x \in I_i}$  satisfies the following properties:

- $\ell_{i, x}$  is  $C^1$  for each  $x \in I_i$ , and  $\sup_{x \in I_i} \sup_{t \in [0, 1]} |\ell'_{i, x}(t)|$  is finite;
- $\ell_{i, x}(0) = x$  and  $\ell_{i, x}(1) = \frac{\eta(0)}{2\eta(x_i)}(x - x_i - \eta(x_i)) \in (-\eta(0), 0)$  for all  $x \in I_i$ ;
- $|\ell_{i, x}(t) - \xi_i(t)| \leq \eta(\xi_i(t))$  for all  $t \in [0, 1]$  and  $x \in I_i$ ; thus,  $|V(\ell_{i, x}(t)) - V(\xi_i(t))| \leq \delta$  for all  $t \in [0, 1]$  and  $x \in I_i$ ;
- for each  $t \in [0, 1]$ , there holds  $\frac{d}{dx} \ell_{i, x}(t) = \frac{1-t+t \frac{\eta(0)}{2\eta(x_i)}}{(1-t)\eta(x_i) + t\eta(0)} \eta(\xi_i(t))$  for  $x \in I_i$ .

As  $\ell_{i, x}(0) = x$  and  $\psi_{\epsilon, n}(\ell_{i, x}(1)) = 0$ , we integrate  $\frac{d}{dt} \psi_{\epsilon, n}(\ell_{i, x}(t)) = \psi'_{\epsilon, n}(\ell_{i, x}(t)) \ell'_{i, x}(t)$  over  $[0, 1]$  to find  $-\psi_{\epsilon, n}(x) = \int_0^1 \psi'_{\epsilon, n}(\ell_{i, x}(t)) \ell'_{i, x}(t) dt$  for  $x \in I_i$ . It follows that

$$\begin{aligned} \int_{I_i} \psi_{\epsilon, n}^2 dx &= \int_{I_i} \left[ \int_0^1 \psi'_{\epsilon, n}(\ell_{i, x}(t)) \ell'_{i, x}(t) dt \right]^2 dx \\ &\leq \sup_{x \in I_i} \max_{t \in [0, 1]} |\ell'_{i, x}(t)|^2 \int_0^1 \int_{I_i} |\psi'_{\epsilon, n}(\ell_{i, x}(t))|^2 dx dt \\ &\leq \sup_{x \in I_i} \max_{t \in [0, 1]} |\ell'_{i, x}(t)|^2 \int_0^1 e^{\frac{V(\xi_i(t)) + \delta}{\epsilon^2}} \int_{I_i} |\psi'_{\epsilon, n}(\ell_{i, x}(t))|^2 e^{-\frac{V(\ell_{i, x}(t))}{\epsilon^2}} dx dt \\ &\leq \sup_{x \in I_i} \max_{t \in [0, 1]} |\ell'_{i, x}(t)|^2 \exp \left\{ \frac{\max_{t \in [0, 1]} V(\xi_i(t)) + \delta}{\epsilon^2} \right\} \int_0^1 \int_{I_i} |\psi'_{\epsilon, n}(\ell_{i, x}(t))|^2 e^{-\frac{V(\ell_{i, x}(t))}{\epsilon^2}} dx dt. \end{aligned}$$

Note that for each  $t \in [0, 1]$ , the function  $x \mapsto \frac{d}{dx} \ell_{i, x}(t)$  is a constant function on  $I_i$ , leading to

$$\begin{aligned} \int_{I_i} |\psi'_{\epsilon, n}(\ell_{i, x}(t))|^2 e^{-\frac{V(\ell_{i, x}(t))}{\epsilon^2}} dx &= \frac{1}{\frac{d}{dx} \ell_{i, x}(t)} \int_{I_i} |\psi'_{\epsilon, n}(\ell_{i, x}(t))|^2 e^{-\frac{V(\ell_{i, x}(t))}{\epsilon^2}} \frac{d}{dx} \ell_{i, x}(t) dx \\ &\leq \frac{1}{\frac{d}{dx} \ell_{i, x}(t)} \int_0^\infty |\psi'_{\epsilon, n}|^2 e^{-\frac{V}{\epsilon^2}} dx. \end{aligned}$$

Setting  $D_i := \sup_{x \in I_i} \max_{t \in [0, 1]} |\ell'_{i, x}(t)|^2 \int_0^1 \frac{1}{\frac{d}{dx} \ell_{i, x}(t)} dt$ , we arrive at

$$\int_{I_i} \psi_{\epsilon, n}^2 dx \leq D_i \exp \left\{ \frac{\max_{t \in [0, 1]} V(\xi_i(t)) + \delta}{\epsilon^2} \right\} \int_0^\infty |\psi'_{\epsilon, n}|^2 e^{-\frac{V}{\epsilon^2}} dx,$$

which together with (3.2) yields

$$1 - \kappa \leq \frac{\sum_{i=1}^N D_i \exp \left\{ \frac{\max_{t \in [0, 1]} V(\xi_i(t)) - V(x_i) + 2\delta}{\epsilon^2} \right\}}{\min_{[\frac{1}{K}, K]} a} \int_0^\infty |\psi'_{\epsilon, n}|^2 e^{-\frac{V}{\epsilon^2}} dx.$$



Since  $\xi_i \in C_{x_i}$  being  $C^1$  is arbitrary and the set of such curves are dense in  $C_{x_i}$ , we find

$$\begin{aligned} 1 - \kappa &\leq \frac{\sum_{i=1}^N D_i \exp \left\{ \frac{\inf_{\xi \in C_{x_i}} \max_{t \in [0,1]} V(\xi(t)) - V(x_i) + 2\delta}{\epsilon^2} \right\}}{\min_{[\frac{1}{K}, K]} a} \int_0^\infty |\psi'_{\epsilon,n}|^2 e^{-\frac{V}{\epsilon^2}} dx \\ &\leq \frac{\sum_{i=1}^N D_i}{\min_{[\frac{1}{K}, K]} a} \exp \left\{ \frac{\sup_{x \in (0,\infty)} [\inf_{\xi \in C_x} \max_{t \in [0,1]} V(\xi(t)) - V(x)] + 2\delta}{\epsilon^2} \right\} \int_0^\infty |\psi'_{\epsilon,n}|^2 e^{-\frac{V}{\epsilon^2}} dx \\ &= \frac{\sum_{i=1}^N D_i}{\min_{[\frac{1}{K}, K]} a} e^{\frac{r_1 + 2\delta}{\epsilon^2}} \int_0^\infty |\psi'_{\epsilon,n}|^2 e^{-\frac{V}{\epsilon^2}} dx. \end{aligned}$$

Thus,

$$\int_0^\infty |\psi'_{\epsilon,n}|^2 e^{-\frac{V}{\epsilon^2}} dx \geq \frac{\min_{[\frac{1}{K}, K]} a}{\sum_{i=1}^N D_i} e^{-\frac{r_1 + 2\delta}{\epsilon^2}} (1 - \kappa), \quad \forall 0 < \epsilon \ll 1 \quad \text{and} \quad n \gg 1.$$

Note that the term  $\frac{\min_{[\frac{1}{K}, K]} a}{\sum_{i=1}^N D_i}$  is independent of the functions  $\{\psi_{\epsilon,n}\}_{\epsilon,n}$ . Letting  $n \rightarrow \infty$  yields

$$\lambda_{\epsilon,1} \geq \frac{\epsilon^2 \min_{[\frac{1}{K}, K]} a}{2 \sum_{i=1}^N D_i} e^{-\frac{r_1 + 2\delta}{\epsilon^2}} (1 - \kappa), \quad \forall 0 < \epsilon \ll 1,$$

which gives  $\liminf_{\epsilon \rightarrow 0} \epsilon^2 \ln \lambda_{\epsilon,1} \geq -r_1 - 2\delta$ . Since  $0 < \delta \ll 1$  is arbitrary, we arrive at the lower bound  $\liminf_{\epsilon \rightarrow 0} \epsilon^2 \ln \lambda_{\epsilon,1} \geq -r_1$ .  $\square$

We make a remark about the proof of the lower bound in Theorem 3.1.

**Remark 3.1.** *While the arguments used in the proof of the lower bound in Theorem 3.1 have the potential to be generalized to treat reversible diffusions in higher dimensions, the proof of the lower bound can be achieved in a much more concise way as  $r_1$  has a much simpler representation:*

$$r_1 := \sup_{x \in (0,\infty)} \left[ \sup_{y \in (0,x)} V(y) - V(x) \right].$$

Let  $\kappa \in (0,1)$  and  $K > 0$  be as in Proposition 2.2. Then,  $\int_0^K \phi_{\epsilon,1}^2 u_\epsilon^G dx \geq 1 - \kappa$  for all  $0 < \epsilon \ll 1$ . Noting that  $\phi_{\epsilon,1}(0+) = 0$ , which is a simple result of Sobolev embedding theorem and the integrability of  $\phi_{\epsilon,1}^2$  with respect to  $u_\epsilon^G$  near 0, we deduce

$$\begin{aligned} 1 - \kappa &\leq \int_0^K \phi_{\epsilon,1}^2 u_\epsilon^G dx = \int_0^K \frac{1}{a} e^{-\frac{V}{\epsilon^2}} \left[ \int_0^x \phi'_{\epsilon,1}(y) dy \right]^2 dx \\ &\leq \int_0^K \frac{x}{a} e^{-\frac{V(x)}{\epsilon^2}} \int_0^x |\phi'_{\epsilon,n}(y)|^2 dy dx \\ &\leq \int_0^K \frac{x}{a} e^{-\frac{V(x)}{\epsilon^2}} e^{\frac{1}{\epsilon^2} \sup_{y \in (0,x)} V(y)} \int_0^x |\phi'_{\epsilon,1}(y)|^2 e^{-\frac{V(y)}{\epsilon^2}} dy dx \\ &\leq \int_0^K \frac{x}{a} e^{\frac{1}{\epsilon^2} [\sup_{y \in (0,x)} V(y) - V(x)]} dx \times \int_0^\infty |\phi'_{\epsilon,1}|^2 e^{-\frac{V}{\epsilon^2}} dy \\ &\leq \left[ \sup_{x \in (0,K)} \frac{x}{a(x)} \right] e^{\frac{r_1}{\epsilon^2}} \lambda_{\epsilon,1}, \quad \forall 0 < \epsilon \ll 1, \end{aligned}$$

where we used the fact  $\lambda_{\epsilon,1} = \int_0^\infty |\phi'_{\epsilon,1}|^2 e^{-\frac{V}{\epsilon^2}} dy$  in the last step. From which, the lower bound  $\liminf_{\epsilon \rightarrow 0} \epsilon^2 \ln \lambda_{\epsilon,1} \geq -r_1$  follows readily.

#### 4. Exponential asymptotics of other eigenvalues

In this section, we prove Theorem A in the case  $i \in \mathbb{N} \setminus \{1\}$ , which is restated as follows.

**Theorem 4.1.** *Assume (H). Then,  $\lim_{\epsilon \rightarrow 0} \epsilon^2 \ln \lambda_{\epsilon, i} = -r_i$  for each  $i \in \mathbb{N} \setminus \{1\}$ .*

The upper and lower bounds are respectively established in Subsection 4.1 and Subsection 4.3. In Subsection 4.2, we state results on asymptotics of the first eigenvalues of some auxiliary operators.

**4.1. Upper bound.** For an open interval  $I \subset \subset (0, \infty)$ , let  $\mathcal{L}_\epsilon^I$  be the generator of the diffusion process  $X_t^\epsilon$  or (1.1) restricted to  $I$  and killed on  $\partial I$ , which is non-degenerate, regular and reversible. Moreover,  $-\mathcal{L}_\epsilon^I$ , considered as an operator in  $L^2(I, u_\epsilon^G dx)$ , has purely discrete spectrum consisting of simple eigenvalues. Let  $\lambda_\epsilon^I > 0$  be the first eigenvalue of  $-\mathcal{L}_\epsilon^I$ . Then,

$$\lambda_\epsilon^I = \inf_{\phi \in C_0^\infty(I)} \frac{\frac{\epsilon^2}{2} \int_I a |\phi'|^2 u_\epsilon^G dx}{\int_I \phi^2 u_\epsilon^G dx} = \inf_{\phi \in C_0^\infty(I)} \frac{\frac{\epsilon^2}{2} \int_I |\phi'|^2 e^{-\frac{V}{\epsilon^2}} dx}{\int_I \frac{\phi^2}{a} e^{-\frac{V}{\epsilon^2}} dx}. \quad (4.1)$$

For  $x \in I$  and  $y \in \partial I$ , let

$$C_{x,y}^I := \{\xi : [0, 1] \rightarrow \bar{I} : \xi \text{ is continuous and satisfies } \xi(0) = x \text{ and } \xi(1) = y\}.$$

Define

$$r^I := \sup_{x \in I} \left[ \inf_{y \in \partial I} \inf_{\xi \in C_{x,y}^I} \sup_{t \in [0, 1]} V(\xi(t)) - V(x) \right].$$

**Proposition 4.1** ([43]). *There holds  $\lim_{\epsilon \rightarrow 0} \epsilon^2 \ln \lambda_\epsilon^I = -r^I$ .*

Recall from Section 1 that for  $r > 0$ ,  $N(r)$  is the number of  $r$ -valleys.

**Proof of Theorem 4.1: Upper Bound.** Fix  $i_* \geq 2$ . We show

$$\limsup_{\epsilon \rightarrow 0} \epsilon^2 \ln \lambda_{\epsilon, i_*} \leq -r_{i_*}. \quad (4.2)$$

We distinguish the following two cases:

**Case 1.** There is  $r > 0$  such that  $N(r) \geq i_*$ .

**Case 2.**  $N(r) < i_*$  for all  $r > 0$ .

**Case 1.** In this case,  $\sup\{r > 0 : N(r) \geq i_*\}$  is well-defined, and hence, by Remark 1.1 (1),

$$r_{i_*} = \sup\{r > 0 : N(r) \geq i_*\}.$$

Let  $0 < \delta_* \ll 1$ . For  $r \in (r_{i_*} - \delta_*, r_{i_*})$ , let  $I_k$ ,  $k = 1, \dots, N(r)$  be the  $r$ -valleys. They are ordered in the nature way:  $\sup I_k \leq \inf I_{k+1}$  for all  $k \in \{1, \dots, N(r) - 1\}$ .

We claim that

$$\limsup_{\epsilon \rightarrow 0} \epsilon^2 \ln \lambda_{\epsilon, N(r)} \leq -r, \quad \forall r \in (r_{i_*} - \delta_*, r_{i_*}). \quad (4.3)$$

Note that  $N(r) \geq i_*$  for all  $r \in (r_{i_*} - \delta_*, r_{i_*})$ , implying  $\lambda_{\epsilon, N(r)} \geq \lambda_{\epsilon, i_*}$  for all  $r \in (r_{i_*} - \delta_*, r_{i_*})$ . Thus, if the claim (4.3) is true, then  $\limsup_{\epsilon \rightarrow 0} \epsilon^2 \ln \lambda_{\epsilon, i_*} \leq -r$  for all  $r \in (r_{i_*} - \delta_*, r_{i_*})$ . Letting  $r \rightarrow r_{i_*}^-$  yields (4.2).

It suffices to show (4.3). Making  $\delta_*$  smaller if necessary, we claim that

$$I_k \subset \subset (0, \infty), \quad \forall k \in \{1, \dots, N(r)\}, \quad r \in (r_{i_*} - \delta_*, r_{i_*}). \quad (4.4)$$

Indeed, the only possible exception takes place if  $\inf I_1 = 0$  for some  $r \in (r_{i_*} - \delta_*, r_{i_*})$ . If this is the case, then  $I_1 = (0, \beta)$ , where  $\beta = \min\{x \in (0, \infty) : V(x) = 0\}$ . Thus,  $\gamma(I_1)$  depends only on  $V$ . Note that there always holds  $\gamma(I_1) \notin (r_{i_*} - \delta_*, r_{i_*})$  if  $\delta_*$  is sufficiently small, which contradicts the fact  $\gamma(I_1) = r \in (r_{i_*} - \delta_*, r_{i_*})$ .

Fix  $0 < \delta \ll 1$ . For  $k \in \{1, \dots, N(r)\}$  and  $0 < \epsilon \ll 1$ , let  $\psi_{\epsilon, k} \in C_0^\infty(I_k)$  be such that

$$\int_{I_k} \psi_{\epsilon, k}^2 u_\epsilon^G dx = 1 \quad \text{and} \quad \frac{\frac{\epsilon^2}{2} \int_{I_k} |\psi'_{\epsilon, k}|^2 e^{-\frac{V}{\epsilon^2}} dx}{\int_{I_k} \frac{\psi_{\epsilon, k}^2}{a} e^{-\frac{V}{\epsilon^2}} dx} \leq \lambda_\epsilon^{I_k} e^{\frac{\delta}{\epsilon^2}},$$

where  $\lambda_\epsilon^{I_k}$  is given in (4.1). Moreover, let  $\alpha_{\epsilon, k}$ ,  $k \in \{1, \dots, N(r)\}$  and  $0 < \epsilon \ll 1$  be such that

$$\sum_{k=1}^{N(r)} \alpha_{\epsilon, k}^2 = 1 \quad \text{and} \quad \sum_{k=1}^{N(r)} \alpha_{\epsilon, k} \int_{I_k} \psi_{\epsilon, k} \phi_{\epsilon, \ell} u_\epsilon^G dx = 0, \quad \forall \ell \in \{1, \dots, N(r) - 1\},$$

where we recall from Proposition 2.1 that  $\phi_{\epsilon, \ell}$  is the normalized eigenfunction of  $\mathcal{L}_\epsilon$  associated to the eigenvalue  $\lambda_{\epsilon, \ell}$ .

Set  $\psi_\epsilon = \sum_{k=1}^{N(r)} \alpha_{\epsilon, k} \psi_{\epsilon, k}$ . Since  $I_k$ ,  $k \in \{1, \dots, N(r)\}$  are pairwise disjoint, we find:

$$\int_0^\infty \psi_\epsilon^2 u_\epsilon^G dx = \sum_{k=1}^{N(r)} \alpha_{\epsilon, k}^2 \int_{I_k} \psi_{\epsilon, k}^2 u_\epsilon^G dx = \sum_{k=1}^{N(r)} \alpha_{\epsilon, k}^2 = 1.$$

Moreover,

$$\int_0^\infty \psi_\epsilon \phi_{\epsilon, \ell} u_\epsilon^G dx = \sum_{k=1}^{N(r)} \alpha_{\epsilon, k} \int_{I_k} \psi_{\epsilon, k} \phi_{\epsilon, \ell} u_\epsilon^G dx = 0, \quad \forall \ell \in \{1, \dots, N(r) - 1\}.$$

In addition, the pairwise disjointness of  $I_k$ ,  $k \in \{1, \dots, N(r)\}$  implies that

$$\begin{aligned} \frac{\epsilon^2}{2} \int_0^\infty |\psi'_\epsilon|^2 e^{-\frac{V}{\epsilon^2}} dx &= \frac{\epsilon^2}{2} \sum_{k=1}^{N(r)} \alpha_{\epsilon, k}^2 \int_{I_k} |\psi'_{\epsilon, k}|^2 e^{-\frac{V}{\epsilon^2}} dx \\ &\leq \frac{\epsilon^2}{2} e^{\frac{\delta}{\epsilon^2}} \sum_{k=1}^{N(r)} \alpha_{\epsilon, k}^2 \lambda_\epsilon^{I_k} \int_{I_k} \frac{\psi_{\epsilon, k}^2}{a} e^{-\frac{V}{\epsilon^2}} dx \\ &\leq \frac{\epsilon^2}{2} e^{\frac{\delta}{\epsilon^2}} \left( \max_{k \in \{1, \dots, N(r)\}} \lambda_\epsilon^{I_k} \right) \sum_{k=1}^{N(r)} \alpha_{\epsilon, k}^2 \int_{I_k} \frac{\psi_{\epsilon, k}^2}{a} e^{-\frac{V}{\epsilon^2}} dx \\ &= \frac{\epsilon^2}{2} e^{\frac{\delta}{\epsilon^2}} \left( \max_{k \in \{1, \dots, N(r)\}} \lambda_\epsilon^{I_k} \right). \end{aligned}$$

The variational formula for  $\lambda_{\epsilon, N(r)}$  given in (2.5) then yields  $\lambda_{\epsilon, N(r)} \leq \frac{\epsilon^2}{2} e^{\frac{\delta}{\epsilon^2}} (\max_{k \in \{1, \dots, N(r)\}} \lambda_\epsilon^{I_k})$ . As  $\ln(\max_{k \in \{1, \dots, N(r)\}} \lambda_\epsilon^{I_k}) = \max_{k \in \{1, \dots, N(r)\}} \ln \lambda_\epsilon^{I_k}$ , we deduce from Proposition 4.1 that

$$\limsup_{\epsilon \rightarrow 0} \epsilon^2 \ln \lambda_{\epsilon, N(r)} \leq \delta - \min_{k \in \{1, \dots, N(r)\}} r^{I_k}.$$

Since  $0 < \delta \ll 1$  is arbitrary, we conclude  $\limsup_{\epsilon \rightarrow 0} \epsilon^2 \ln \lambda_{\epsilon, N(r)} \leq -\min_{k \in \{1, \dots, N(r)\}} r^{I_k}$ . As each  $I_k$  is a  $r$ -valley, the definition of  $r^{I_k}$  ensures that  $r^{I_k} = r$ . This proves (4.3), and thus, finishes the proof in this case.

**Case 2.** In this case,  $r_{i_*} = 0$  (see Remark 1.1 (1)) so that (4.2) reads

$$\limsup_{\epsilon \rightarrow 0} \epsilon^2 \ln \lambda_{\epsilon, i_*} \leq 0. \quad (4.5)$$

Let  $I_k \subset\subset (0, \infty)$ ,  $k \in \{1, \dots, i_*\}$  be pairwise disjoint, and on at least one of them,  $V$  is monotone. As a result, at least one of  $r^{I_k}$ ,  $k \in \{1, \dots, i_*\}$  is zero, that is,  $\min_{k \in \{1, \dots, i_*\}} r^{I_k} = 0$ .

Fix  $0 < \delta \ll 1$ . For  $k \in \{1, \dots, i_*\}$  and  $0 < \epsilon \ll 1$ , let  $\psi_{\epsilon, k} \in C_0^\infty(I_k)$  be such that

$$\int_{I_k} \psi_{\epsilon, k}^2 u_\epsilon^G dx = 1 \quad \text{and} \quad \frac{\frac{\epsilon^2}{2} \int_{I_k} |\psi'_{\epsilon, k}|^2 e^{-\frac{V}{\epsilon^2}} dx}{\int_{I_k} \frac{\psi_{\epsilon, k}^2}{a} e^{-\frac{V}{\epsilon^2}} dx} \leq \lambda_\epsilon^{I_k} e^{\frac{\delta}{\epsilon^2}}.$$

Moreover, let  $\alpha_{\epsilon, k}$ ,  $k \in \{1, \dots, i_*\}$  and  $0 < \epsilon \ll 1$  be such that

$$\sum_{k=1}^{i_*} \alpha_{\epsilon, k}^2 = 1 \quad \text{and} \quad \sum_{k=1}^{i_*} \alpha_{\epsilon, k} \int_{I_k} \psi_{\epsilon, k} \phi_{\epsilon, \ell} u_\epsilon^G dx = 0. \quad \forall \ell \in \{1, \dots, i_* - 1\}.$$

Set  $\psi_\epsilon = \sum_{k=1}^{i_*} \alpha_{\epsilon, k} \psi_{\epsilon, k}$ . Arguing as in **Case 1**, we find

$$\int_0^\infty \psi_\epsilon^2 u_\epsilon^G dx = 1, \quad \int_0^\infty \psi_\epsilon \phi_{\epsilon, \ell} u_\epsilon^G dx = 0, \quad \forall \ell \in \{1, \dots, i_* - 1\},$$

$$\frac{\epsilon^2}{2} \int_0^\infty |\psi'_\epsilon|^2 e^{-\frac{V}{\epsilon^2}} dx \leq \frac{\epsilon^2}{2} e^{\frac{\delta}{\epsilon^2}} \left( \max_{k \in \{1, \dots, i_*\}} \lambda_\epsilon^{I_k} \right).$$

Hence,  $\lambda_{\epsilon, i_*} \leq \frac{\epsilon^2}{2} e^{\frac{\delta}{\epsilon^2}} (\max_{k \in \{1, \dots, i_*\}} \lambda_\epsilon^{I_k})$ , implying  $\limsup_{\epsilon \rightarrow 0} \epsilon^2 \ln \lambda_{\epsilon, i_*} \leq \delta - \min_{k \in \{1, \dots, i_*\}} r^{I_k} = \delta$ . As  $0 < \delta \ll 1$  is arbitrary, the upper bound (4.5) follows.  $\square$

**4.2. Auxiliary eigenvalue asymptotics.** In this subsection, we consider the first eigenvalue asymptotics of three types of operators who are generators of the diffusion process  $X_t^\epsilon$  or (1.1) restricted to different types of intervals with reflection on boundaries. More precisely, let  $I \subset (0, \infty)$  be a proper open interval. There are only three types:

Type 1 :  $I = (0, b)$  for  $0 < b < \infty$ ,

Type 2 :  $I = (a, b)$  for  $0 < a < b < \infty$ ,

Type 3 :  $I = (a, \infty)$  for  $0 < a < \infty$ .

Let  $\mathcal{L}_\epsilon^{I, R}$  be the generator of the diffusion process  $X_t^\epsilon$  or (1.1) restricted to  $I$  and reflected on the right boundary (Type 1), both boundaries (Type 2) and the left boundary (Type 3). The superscript “ $R$ ” stands for reflection. This operator extends to a self-adjoint operator in  $L^2(I, u_\epsilon^G dx)$ , still denoted by  $\mathcal{L}_\epsilon^{I, R}$ . The rigorous formulation can be done using Dirichlet forms as done in Section 1 for the operator  $\mathcal{L}_\epsilon$ . Proposition 2.1 has a version for  $-\mathcal{L}_\epsilon^{I, R}$ .

Note that for a Type 2 or Type 3 interval  $I$ ,  $u_\epsilon^G$  is integrable on  $I$ , and hence,  $u_\epsilon^{Gibbs} := \frac{u_\epsilon^G}{\int_I u_\epsilon^G dx}$  is the Gibbs density, namely, the density of the unique stationary distribution of the restricted diffusion process. In particular, the first eigenvalue of  $-\mathcal{L}_\epsilon^{I, R}$  is zero.

Let

$$\lambda_\epsilon^{I, R} = \begin{cases} \text{the first eigenvalue of } -\mathcal{L}_\epsilon^{I, R}, & \text{if } I \text{ is of Type 1,} \\ \text{the first non-zero eigenvalue of } -\mathcal{L}_\epsilon^{I, R}, & \text{if } I \text{ is of Type 2 or Type 3.} \end{cases}$$

Recall that

- if  $I$  is of Type 1, then

$$\lambda_\epsilon^{I,R} = \inf_{\substack{\phi \in C^\infty(I) \setminus \{0\} \\ \inf \text{supp}(\phi) > 0}} \frac{\frac{\epsilon^2}{2} \int_I a |\phi'|^2 u_\epsilon^G dx}{\int_I \phi^2 u_\epsilon^G dx}, \quad (4.6)$$

- if  $I$  is of Type 2, then

$$\lambda_\epsilon^{I,R} = \inf_{\phi \in C^\infty(I) \setminus \{0\}} \frac{\frac{\epsilon^2}{2} \int_I a |\phi'|^2 u_\epsilon^G dx}{\int_I \left( \phi - \int_I \phi u_\epsilon^{\text{Gibbs}} dx \right)^2 u_\epsilon^G dx}, \quad (4.7)$$

- if  $I$  is of Type 3, then

$$\lambda_\epsilon^{I,R} = \inf_{\substack{\phi \in C^\infty(I) \setminus \{0\} \\ \sup \text{supp}(\phi) < \infty}} \frac{\frac{\epsilon^2}{2} \int_I a |\phi'|^2 u_\epsilon^G dx}{\int_I \left( \phi - \int_I \phi u_\epsilon^{\text{Gibbs}} dx \right)^2 u_\epsilon^G dx}. \quad (4.8)$$

We define the following quantity:

$$r^{I,R} = \begin{cases} \sup_{x \in I} \left[ \inf_{\xi \in C_x^{I,R}} \sup_{t \in [0,1]} V(\xi(t)) - V(x) \right], & \text{if } I \text{ is of Type 1,} \\ \sup_{x,y \in I} \left[ \inf_{\xi \in C_{x,y}^{I,R}} \sup_{t \in [0,1]} V(\xi(t)) - V(x) - V(y) \right] + \inf_I V, & \text{if } I \text{ is of Type 2 or Type 3,} \end{cases} \quad (4.9)$$

where for  $x, y \in I$ ,

$$\begin{aligned} C_x^{I,R} &= \{ \xi : [0, 1] \rightarrow \bar{I} : \xi \text{ is continuous and satisfies } \xi(0) = x \text{ and } \xi(1) = 0 \}, \\ C_{x,y}^{I,R} &= \{ \xi : [0, 1] \rightarrow \bar{I} : \xi \text{ is continuous and satisfies } \xi(0) = x \text{ and } \xi(1) = y \}. \end{aligned}$$

**Proposition 4.2.** *There holds  $\lim_{\epsilon \rightarrow 0} \epsilon^2 \ln \lambda_\epsilon^{I,R} = -r^{I,R}$ .*

*Proof.* If  $I$  is of Type 2, the result is proven in [30, 42]. As  $V(\infty) = \infty$ , the role played by a reflection boundary at a finite right endpoint is similar to that of  $\infty$ . In consideration of this, when  $I$  is of Type 1, the proof is similar to that of Theorem A in the case  $i = 1$ . If  $I$  is of Type 3, the proof is similar to that when  $I$  is of Type 2.  $\square$

**4.3. Lower bound.** We prove the lower bound in this subsection.

**Proof of Theorem 4.1: Lower Bound.** Fix  $i_* \geq 2$ . We prove

$$\liminf_{\epsilon \rightarrow 0} \epsilon^2 \ln \lambda_{\epsilon, i_*} \geq -r_{i_*}. \quad (4.10)$$

Let  $0 < \delta_* \ll 1$  be fixed. For  $r \in (r_{i_*}, r_{i_*} + \delta_*)$ , let  $I_k(r)$ ,  $k \in \{1, \dots, N(r)\}$  be the  $r$ -valleys. They are ordered in the natural way:  $\sup I_k(r) \leq \inf I_{k+1}(r)$  for all  $k \in \{1, \dots, N(r) - 1\}$ .

We claim that

$$\liminf_{\epsilon \rightarrow 0} \epsilon^2 \ln \lambda_{\epsilon, N(r)+1} \geq -r, \quad \forall r \in (r_{i_*}, r_{i_*} + \delta_*). \quad (4.11)$$

Note that making  $\delta_*$  smaller if necessary, there exists  $n_0 \in \mathbb{N}$  such that

$$N(r) + n_0 = i_*, \quad \forall r \in (r_{i_*}, r_{i_*} + \delta_*). \quad (4.12)$$

Thus, if the claim (4.11) is true, then

$$\limsup_{\epsilon \rightarrow 0} \epsilon^2 \ln \lambda_{\epsilon, i_*} = \limsup_{\epsilon \rightarrow 0} \epsilon^2 \ln \lambda_{\epsilon, N(r)+n_0} \geq \liminf_{\epsilon \rightarrow 0} \epsilon^2 \ln \lambda_{\epsilon, N(r)+1} \geq -r, \quad \forall r \in (r_{i_*}, r_{i_*} + \delta_*).$$

Letting  $r \rightarrow r_{i_*}^+$  yields (4.10). Hence, it suffices to prove (4.11), which is done in three steps.

**Step 1.** Note that making  $\delta_*$  smaller if necessary, there hold for each  $r \in (r_{i_*}, r_{i_*} + \delta_*)$ ,

$$\begin{aligned} I_k(r) &\subset\subset (0, \infty), \quad \forall k \in \{1, \dots, N(r)\}, \\ \sup I_k(r) &< \inf I_{k+1}(r), \quad \forall k = 1, \dots, N(r) - 1. \end{aligned}$$

The first conclusion follows from arguments leading to (4.4). The second one follows from (4.12) saying that the number of  $r$ -valleys remains constant for  $r \in (r_{i_*}, r_{i_*} + \delta_*)$ . Otherwise, two or more of them merge as  $r$  increases so that the number of  $r$ -valleys drops.

The purpose of this step is to find for each  $r \in (r_{i_*}, r_{i_*} + \delta_*)$  some special points in the following disjoint intervals:

$$(0, \inf I_1(r)), \quad (\sup I_k(r), \inf I_{k+1}(r)), \quad \forall k \in \{1, \dots, N(r) - 1\}$$

in order to re-partition  $(0, \infty)$ .

We first treat the intervals  $(\sup I_k(r), \inf I_{k+1}(r))$  for  $k \in \{1, \dots, N(r) - 1\}$  and  $r \in (r_{i_*}, r_{i_*} + \delta_*)$ . Note that

$$\sup_{(\sup I_k(r), \inf I_{k+1}(r))} V > \max \{V(\sup I_k(r)), V(\inf I_{k+1}(r))\}, \quad \forall k \in \{1, \dots, N(r) - 1\}, \quad r \in (r_{i_*}, r_{i_*} + \delta_*). \quad (4.13)$$

In fact, if (4.13) fails for some  $r_0 \in (r_{i_*}, r_{i_*} + \delta_*)$  and  $k_0 \in \{1, \dots, N(r_0) - 1\}$ , then for all  $r \in (r_0, r_{i_*} + \delta_*)$ ,  $I_{k_0}(r)$  and  $I_{k_0+1}(r)$  merge, and therefore,  $N(r) < N(r_0)$  for all  $r \in (r_0, r_{i_*} + \delta_*)$ , which contradicts (4.12).

According to (4.13), for each  $r \in (r_{i_*}, r_{i_*} + \delta_*)$  and  $k \in \{1, \dots, N(r) - 1\}$  there exists  $x_k(r) \in (\sup I_k(r), \inf I_{k+1}(r))$  such that

$$V(x_k(r)) = \sup_{(\sup I_k(r), \inf I_{k+1}(r))} V.$$

Now, we treat the intervals  $(0, \inf I_1(r))$  for  $r \in (r_{i_*}, r_{i_*} + \delta_*)$ . Note that

$$\sup_{(0, \inf I_1(r))} V > V(\inf I_1(r)), \quad \forall r \in (r_{i_*}, r_{i_*} + \delta_*). \quad (4.14)$$

Indeed, if (4.14) fails for some  $r_0 \in (r_{i_*}, r_{i_*} + \delta_*)$ , then  $I_1(r)$  is no longer a  $r$ -valley for all  $r \in (r_0, r_{i_*} + \delta_*)$ , leading to a contradiction.

For  $r \in (r_{i_*}, r_{i_*} + \delta_*)$ , set  $E(r) = \left\{x \in (0, \inf I_1(r)) : V(x) = \sup_{(0, \inf I_1(r))} V\right\}$ . We claim that there holds either

$$E(r) \neq \emptyset, \quad \forall r \in (r_{i_*}, r_{i_*} + \delta_*), \quad (4.15)$$

or

$$E(r) = \emptyset, \quad \forall r \in (r_{i_*}, r_{i_*} + \delta_*). \quad (4.16)$$

Suppose on the contrary that there are  $r_1, r_2 \in (r_{i_*}, r_{i_*} + \delta_*)$  such that  $E(r_1) \neq \emptyset$  and  $E(r_2) = \emptyset$ . If  $r_1 > r_2$ , then  $I_1(r_1) \supset I_1(r_2)$ . It follows that  $E(r_2) = E(r_1)$ , leading to a contradiction. If  $r_1 < r_2$ , then  $I_1(r_1) \subset I_1(r_2)$ . By (4.14) and the assumptions, there holds

$$V(\inf I_1(r_2)) < V(0+) \leq \sup_{(0, \inf I_1(r_1))} V,$$

which implies  $\inf I_1(r_2) > \max E(r_1)$ . It follows that  $E(r_2) = E(r_1)$ , leading to a contradiction. Hence, either (4.15) or (4.16) holds.

If (4.15) is true, we fix any

$$x_0(r) \in E(r), \quad \forall r \in (r_{i_*}, r_{i_*} + \delta_*).$$

If (4.16) is the case, we see from (4.14) that  $V(0+) > V(x)$  for all  $x \in (0, \inf I_1(r))$  and  $r \in (r_{i_*}, r_{i_*} + \delta_*)$ . For each  $r \in (r_{i_*}, r_{i_*} + \delta_*)$ , let

$$x_0(r) := \begin{cases} \min \{x \in (0, \inf I_1(r)) : V(0+) - V(x) = r\}, & \text{if } V(0+) > \inf_{(0, \inf I_1(r))} V + r, \\ \inf I_1(r), & \text{if } V(0+) \leq \inf_{(0, \inf I_1(r))} V + r. \end{cases}$$

Hence, we have defined the points:  $x_k(r)$ ,  $k \in \{0, 1, \dots, N(r) - 1\}$ ,  $r \in (r_{i_*}, r_{i_*} + \delta_*)$ .

**Step 2.** We re-partition  $(0, \infty)$  by defining the following intervals: for each  $r \in (r_{i_*}, r_{i_*} + \delta_*)$ ,

$$\begin{aligned} \tilde{I}_0(r) &= (0, x_0(r)), \\ \tilde{I}_k(r) &= (x_{k-1}(r), x_k(r)), \quad k \in \{1, \dots, N(r) - 1\}, \\ \tilde{I}_{N(r)}(r) &= (x_{N(r)-1}(r), \infty). \end{aligned}$$

Note that  $\tilde{I}_0(r)$  is of Type 1,  $\tilde{I}_k(r)$  is of Type 2 for each  $k \in \{1, \dots, N(r) - 1\}$ , and  $\tilde{I}_{N(r)}(r)$  is of Type 3. Recall from (4.9) the definition of  $r^{\tilde{I}_k(r), R}$ . We claim that

$$r^{\tilde{I}_k(r), R} \leq r, \quad \forall k \in \{0, 1, \dots, N(r)\}, \quad r \in (r_{i_*}, r_{i_*} + \delta_*). \quad (4.17)$$

The rest of this step is devoted to the proof of (4.17). Note that  $I_k(r)$  is the only  $r$ -valley contained in  $\tilde{I}_k(r)$ . Otherwise, the number of  $r$ -valleys changes as  $r$  varies.

By the definition of  $x_0(r)$ , it is not hard to see that  $V(0+) - \inf_{(0, x_0(r))} V \leq r$ . This is trivial if (4.16) is the case. If (4.15) is the case, there must hold  $V(0+) - \inf_{(0, x_0(r))} V < r$  as there exists no  $r$ -valley contained in  $(0, x_0(r))$ . The claim (4.17) in the case  $k = 0$  follows.

Let  $r \in (r_{i_*}, r_{i_*} + \delta_*)$  and  $k \in \{1, \dots, N(r)\}$ . Note that

$$r^{\tilde{I}_k(r), R} = \sup_{\substack{x, y \in \tilde{I}_k(r) \\ x < y}} \left[ \inf_{\xi \in C_{x, y}^{\tilde{I}_k(r), R}} \sup_{t \in [0, 1]} V(\xi(t)) - V(x) - V(y) \right] + \inf_{\tilde{I}_k(r)} V.$$

Let  $E_k(r)$  be the set of local maximal points of  $V|_{\tilde{I}_k(r)}$  with connected components attached to  $\partial \tilde{I}_k(r)$  (if there is any) removed. If  $E_k(r) = \emptyset$ , then it is easy to check that  $r_1^{\tilde{I}_k(r), R} = 0$ . Suppose  $E_k(r) \neq \emptyset$ . Then, for each  $x, y \in \tilde{I}_k(r)$  with  $x < y$ , there holds

$$\inf_{\xi \in C_{x, y}^{\tilde{I}_k(r), R}} \sup_{t \in [0, 1]} V(\xi(t)) = \begin{cases} \max \{V(x), V(y)\}, & \text{if } (x, y) \cap E_k(r) = \emptyset, \\ \max_{\{x, y\} \cup ((x, y) \cap E_k(r))} V, & \text{if } (x, y) \cap E_k(r) \neq \emptyset, \end{cases}$$

leading to

$$\begin{aligned} & \inf_{\xi \in C_{x, y}^{\tilde{I}_k(r), R}} \sup_{t \in [0, 1]} V(\xi(t)) - V(x) - V(y) \\ &= \begin{cases} -\min \{V(x), V(y)\}, & \text{if } (x, y) \cap E_k(r) = \emptyset, \\ \max_{\{x, y\} \cup ((x, y) \cap E_k(r))} V - V(x) - V(y), & \text{if } (x, y) \cap E_k(r) \neq \emptyset. \end{cases} \end{aligned}$$



Thus,

$$\sup_{\substack{x,y \in \tilde{I}_k(r) \\ (x,y) \cap E_k(r) = \emptyset}} \left[ \inf_{\xi \in C_{x,y}^{\tilde{I}_k(r),R}} \sup_{t \in [0,1]} V(\xi(t)) - V(x) - V(y) \right] = - \inf_{\tilde{I}_k(r)} V.$$

We claim that

$$\tilde{r}_{\tilde{I}_k(r),R} := \sup_{\substack{x,y \in \tilde{I}_k(r) \\ (x,y) \cap E_k(r) \neq \emptyset}} \left[ \inf_{\xi \in C_{x,y}^{\tilde{I}_k(r),R}} \sup_{t \in [0,1]} V(\xi(t)) - V(x) - V(y) \right] \leq r - \inf_{\tilde{I}_k(r)} V. \quad (4.18)$$

Let  $[\alpha_1, \beta_1], \dots, [\alpha_{\tilde{N}}, \beta_{\tilde{N}}]$  for some  $\tilde{N} = \tilde{N}(k, r) \in \mathbb{N}$  be the connected components of  $E_k(r)$ . They are ordered in the natural way:  $\beta_\ell < \alpha_{\ell+1}$  for all  $\ell \in \{1, \dots, \tilde{N} - 1\}$ . Set  $\beta_0 := \inf \tilde{I}_k(r)$  and  $\alpha_{\tilde{N}+1} := \sup \tilde{I}_k(r)$ . Clearly,  $\beta_0 < \alpha_1$  and  $\beta_{\tilde{N}} < \alpha_{\tilde{N}+1}$ .

Note that if  $x_0 \in [\alpha_{\ell_0}, \beta_{\ell_0}]$  for some  $\ell_0 \in \{1, \dots, \tilde{N}\}$  and  $y \in (x_0, \sup \tilde{I}_k(r))$ , then

$$\inf_{\xi \in C_{x_0,y}^{\tilde{I}_k(r),R}} \sup_{t \in [0,1]} V(\xi(t)) - V(x_0) - V(y) \leq \inf_{\xi \in C_{x,y}^{\tilde{I}_k(r),R}} \sup_{t \in [0,1]} V(\xi(t)) - V(x) - V(y)$$

for all  $x < \alpha_{\ell_0}$  with  $\alpha_{\ell_0} - x \ll 1$ . Similarly, if  $y_0 \in [\alpha_{\ell_0}, \beta_{\ell_0}]$  for some  $\ell_0 \in \{1, \dots, \tilde{N}\}$  and  $x \in (\inf \tilde{I}_k(r), y_0)$ , then

$$\inf_{\xi \in C_{x,y_0}^{\tilde{I}_k(r),R}} \sup_{t \in [0,1]} V(\xi(t)) - V(x) - V(y_0) \leq \inf_{\xi \in C_{x,y}^{\tilde{I}_k(r),R}} \sup_{t \in [0,1]} V(\xi(t)) - V(x) - V(y)$$

for all  $y > \beta_{\ell_0}$  with  $y - \beta_{\ell_0} \ll 1$ . Therefore,

$$\begin{aligned} \tilde{r}_{\tilde{I}_k(r),R} &= \sup_{\substack{\ell_1, \ell_2 \in \{1, \dots, \tilde{N}\} \\ \ell_1 \leq \ell_2}} \sup_{\substack{x \in (\beta_{\ell_1-1}, \alpha_{\ell_1}) \\ y \in (\beta_{\ell_2}, \alpha_{\ell_2+1})}} \left[ \inf_{\xi \in C_{x,y}^{\tilde{I}_k(r),R}} \sup_{t \in [0,1]} V(\xi(t)) - V(x) - V(y) \right] \\ &= \sup_{\substack{\ell_1, \ell_2 \in \{1, \dots, \tilde{N}\} \\ \ell_1 \leq \ell_2}} \sup_{\substack{x \in (\beta_{\ell_1-1}, \alpha_{\ell_1}) \\ y \in (\beta_{\ell_2}, \alpha_{\ell_2+1})}} \left[ \max_{\{x,y\} \cup (\cup_{\ell=\ell_1}^{\ell_2} [\alpha_\ell, \beta_\ell])} V - V(x) - V(y) \right] \\ &= \sup_{\substack{\ell_1, \ell_2 \in \{1, \dots, \tilde{N}\} \\ \ell_1 \leq \ell_2}} \left[ \max_{\cup_{\ell=\ell_1}^{\ell_2} [\alpha_\ell, \beta_\ell]} V - \min_{(\beta_{\ell_1-1}, \alpha_{\ell_1})} V - \min_{(\beta_{\ell_2}, \alpha_{\ell_2+1})} V \right], \end{aligned} \quad (4.19)$$

where the minimals are attained due to the definition of  $E_k(r)$ .

We show that for each  $\ell_1, \ell_2 \in \{1, \dots, \tilde{N}\}$  with  $\ell_1 \leq \ell_2$  there holds

$$\min \left\{ \max_{\cup_{\ell=\ell_1}^{\ell_2} [\alpha_\ell, \beta_\ell]} V - \min_{(\beta_{\ell_1-1}, \alpha_{\ell_1})} V, \max_{\cup_{\ell=\ell_1}^{\ell_2} [\alpha_\ell, \beta_\ell]} V - \min_{(\beta_{\ell_2}, \alpha_{\ell_2+1})} V \right\} \leq r. \quad (4.20)$$

To see this, we recall that  $I_k(r)$  is the only  $r$ -valley contained in  $\tilde{I}_k(r)$ , and the following hold:

$$\begin{aligned} V(\inf \tilde{I}_k(r)) &= \sup_{(\inf \tilde{I}_k(r), \sup I_k(r))} V, & V(\sup \tilde{I}_k(r)) &= \sup_{(\inf I_k(r), \sup \tilde{I}_k(r))} V, \\ \max \left\{ V(\inf \tilde{I}_k(r)), V(\sup \tilde{I}_k(r)) \right\} &= \sup_{\tilde{I}_k(r)} V > \sup_{I_k(r)} V. \end{aligned} \quad (4.21)$$

Suppose for contradiction that (4.20) fails for some  $\ell_1, \ell_2 \in \{1, \dots, \tilde{N}\}$  with  $\ell_1 \leq \ell_2$ , that is,

$$\min \left\{ \max_{\cup_{\ell=\ell_1}^{\ell_2} [\alpha_\ell, \beta_\ell]} V - \min_{(\beta_{\ell_1-1}, \alpha_{\ell_1})} V, \max_{\cup_{\ell=\ell_1}^{\ell_2} [\alpha_\ell, \beta_\ell]} V - \min_{(\beta_{\ell_2}, \alpha_{\ell_2+1})} V \right\} > r.$$

Let  $x_* \in \cup_{\ell=\ell_1}^{\ell_2} [\alpha_\ell, \beta_\ell]$  be such that  $V(x_*) = \max_{\cup_{\ell=\ell_1}^{\ell_2} [\alpha_\ell, \beta_\ell]} V$ . Then, the third identity in (4.21) ensures that there must hold

$$\text{either } I_k(r) \subset (\inf \tilde{I}_k(r), x_*) \text{ or } I_k(r) \subset (x_*, \sup \tilde{I}_k(r)). \quad (4.22)$$

If the first inclusion in (4.22) is the case, we deduce from the second identity in (4.21) that

$$V(\sup \tilde{I}_k(r)) = \sup_{(\inf I_k(r), \sup \tilde{I}_k(r))} V \geq \sup_{(x_*, \sup \tilde{I}_k(r))} V \geq V(x_*) > \min_{(\beta_{\ell_2}, \alpha_{\ell_2+1})} V + r.$$

As a result, there exists a  $r$ -valley contained in  $(x_*, \sup \tilde{I}_k(r))$ , which contradicts the fact that  $I_k(r)$  is the only  $r$ -valley contained in  $\tilde{I}_k(r)$ . Similarly, a contradiction can be deduced if the second inclusion in (4.22) is the case. Hence, (4.20) is true.

Obviously, (4.20) implies that for each  $\ell_1, \ell_2 \in \{1, \dots, \tilde{N}\}$  with  $\ell_1 \leq \ell_2$  there holds

$$\max_{\cup_{\ell=\ell_1}^{\ell_2} [\alpha_\ell, \beta_\ell]} V - \min_{(\beta_{\ell_1-1}, \alpha_{\ell_1})} V - \min_{(\beta_{\ell_2}, \alpha_{\ell_2+1})} V \leq r - \inf_{\tilde{I}_k(r)} V.$$

This together with (4.19) implies (4.18). The claim (4.17) follows.

**Step 3.** By the min-max principle, there holds for each  $0 < \epsilon \ll 1$  and  $r \in (r_{i_*}, r_{i_*} + \delta_*)$

$$\lambda_{\epsilon, N(r)+1} \geq \inf \left\{ \frac{\frac{\epsilon^2}{2} \int_0^\infty a |\phi'|^2 u_\epsilon^G dx}{\int_0^\infty \phi^2 u_\epsilon^G dx} : \phi \in C_0^\infty((0, \infty)), \int_{\tilde{I}_k(r)} \phi u_\epsilon^G dx = 0, \forall k \in \{1, \dots, N(r)\} \right\}.$$

Fix  $0 < \epsilon \ll 1$  and  $r \in (r_{i_*}, r_{i_*} + \delta_*)$ . Let  $\phi \in C_0^\infty((0, \infty))$  satisfy  $\int_{\tilde{I}_k(r)} \phi u_\epsilon^G dx = 0$  for all  $k = 1, \dots, N(r)$ . According to (4.6),

$$\int_{\tilde{I}_0(r)} \phi^2 u_\epsilon^G dx \leq \frac{1}{\lambda_\epsilon^{\tilde{I}_0(r), R}} \frac{\epsilon^2}{2} \int_{\tilde{I}_0(r)} a |\phi'|^2 u_\epsilon^G dx.$$

According to (4.7), (4.8) and the choice of  $\phi$ , we find

$$\sum_{k=1}^{N(r)} \int_{\tilde{I}_k(r)} \phi^2 u_\epsilon^G dx \leq \sum_{k=1}^{N(r)} \frac{1}{\lambda_\epsilon^{\tilde{I}_k(r), R}} \frac{\epsilon^2}{2} \int_{\tilde{I}_k(r)} a |\phi'|^2 u_\epsilon^G dx.$$

Note that  $\int_0^\infty \phi^2 u_\epsilon^G dx = \int_{\tilde{I}_0(r)} \phi^2 u_\epsilon^G dx + \sum_{k=1}^{N(r)} \int_{\tilde{I}_k(r)} \phi^2 u_\epsilon^G dx$ . It follows that

$$\frac{\frac{\epsilon^2}{2} \int_0^\infty a |\phi'|^2 u_\epsilon^G dx}{\int_0^\infty \phi^2 u_\epsilon^G dx} \geq \min_{k \in \{0, 1, \dots, N(r)\}} \lambda_\epsilon^{\tilde{I}_k(r), R},$$

implying  $\lambda_{\epsilon, N(r)+1} \geq \min_{k \in \{0, 1, \dots, N(r)\}} \lambda_\epsilon^{\tilde{I}_k(r), R}$  for all  $0 < \epsilon \ll 1$ , which together with (4.17) yields

$$\liminf_{\epsilon \rightarrow 0} \epsilon^2 \ln \lambda_{\epsilon, N(r)+1} \geq - \max_{k \in \{0, 1, \dots, N(r)\}} r^{\tilde{I}_k(r), R} \geq -r.$$

This proves (4.11), and hence, finishes the proof.  $\square$

## 5. Non-exponential asymptotics of eigenvalues

In Subsection 5.1, we state an auxiliary result on the asymptotics of eigenvalues of a family of Schrödinger operators. This is used in Subsection 5.2 to prove Theorem B. In Subsection 5.3, we establish auxiliary upper bounds of the eigenvalues  $\lambda_{\epsilon,i}$ ,  $i \in \mathbb{N}$ .

**5.1. Asymptotics of Schrödinger eigenvalues.** Recall that  $y_\infty$  is defined in Subsection 2.2. Consider the following family of one-dimensional Schrödinger operators

$$H_h = -h^2 \frac{d^2}{dy^2} + f(y) + hg(y) : L^2((0, y_\infty)) \rightarrow L^2((0, y_\infty)), \quad (5.1)$$

where  $0 < h \ll 1$ ,  $f : (0, y_\infty) \rightarrow [0, \infty)$  and  $g : (0, y_\infty) \rightarrow \mathbb{R}$  are continuous functions and satisfy:

- $\lim_{y \rightarrow y_\infty^-} f(y) = \infty$ ;  $f$  admits a unique zero  $y^*$ , and is twice continuously differentiable near  $y^*$  with  $f''(y^*) > 0$ ;
- $g$  is lower bounded, and satisfies  $g \leq C_1 f + C_2$  on  $(0, y_\infty)$  for some  $C_1, C_2 > 0$ .

For each  $0 < h \ll 1$ , the spectrum of  $H_h$ , as a self-adjoint realization with core  $C_0^\infty((0, \infty))$ , is purely discrete and is denoted by:  $E_{h,1} < E_{h,2} < E_{h,3} \cdots \rightarrow \infty$ . The following result concerning the asymptotics of  $E_{h,i}$  as  $h \rightarrow 0$  is well-known (see e.g. [14, 54, 37]).

**Proposition 5.1.** *For each  $i \in \mathbb{N}$ , there holds  $\lim_{h \rightarrow 0^+} \frac{E_{h,i}}{h} = (2i - 1) \sqrt{\frac{f''(y^*)}{2}} + g(y^*)$ .*

**5.2. Proof of Theorem B.** Let the assumptions in Theorem B be satisfied. Recall  $\mathcal{L}_\epsilon^S$  from (2.1) and consider the Schrödinger operator

$$H_\epsilon := -2\epsilon^2 \mathcal{L}_\epsilon^S = -\epsilon^4 \frac{d^2}{dy^2} + \left[ \frac{b^2}{a} + \epsilon^2 \left( b' - \frac{a'b}{a} \right) + \epsilon^4 \left( \frac{3}{16} \frac{(a')^2}{a} - \frac{a''}{4} \right) \right] (\xi^{-1}(y)). \quad (5.2)$$

Denote by  $E_{\epsilon,1} < E_{\epsilon,2} < E_{\epsilon,3} < \cdots$  the eigenvalues of  $H_\epsilon$ , and recall that the eigenvalues of  $-\mathcal{L}_\epsilon^\epsilon$  are  $\lambda_{\epsilon,1} < \lambda_{\epsilon,2} < \lambda_{\epsilon,3} < \cdots$ . Obviously,

$$E_{\epsilon,i} = 2\epsilon^2 \lambda_{\epsilon,i}, \quad \forall i \in \mathbb{N} \quad \text{and} \quad 0 < \epsilon \ll 1. \quad (5.3)$$

Note that with  $\epsilon^2 = h$ , the operator  $H_\epsilon$  is almost in the form of  $H_h$  given in (5.1). A careful analysis of the extra term  $\epsilon^4 \left( \frac{3}{16} \frac{(a')^2}{a} - \frac{a''}{4} \right) \circ \xi^{-1}$ , which is unbounded near 0, allows us to apply Proposition 5.1 to prove the following result.

**Lemma 5.1.** *For each  $i \in \mathbb{N}$ , there holds  $\lim_{\epsilon \rightarrow 0} \frac{E_{\epsilon,i}}{\epsilon^2} = 2b'(x^*)(1 - i)$ .*

*Proof.* In order to apply Proposition 5.1, we construct operators that are in the form of (5.1) and comparable in the sense of quadratic forms to  $H_\epsilon$ .

Let  $c : (0, \infty) \rightarrow [0, \infty)$  be continuous and satisfy the following conditions:

- $c(x) = \frac{1}{x}$  for  $x \in (0, \frac{x^*}{2})$ ,  $c(x) = \frac{b^2(x)}{a(x)}$  for  $x \gg x^*$ ;
- $c$  is twice continuously differentiable near  $x^*$ , and satisfies  $c(x^*) = c'(x^*) = c''(x^*) = 0$ .

By (H), for each  $0 < \delta \ll 1$ , there exists  $0 < \epsilon_\delta \ll 1$  such that  $\epsilon^4 \left( \frac{3}{16} \frac{(a')^2}{a} - \frac{a''}{4} \right) \leq c + \epsilon^2 \delta$  on  $(0, \infty)$  for all  $\epsilon \in (0, \epsilon_\delta)$ , and thus, in the sense of quadratic forms,

$$H_\epsilon \leq \bar{H}_\epsilon^\delta := -\epsilon^4 \frac{d^2}{dy^2} + \bar{f}(y) + \epsilon^2 \bar{g}^\delta(y), \quad (5.4)$$

where  $\bar{f} = \left( \frac{b^2}{a} + c \right) \circ \xi^{-1}$  and  $\bar{g}^\delta = \left( b' - \frac{a'b}{a} + \delta \right) \circ \xi^{-1}$ .

Obviously,  $\bar{f}$  is continuously differentiable, satisfies  $\bar{f}(y) \rightarrow \infty$  as  $y \rightarrow y_\infty^-$ , has  $y^* := \xi(x^*)$  as the unique zero, and is twice continuously differentiable near  $y^*$  with  $\bar{f}''(y^*) = 2(b'(x^*))^2$ . By **(H)** and the construction of the function  $c$ , it is easy to see that  $\bar{g}^\delta$  is lower bounded, and there are  $C_1, C_2 > 0$  such that  $\bar{g}^\delta \leq C_1 \bar{f} + C_2$  for all  $0 < \delta \ll 1$ .

Denote by  $\bar{E}_{\epsilon,1}^\delta \leq \bar{E}_{\epsilon,2}^\delta \leq \dots$  the eigenvalues of  $\bar{H}_\epsilon^\delta$ . We apply Proposition 5.1 to conclude that

$$\lim_{\epsilon \rightarrow 0} \frac{\bar{E}_{\epsilon,i}^\delta}{\epsilon^2} = (2i-1) \sqrt{\frac{\bar{f}''(y^*)}{2}} + \bar{g}^\delta(y^*) = 2b'(x^*)(1-i) + \delta, \quad \forall 0 < \delta \ll 1 \text{ and } i \in \mathbb{N}.$$

It follows from (5.4) that  $E_{\epsilon,i} \leq \bar{E}_{\epsilon,i}^\delta$ , leading to  $\limsup_{\epsilon \rightarrow 0} \frac{E_{\epsilon,i}}{\epsilon^2} \leq 2b'(x^*)(1-i)$  for all  $i \in \mathbb{N}$ .

It remains to establish the lower bound. Fix  $x_0 \gg x^*$  and let  $\eta : (0, \infty) \rightarrow [0, \frac{1}{2}]$  be smooth and satisfy  $\eta(x) = 0$  for  $x \leq x_0$  and  $\eta(x) = \frac{1}{2}$  for  $x \geq 2x_0$ . Then, for each  $0 < \delta \ll 1$ , there exists  $0 < \epsilon_\delta \ll 1$  such that  $\eta \frac{b^2}{a} + \epsilon^4 \left( \frac{3}{16} \frac{(a')^2}{a} - \frac{a''}{4} \right) \geq -\epsilon^2 \delta$  on  $(0, \infty)$  for all  $\epsilon \in (0, \epsilon_\delta)$ , which gives

$$H_\epsilon \geq \underline{H}_\epsilon^\delta := -\epsilon^4 \frac{d^2}{dy^2} + \underline{f}(y) + \epsilon^2 \underline{g}^\delta(y), \quad (5.5)$$

where  $\underline{f} = \left( (1-\eta) \frac{b^2}{a} \right) \circ \xi^{-1}$  and  $\underline{g}^\delta = \left( b' - \frac{a'b}{a} - \delta \right) \circ \xi^{-1}$ .

Denote by  $\underline{E}_{\epsilon,1}^\delta \leq \underline{E}_{\epsilon,2}^\delta \leq \dots$  the eigenvalues of  $\underline{H}_\epsilon^\delta$ . We apply Proposition 5.1 to conclude that

$$\lim_{\epsilon \rightarrow 0} \frac{\underline{E}_{\epsilon,i}^\delta}{\epsilon^2} = (2i-1) \sqrt{\frac{\underline{f}''(y^*)}{2}} + \underline{g}^\delta(y^*) = 2b'(x^*)(1-i) - \delta, \quad \forall 0 < \delta \ll 1 \text{ and } i \in \mathbb{N}.$$

It follows from (5.5) that  $\liminf_{\epsilon \rightarrow 0} \frac{E_{\epsilon,i}}{\epsilon^2} \geq 2b'(x^*)(1-i)$  for all  $i \in \mathbb{N}$ . This completes the proof.  $\square$

Theorem B now is a simple consequence of Lemma 5.1.

**Proof of Theorem B.** By Lemma 5.1 and (5.3),  $\lim_{\epsilon \rightarrow 0} \lambda_{\epsilon,i} = b'(x^*)(1-i)$  for all  $i \in \mathbb{N}$ .  $\square$

**5.3. Upper bounds of eigenvalues.** In this subsection, we use Proposition 5.1 to establish non-exponential upper bounds of the eigenvalues  $\lambda_{\epsilon,i}$ ,  $i \in \mathbb{N}$  that complement Theorem A, which provides almost no information about  $\lambda_{\epsilon,i}$  when  $r_i = 0$ .

**Proposition 5.2.** *Assume (H). There are  $\Lambda_1, \Lambda_2 > 0$  such that  $\lambda_{\epsilon,i} \leq \Lambda_1 i + \Lambda_2$  for all  $i \in \mathbb{N}$  and  $0 < \epsilon \ll 1$ .*

*Proof.* Recall that  $\lambda_{\epsilon,i}$ ,  $i \in \mathbb{N}$  are also the eigenvalues of the Schrödinger operator  $-\mathcal{L}_\epsilon^S$ . The operator  $H_\epsilon$  is given in (5.2). Its eigenvalues  $E_{\epsilon,i}$ ,  $i \in \mathbb{N}$  satisfy  $E_{\epsilon,i} = 2\epsilon^2 \lambda_{\epsilon,i}$  for all  $i \in \mathbb{N}$ .

Arguing as in the proof of Lemma 5.1, it is not hard to find smooth functions  $\tilde{f} : (0, y_\infty) \rightarrow [0, \infty)$  and  $\tilde{g} : (0, y_\infty) \rightarrow \mathbb{R}$  satisfying

- $\tilde{f}(y) \rightarrow \infty$  as  $y \rightarrow 0^+$  and  $y \rightarrow y_\infty^-$ ;  $\tilde{f}$  admits a unique zero at  $\tilde{y}_*$  and satisfies  $\tilde{f}''(\tilde{y}_*) > 0$ ,
- $\tilde{g}$  is lower bounded and there are constants  $C_1, C_2 > 0$  such that  $\tilde{g} \leq C_1 \tilde{f} + C_2$ ,

such that  $H_\epsilon \leq \tilde{H}_\epsilon := -\epsilon^4 \frac{d^2}{dy^2} + \tilde{f}(y) + \epsilon^2 \tilde{g}(y)$  in the sense of quadratic forms. Denote by  $\tilde{E}_{\epsilon,1} < \tilde{E}_{\epsilon,2} < \tilde{E}_{\epsilon,3} \dots \rightarrow \infty$  the eigenvalues of  $\tilde{H}_\epsilon$ . We apply Proposition 5.1 with  $h = \epsilon^2$  to  $\tilde{H}_\epsilon$  to conclude

$$\lim_{\epsilon \rightarrow 0} \frac{\tilde{E}_{\epsilon,i}}{\epsilon^2} = (2i-1) \sqrt{\frac{\tilde{f}''(\tilde{y}_*)}{2}} + \tilde{g}(\tilde{y}_*), \quad \forall i \in \mathbb{N}.$$

It follows that  $\limsup_{\epsilon \rightarrow 0} \lambda_{\epsilon,i} \leq \frac{2i-1}{2} \sqrt{\frac{\tilde{f}''(\tilde{y}_*)}{2}} + \frac{\tilde{g}(\tilde{y}_*)}{2}$  for all  $i \in \mathbb{N}$ . This completes the proof.  $\square$

## 6. Qualitative characterizations of multi-scale dynamics

In this section, we study qualitative characterization of the multi-scale dynamics of  $X_t^\epsilon$ , and prove Theorem C.

We need the following result. Recall from Proposition 2.1 that  $(P_t^\epsilon)_{t \geq 0}$  is the positive analytic semigroup generated by  $\mathcal{L}_\epsilon$ , and  $Q_k^\epsilon$  is the spectral projection of  $\mathcal{L}_\epsilon$  corresponding to  $\{\lambda_{\epsilon,j}\}_{j \geq k}$ .

**Lemma 6.1.** *For each  $k \in \mathbb{N}$ , there hold the following statements.*

(1) *There exists  $C > 0$  such that for each  $0 < \epsilon \ll 1$ ,*

$$|P_t^\epsilon Q_k^\epsilon f| \leq \frac{C}{\epsilon} a^{\frac{1}{4}} e^{\frac{V}{2\epsilon^2}} e^{-\lambda_{\epsilon,k} t} \|f\|_{L^2(u_\epsilon^G)} \quad \text{in } (0, \infty), \quad \forall f \in L^2(u_\epsilon^G) \text{ and } t > 1.$$

(2) *There exist  $C > 0$  such that for each  $0 < \epsilon \ll 1$ ,*

$$|P_t^\epsilon Q_k^\epsilon f| \leq a^{\frac{1}{4}} e^{\frac{V+C}{2\epsilon^2}} e^{-\lambda_{\epsilon,k} t} \|f\|_\infty \quad \text{in } (0, \infty), \quad \forall f \in C_b([0, \infty)) \text{ and } t > 2.$$

*Proof.* Fix  $k \in \mathbb{N}$  and let  $0 < \epsilon \ll 1$ . For  $t \geq 0$ , we define  $\tilde{P}_t^\epsilon := \tilde{U}_\epsilon U_\epsilon P_t^\epsilon U_\epsilon^{-1} \tilde{U}_\epsilon^{-1}$ , where  $U_\epsilon$  and  $\tilde{U}_\epsilon$  are unitary transforms specified in Subsection 2.2. Then,  $(\tilde{P}_t^\epsilon)_{t \geq 0}$  is an analytic semigroup of contractions on  $L^2((0, y_\infty))$  with generator  $\mathcal{L}_\epsilon^S$ . The spectrum of  $\mathcal{L}_\epsilon^S$  is the same as that of  $\mathcal{L}_\epsilon$  consisting of simple eigenvalues  $\{-\lambda_{\epsilon,k}\}_{k \in \mathbb{N}}$ .

(1) Set  $M := 1 + |\inf_{0 < \epsilon \ll 1} \min V_\epsilon|$ , which is finite thanks to Lemma 2.2 (4). We claim that for each  $p \in (2, \infty]$ , there is  $\tilde{C}_1 = \tilde{C}_1(p) > 0$  such that

$$\|\tilde{P}_t^\epsilon \tilde{f}\|_{L^p} \leq \frac{\tilde{C}_1}{t^\epsilon} e^{Mt} \|\tilde{f}\|_{L^2}, \quad \forall \tilde{f} \in L^2((0, y_\infty)) \text{ and } t > 0. \quad (6.1)$$

Since  $\|(\lambda - \mathcal{L}_\epsilon^S)^{-1}\|_{L^2 \rightarrow L^2} = \frac{1}{\text{dist}(\lambda, \sigma(\mathcal{L}_\epsilon^S))}$  for all  $\lambda \in \rho(\mathcal{L}_\epsilon^S)$  and  $\sigma(\mathcal{L}_\epsilon^S) \subset (-\infty, 0)$ , we find

$$\|(\lambda - (\mathcal{L}_\epsilon^S - M))^{-1}\|_{L^2 \rightarrow L^2} \leq \frac{1}{|\lambda|}, \quad \forall \lambda \in \mathbb{C} \text{ with } \Re \lambda > 0. \quad (6.2)$$

As  $\mathcal{L}_\epsilon^S - M$  generates the analytic semigroup  $(e^{-Mt} \tilde{P}_t^\epsilon)_{t \geq 0}$  of contractions on  $L^2((0, y_\infty))$ , and the right-hand side of (6.2) is independent of  $\epsilon$ , we apply [50, Theorem 2.5.2] to find  $C_1 > 0$  such that

$$\|(\mathcal{L}_\epsilon^S - M) e^{-Mt} \tilde{P}_t^\epsilon \tilde{f}\|_{L^2} \leq \frac{C_1}{t} \|\tilde{f}\|_{L^2}, \quad \forall \tilde{f} \in L^2((0, y_\infty)) \text{ and } t > 0. \quad (6.3)$$

Let  $D(\mathcal{L}_\epsilon^S)$  be the domain of  $\mathcal{L}_\epsilon^S$ . Note that

$$\langle -(\mathcal{L}_\epsilon^S - M)u, u \rangle_{L^2} = \frac{\epsilon^2}{2} \int_0^{y_\infty} |u'|^2 dy + \int_0^{y_\infty} (V_\epsilon + M)|u|^2 dy, \quad \forall u \in D(\mathcal{L}_\epsilon^S),$$

which together with (6.3) and the fact  $\inf_{0 < \epsilon \ll 1} (\min V_\epsilon + M) \geq 1$  leads to

$$\begin{aligned} & \frac{\epsilon^2}{2} \int_0^{y_\infty} |\partial_y \tilde{P}_t^\epsilon \tilde{f}|^2 dy + \int_0^{y_\infty} (V_\epsilon + M) |\tilde{P}_t^\epsilon \tilde{f}|^2 dy \\ & \leq \|(\mathcal{L}_\epsilon^S - M) \tilde{P}_t^\epsilon \tilde{f}\|_{L^2} \|\tilde{P}_t^\epsilon \tilde{f}\|_{L^2} \\ & \leq \frac{C_1 e^{Mt}}{t} \|\tilde{f}\|_{L^2} \|\tilde{P}_t^\epsilon \tilde{f}\|_{L^2} \leq \frac{C_1 e^{Mt}}{t} \|\tilde{f}\|_{L^2} \left( \frac{\epsilon^2}{2} \int_0^{y_\infty} |\partial_y \tilde{P}_t^\epsilon \tilde{f}|^2 dy + \int_0^{y_\infty} (V_\epsilon + M) |\tilde{P}_t^\epsilon \tilde{f}|^2 dy \right)^{\frac{1}{2}}, \end{aligned}$$

yielding

$$\frac{\epsilon^2}{2} \int_0^{y_\infty} |\partial_y \tilde{P}_t^\epsilon \tilde{f}|^2 dy + \int_0^{y_\infty} (V_\epsilon + M) |\tilde{P}_t^\epsilon \tilde{f}|^2 dy \leq \frac{C_1^2 e^{2Mt}}{t^2} \|\tilde{f}\|_{L^2}^2, \quad \forall \tilde{f} \in L^2((0, y_\infty)), \quad t > 0.$$

By Sobolev's embedding theorem, we find for each  $p \in (2, \infty]$  the existence of  $C_2(p) > 0$  such that

$$\|\tilde{P}_t^\epsilon \tilde{f}\|_{L^p} \leq C_2(p) \left( \|\partial_y \tilde{P}_t^\epsilon \tilde{f}\|_{L^2} + \|\tilde{P}_t^\epsilon \tilde{f}\|_{L^2} \right) \leq \frac{2C_1 C_2(p)}{t\epsilon} e^{Mt} \|\tilde{f}\|_{L^2}, \quad \forall \tilde{f} \in L^2((0, y_\infty)), \quad t > 0.$$

This gives (6.1) with  $\tilde{C}_1 := 2C_1 C_2(p)$ .

Now, let  $f \in L^2(u_\epsilon^G)$  and set  $\tilde{f}_\epsilon := \tilde{U}_\epsilon U_\epsilon f$ . Note that  $\|\tilde{P}_t^\epsilon \tilde{Q}_k^\epsilon \tilde{f}_\epsilon\|_{L^2} \leq e^{-\lambda_{\epsilon,k} t} \|\tilde{f}_\epsilon\|_{L^2}$  for all  $t > 0$  and  $0 < \epsilon \ll 1$ , where  $\tilde{Q}_k^\epsilon := \tilde{U}_\epsilon U_\epsilon Q_k^\epsilon U_\epsilon^{-1} \tilde{U}_\epsilon^{-1}$  is the spectral projection of  $\mathcal{L}_\epsilon^S$  corresponding to  $\{-\lambda_{\epsilon,j}\}_{j \geq k}$ . Applying (6.1) with  $p = \infty$ , we deduce

$$\|\tilde{P}_t^\epsilon \tilde{Q}_k^\epsilon \tilde{f}_\epsilon\|_{L^\infty} \leq \frac{\tilde{C}_1 e^M}{\epsilon} \|\tilde{P}_{t-1}^\epsilon \tilde{Q}_k^\epsilon \tilde{f}_\epsilon\|_{L^2} \leq \frac{\tilde{C}_1 e^M}{\epsilon} e^{-\lambda_{\epsilon,k}(t-1)} \|\tilde{f}_\epsilon\|_{L^2}, \quad \forall t > 1, \quad (6.4)$$

leading to

$$\begin{aligned} |P_t^\epsilon Q_k^\epsilon f| &= |U_\epsilon^{-1} \tilde{U}_\epsilon^{-1} \tilde{P}_t^\epsilon \tilde{Q}_k^\epsilon \tilde{f}_\epsilon| \\ &= |(\tilde{P}_t^\epsilon \tilde{Q}_k^\epsilon \tilde{f}_\epsilon) \circ \xi| \frac{\sqrt{\xi'}}{\sqrt{u_\epsilon^G}} \leq \|\tilde{P}_t^\epsilon \tilde{Q}_k^\epsilon \tilde{f}_\epsilon\|_{L^\infty} a^{\frac{1}{4}} e^{\frac{V}{2\epsilon^2}} \leq \frac{\tilde{C}_1 e^{M+\lambda_{\epsilon,1}}}{\epsilon} a^{\frac{1}{4}} e^{\frac{V}{2\epsilon^2}} e^{-\lambda_{\epsilon,k} t} \|f\|_{L^2(u_\epsilon^G)}, \quad \forall t > 1. \end{aligned}$$

Since  $\lambda_{\epsilon,1} \rightarrow 0$  as  $\epsilon \rightarrow 0$  by Theorem A, the conclusion of (1) follows.

(2) Fix  $f \in C_b([0, \infty))$  and set  $\tilde{f}_\epsilon := \tilde{U}_\epsilon U_\epsilon f$ . We claim that there exist  $\tilde{C}_2, \tilde{C}_3 > 0$  such that

$$\|\tilde{P}_1^\epsilon \tilde{f}_\epsilon\|_{L^2} \leq \frac{\tilde{C}_2}{\epsilon} e^{\frac{\tilde{C}_3}{\epsilon^2}} \|f\|_\infty. \quad (6.5)$$

To do so, we fix  $p > 2$ . By (6.1),  $\tilde{P}_1^\epsilon : L^2((0, y_\infty)) \rightarrow L^p((0, y_\infty))$  is linear and bounded and satisfies  $\|\tilde{P}_1^\epsilon\|_{L^2 \rightarrow L^p} \leq \frac{\tilde{C}_1}{\epsilon} e^M$ . This together with the symmetry of  $\tilde{P}_1^\epsilon : L^2((0, y_\infty)) \rightarrow L^2((0, y_\infty))$  yields

$$\|\tilde{P}_1^\epsilon \tilde{f}\|_{L^2} \leq \frac{\tilde{C}_1}{\epsilon} e^M \|\tilde{f}\|_{L^{p'}}, \quad \forall \tilde{f} \in L^2((0, y_\infty)) \cap L^{p'}((0, y_\infty)), \quad (6.6)$$

where  $p' = \frac{p}{p-1} \in (1, 2)$  is the dual exponent of  $p$ . As a consequence,  $\tilde{P}_1^\epsilon$  uniquely extends to be a bounded linear operator from  $L^{p'}((0, y_\infty))$  to  $L^2((0, y_\infty))$ , and satisfies  $\|\tilde{P}_1^\epsilon\|_{L^{p'} \rightarrow L^2} \leq \frac{\tilde{C}_1}{\epsilon} e^M$ .

Straightforward calculations give

$$\begin{aligned} \int_0^{y_\infty} |\tilde{f}_\epsilon|^{p'} dy &= \int_0^{y_\infty} |f \circ \xi^{-1}|^{p'} \left| \frac{u_\epsilon^G \circ \xi^{-1}}{\xi' \circ \xi^{-1}} \right|^{\frac{p'}{2}} dy \\ &= \int_0^\infty |f|^{p'} |u_\epsilon^G|^{\frac{p'}{2}} |\xi'|^{1-\frac{p'}{2}} dx = \int_0^\infty |f|^{p'} a^{-\frac{1}{2}-\frac{p'}{4}} e^{-\frac{p'V}{2\epsilon^2}} dx. \end{aligned}$$

Since  $\int_0^\infty a^{-\frac{1}{2}-\frac{p'}{4}} e^{-\frac{p'V}{2\epsilon^2}} dx \leq C_3^{p'} e^{\frac{C_4 p'}{\epsilon^2}}$  for some  $C_3, C_4 > 0$ , we find  $\|\tilde{f}_\epsilon\|_{L^{p'}} \leq C_3 e^{\frac{C_4}{\epsilon^2}} \|f\|_\infty$ , which together with (6.6) gives  $\|\tilde{P}_1^\epsilon \tilde{f}_\epsilon\|_{L^2} \leq \frac{\tilde{C}_1 C_3}{\epsilon} e^M e^{\frac{C_4}{\epsilon^2}} \|f\|_\infty$ . The claim then follows readily by setting  $\tilde{C}_2 := \tilde{C}_1 C_3 e^M$  and  $\tilde{C}_3 := C_4$ .

For  $t > 2$ , we deduce from (6.4) and (6.5) that

$$\begin{aligned} \|\tilde{P}_t^\epsilon \tilde{Q}_k^\epsilon \tilde{f}_\epsilon\|_{L^\infty} &\leq \frac{\tilde{C}_1 e^M}{\epsilon} \|\tilde{P}_{t-1}^\epsilon \tilde{Q}_k^\epsilon \tilde{f}_\epsilon\|_{L^2} \\ &\leq \frac{\tilde{C}_1 e^M}{\epsilon} e^{-\lambda_{\epsilon,k}(t-2)} \|\tilde{P}_1^\epsilon \tilde{Q}_k^\epsilon \tilde{f}_\epsilon\|_{L^2} \leq \frac{\tilde{C}_1 \tilde{C}_2 e^{M+2\lambda_{\epsilon,k}}}{\epsilon^2} e^{-\lambda_{\epsilon,k} t} e^{\frac{\tilde{C}_3}{\epsilon^2}} \|f\|_\infty, \end{aligned}$$

and hence,

$$|P_t^\epsilon Q_k^\epsilon f| = |U_\epsilon^{-1} \tilde{U}_\epsilon^{-1} \tilde{P}_t^\epsilon \tilde{Q}_k^\epsilon \tilde{f}_\epsilon| = |(\tilde{P}_t^\epsilon \tilde{Q}_k^\epsilon \tilde{f}_\epsilon) \circ \xi| \frac{\sqrt{\xi'}}{\sqrt{u_\epsilon^G}} \leq \frac{\tilde{C}_1 \tilde{C}_2 e^{M+2\lambda_{\epsilon,k}}}{\epsilon^2} e^{-\lambda_{\epsilon,k} t} e^{\frac{C_3}{\epsilon^2}} \|f\|_\infty \times a^{\frac{1}{4}} e^{\frac{V}{2\epsilon^2}}.$$

By Proposition 5.2, it is easy to see the existence of  $\tilde{C}_4 > 0$  (depending on  $k$ ) such that  $|P_t^\epsilon Q_k^\epsilon f| \leq a^{\frac{1}{4}} e^{\frac{V+\tilde{C}_4}{2\epsilon^2}} e^{-\lambda_{\epsilon,k} t} \|f\|_\infty$  for all  $t > 2$  and  $0 < \epsilon \ll 1$ . This completes the proof.  $\square$

**Proof of Theorem C.** (1) Let  $K \subset [0, \infty)$  be a fixed compact set, and  $\mu \in \mathcal{P}([0, \infty))$  satisfy  $\text{supp}(\mu) \subset K$ . Fix  $f \in C_b([0, \infty))$  and  $0 < \epsilon \ll 1$ . Then, Proposition 2.1 (5) gives

$$\begin{aligned} \mathbb{E}_\mu^\epsilon[f(X_t^\epsilon) \mathbb{1}_{t < T_0^\epsilon}] &= \langle f, \phi_{\epsilon,1} \rangle_{L^2(u_\epsilon^G)} \|\phi_{\epsilon,1}\|_{L^1(\mu)} e^{-\lambda_{\epsilon,1} t} + \int_0^\infty P_t^\epsilon Q_2^\epsilon f d\mu \\ &= \alpha_\epsilon(\mu) e^{-\lambda_{\epsilon,1} t} \int_0^\infty f d\mu_\epsilon + \int_0^\infty P_t^\epsilon Q_2^\epsilon f d\mu, \quad \forall t > 0, \end{aligned}$$

where  $\alpha_\epsilon(\mu) := \|\phi_{\epsilon,1}\|_{L^1(u_\epsilon^G)} \|\phi_{\epsilon,1}\|_{L^1(\mu)}$  and  $\mu_\epsilon$  is the unique QSD of  $X_t^\epsilon$  (see Proposition 2.3). In particular, setting  $f = \mathbb{1}$  yields

$$\mathbb{P}_\mu^\epsilon[t < T_0^\epsilon] = \alpha_\epsilon(\mu) e^{-\lambda_{\epsilon,1} t} + \int_0^\infty P_t^\epsilon Q_2^\epsilon \mathbb{1} d\mu, \quad \forall t > 0. \quad (6.7)$$

It follows that

$$\begin{aligned} \mathbb{E}_\mu^\epsilon[f(X_t^\epsilon)] &= \mathbb{E}_\mu^\epsilon[f(X_t^\epsilon) \mathbb{1}_{t < T_0^\epsilon}] + \mathbb{E}_\mu^\epsilon[f(0) \mathbb{1}_{t \geq T_0^\epsilon}] \\ &= \alpha_\epsilon(\mu) e^{-\lambda_{\epsilon,1} t} \int_0^\infty f d\mu_\epsilon + \int_0^\infty P_t^\epsilon Q_2^\epsilon f d\mu + f(0) \left(1 - \alpha_\epsilon(\mu) e^{-\lambda_{\epsilon,1} t} - \int_0^\infty P_t^\epsilon Q_2^\epsilon \mathbb{1} d\mu\right) \\ &= \alpha_\epsilon(\mu) e^{-\lambda_{\epsilon,1} t} \int_0^\infty f d\mu_\epsilon + (1 - \alpha_\epsilon(\mu) e^{-\lambda_{\epsilon,1} t}) f(0) \\ &\quad + \int_0^\infty P_t^\epsilon Q_2^\epsilon f d\mu - f(0) \int_0^\infty P_t^\epsilon Q_2^\epsilon \mathbb{1} d\mu, \quad \forall t > 0. \end{aligned}$$

Lemma 6.1 (2) (with  $k = 2$ ) yields the existence of  $C_1 > 0$  such that

$$\begin{aligned} &\left| \mathbb{E}_\mu^\epsilon[f(X_t^\epsilon)] - \left[ \alpha_\epsilon(\mu) e^{-\lambda_{\epsilon,1} t} \int_0^\infty f d\mu_\epsilon + (1 - \alpha_\epsilon(\mu) e^{-\lambda_{\epsilon,1} t}) f(0) \right] \right| \\ &\leq \int_0^\infty (|P_t^\epsilon Q_2^\epsilon f| + |P_t^\epsilon Q_2^\epsilon \mathbb{1}| |f(0)|) d\mu \leq 2 \int_0^\infty a^{\frac{1}{4}} e^{\frac{V+C_1}{2\epsilon^2}} d\mu e^{-\lambda_{\epsilon,2} t} \|f\|_\infty, \quad \forall t > 2. \end{aligned}$$

Clearly, there is  $C_2 = C_2(K) > 0$  such that  $2 \int_0^\infty a^{\frac{1}{4}} e^{\frac{V+C_1}{2\epsilon^2}} d\mu \leq e^{\frac{C_2}{\epsilon^2}}$ , leading to

$$\left\| \mathbb{P}_\mu^\epsilon[X_t^\epsilon \in \bullet] - [\alpha_\epsilon(\mu) e^{-\lambda_{\epsilon,1} t} \mu_\epsilon + (1 - \alpha_\epsilon(\mu) e^{-\lambda_{\epsilon,1} t}) \delta_0] \right\|_{TV} \leq \exp \left\{ \frac{C_2}{\epsilon^2} - \lambda_{\epsilon,2} t \right\}, \quad \forall t > 2.$$

It remains to estimate  $\alpha_\epsilon(\mu)$ . Note from (6.7) that

$$\begin{aligned} \alpha_\epsilon(\mu) &= e^{\lambda_{\epsilon,1} t} \left( \mathbb{P}_\mu^\epsilon[t < T_0^\epsilon] - \int_0^\infty P_t^\epsilon Q_2^\epsilon \mathbb{1} d\mu \right) \\ &\leq e^{\lambda_{\epsilon,1} t} \left( 1 + \int_0^\infty a^{\frac{1}{4}} e^{\frac{V+C_1}{2\epsilon^2}} d\mu e^{-\lambda_{\epsilon,2} t} \right) \leq e^{\lambda_{\epsilon,1} t} \left( 1 + e^{\frac{C_2}{\epsilon^2} - \lambda_{\epsilon,2} t} \right), \quad \forall t > 0. \end{aligned}$$



Since the function  $t \mapsto e^{\lambda_{\epsilon,1}t} \left(1 + e^{\frac{C_2}{\epsilon^2} - \lambda_{\epsilon,2}t}\right)$  is minimized at  $t = \left(\frac{C_2}{\epsilon^2} - \ln \frac{\lambda_{\epsilon,1}}{\lambda_{\epsilon,2} - \lambda_{\epsilon,1}}\right) \frac{1}{\lambda_{\epsilon,2}}$ , we find

$$\alpha_{\epsilon}(\mu) \leq \exp \left\{ \left( \frac{C_2}{\epsilon^2} - \ln \frac{\lambda_{\epsilon,1}}{\lambda_{\epsilon,2} - \lambda_{\epsilon,1}} \right) \frac{\lambda_{\epsilon,1}}{\lambda_{\epsilon,2}} \right\} \left( 1 + \frac{\lambda_{\epsilon,1}}{\lambda_{\epsilon,2} - \lambda_{\epsilon,1}} \right).$$

By the asymptotics of  $\lambda_{\epsilon,1}$  and  $\lambda_{\epsilon,2}$  (see Theorem A), we find the existence of some small  $\epsilon_* = \epsilon_*(C_2) = \epsilon_*(K) > 0$  such that  $\alpha_{\epsilon}(\mu) \leq 1 + e^{-\frac{r_1 - r_2}{2\epsilon^2}}$  for all  $\epsilon \in (0, \epsilon_*)$ . This proves (1).

(2) We prove the result in the global attractor case; the local attractor case can be treated in the same manner. Let  $0 < \delta \ll 1$  and  $K_1, K_2$  be compact sets in  $(0, \infty)$ . Let  $\mu \in \mathcal{P}([0, \infty))$  satisfy  $\text{supp}(\mu) \subset K_1$ , and  $f$  be a continuous function on  $(0, \infty)$  with  $\text{supp}(f) \subset K_2$ .

Recall that  $\mathcal{A}$  is the global attractor of (1.2). Let  $T = T(K_1, \delta) > 0$  be the smallest time such that  $\sup_{x_0 \in K} \text{dist}(x_t, \mathcal{A}) < \frac{\delta}{2}$  for all  $t > T$ , where  $x_t$  denotes the solution of (1.2). By the sample path large deviation principle, there are  $D_1 = D_1(K_1, \delta) > 0$  and  $D_2 = D_2(K_1, \delta) > 0$  such that

$$\sup_{x_0 \in K_1} \mathbb{P}_{x_0}^{\epsilon} \left[ \sup_{t \in [0, T]} |X_t^{\epsilon} - x_t| > \frac{\delta}{2} \right] \leq D_1 e^{-\frac{D_2}{\epsilon^2}}, \quad \forall 0 < \epsilon \ll 1,$$

resulting in

$$\mathbb{P}_{\mu}^{\epsilon} [X_T^{\epsilon} \in \mathcal{A}_{\delta}^c] \leq \mathbb{P}_{\mu}^{\epsilon} \left[ |X_T^{\epsilon} - x_T| > \frac{\delta}{2} \right] \leq D_1 e^{-\frac{D_2}{\epsilon^2}}, \quad \forall 0 < \epsilon \ll 1, \quad (6.8)$$

where  $\mathcal{A}_{\delta} := \{x > 0 : \text{dist}(x, \mathcal{A}) < \delta\}$  and  $\mathcal{A}_{\delta}^c := (0, \infty) \setminus \mathcal{A}_{\delta}$ .

Setting  $\mu_T^{\epsilon} := \mathbb{P}_{\mu}^{\epsilon} [X_T^{\epsilon} \in \bullet]$ , we use the Markov property of  $X_t^{\epsilon}$  to write

$$\begin{aligned} \mathbb{E}_{\mu}^{\epsilon} [f(X_t^{\epsilon})] &= \mathbb{E}_{\mu}^{\epsilon} \left[ \mathbb{E}_{\mu}^{\epsilon} [f(X_t^{\epsilon}) | X_T^{\epsilon}] \right] \\ &= \int_{\mathcal{A}_{\delta}} \mathbb{E}_{\bullet}^{\epsilon} [f(X_{t-T}^{\epsilon})] d\mu_T^{\epsilon} + \int_{\mathcal{A}_{\delta}^c} \mathbb{E}_{\bullet}^{\epsilon} [f(X_{t-T}^{\epsilon})] d\mu_T^{\epsilon} =: E_1^{\epsilon} + E_2^{\epsilon}, \quad \forall t \geq T. \end{aligned} \quad (6.9)$$

We estimate  $E_1^{\epsilon}$ . Since Proposition 2.1 gives

$$\mathbb{E}_{\bullet}^{\epsilon} [f(X_{t-T}^{\epsilon})] - \|\phi_{\epsilon,1}\|_{L^1(u_{\epsilon}^G)} \phi_{\epsilon,1} e^{-\lambda_{\epsilon,1}(t-T)} \int_0^{\infty} f d\mu_{\epsilon} = P_{t-T}^{\epsilon} Q_2^{\epsilon} f, \quad \forall t > T,$$

we apply Lemma 6.1 (1) (with  $k = 2$ ) to find  $D_3 > 0$  such that

$$\begin{aligned} & \left| E_1^{\epsilon} - \|\phi_{\epsilon,1}\|_{L^1(u_{\epsilon}^G)} \int_{\mathcal{A}_{\delta}} \phi_{\epsilon,1} d\mu_T^{\epsilon} e^{-\lambda_{\epsilon,1}(t-T)} \int_0^{\infty} f d\mu_{\epsilon} \right| \\ & \leq \frac{D_3}{\epsilon} e^{-\lambda_{\epsilon,2}(t-T)} \left( \int_{\mathcal{A}_{\delta}} a^{\frac{1}{4}} e^{\frac{V}{2\epsilon^2}} d\mu_T^{\epsilon} \right) \left( \int_0^{\infty} |f|^2 u_{\epsilon}^G dx \right)^{\frac{1}{2}} \\ & = \frac{D_3}{\epsilon} e^{-\lambda_{\epsilon,2}(t-T)} \left( \int_{\mathcal{A}_{\delta}} a^{\frac{1}{4}} e^{\frac{V}{2\epsilon^2}} d\mu_T^{\epsilon} \right) \left( \int_0^{\infty} \frac{|f|^2}{a} e^{-\frac{V}{\epsilon^2}} dx \right)^{\frac{1}{2}} \\ & \leq \frac{D_3}{\epsilon} \left( \sup_{\mathcal{A}_{\delta}} a^{\frac{1}{4}} \right) \left( \int_{K_2} \frac{1}{a} dx \right)^{\frac{1}{2}} \exp \left\{ \frac{\sup_{\mathcal{A}_{\delta}} V}{2\epsilon^2} - \frac{\inf_{K_2} V}{2\epsilon^2} - \lambda_{\epsilon,2}(t-T) \right\} \|f\|_{\infty} \\ & \leq \frac{D_4}{\epsilon} \exp \left\{ \frac{\sup_{\mathcal{A}_{\delta}} V}{2\epsilon^2} - \frac{\inf_{K_2} V}{2\epsilon^2} - \lambda_{\epsilon,2}(t-T) \right\} \|f\|_{\infty}, \quad \forall t > T + 1, \end{aligned} \quad (6.10)$$

where  $D_4 = D_4(K_2) := D_3 \left( \sup_{\mathcal{A}_1} a^{\frac{1}{4}} \right) \left( \int_{K_2} \frac{1}{a} dx \right)^{\frac{1}{2}}$ .

To estimate  $E_2^\epsilon$ , we see from (6.8) that

$$E_2^\epsilon \leq \mathbb{P}_\mu^\epsilon[X_T^\epsilon \in \mathcal{A}_\delta^c] \|f\|_\infty \leq D_1 e^{-\frac{D_2}{\epsilon^2}} \|f\|_\infty, \quad \forall t > T.$$

Since Lemma 6.1 (2) (with  $k = 1$ ) gives  $D_5 > 0$  such that

$$E_2^\epsilon \leq \mathbb{E}_\mu^\epsilon[f(X_t^\epsilon)] \leq \int_0^\infty a^{\frac{1}{4}} e^{\frac{V+D_5}{\epsilon^2}} d\mu e^{-\lambda_{\epsilon,1}(t-T)} \|f\|_\infty, \quad \forall t > T + 2,$$

there exists  $D_6 = D_6(K_1) > 0$  such that

$$E_2^\epsilon \leq \exp\left\{\frac{D_6}{\epsilon^2} - \lambda_{\epsilon,1}(t-T)\right\} \|f\|_\infty, \quad \forall t > T + 2.$$

Clearly, there exist  $p = p(K_1, \delta) > 1$  and  $D_7 = D_7(K_1, \delta) > 0$  such that  $-\frac{D_2}{p} + \frac{D_6}{p'} < -D_7$ , where  $p' = \frac{p}{p-1}$  is the dual exponent of  $p$ . Then,

$$\begin{aligned} E_2^\epsilon &= (E_2^\epsilon)^{\frac{1}{p}} (E_2^\epsilon)^{\frac{1}{p'}} \leq D_1^{\frac{1}{p}} \exp\left\{-\frac{D_2}{p\epsilon^2} + \frac{D_6}{p'\epsilon^2} - \frac{\lambda_{\epsilon,1}(t-T)}{p'}\right\} \|f\|_\infty \\ &\leq D_1^{\frac{1}{p}} \exp\left\{-\frac{D_7}{\epsilon^2} - \frac{\lambda_{\epsilon,1}(t-T)}{p'}\right\} \|f\|_\infty, \quad \forall t > T + 2. \end{aligned} \quad (6.11)$$

Setting  $\hat{\alpha}_\epsilon(\mu, \delta) := \|\phi_{\epsilon,1}\|_{L^1(u_\epsilon^G)} \int_{\mathcal{A}_\delta} \phi_{\epsilon,1} d\mu_T^\epsilon e^{\lambda_{\epsilon,1}T}$ , we find from (6.9), (6.10) and (6.11) that

$$\begin{aligned} \left| \mathbb{E}_\mu^\epsilon[f(X_t^\epsilon)] - \hat{\alpha}_\epsilon(\mu, \delta) e^{-\lambda_{\epsilon,1}t} \int_0^\infty f d\mu_\epsilon \right| &\leq \left| E_1^\epsilon - \hat{\alpha}_\epsilon(\mu, \delta) e^{-\lambda_{\epsilon,1}t} \int_0^\infty f d\mu_\epsilon \right| + E_2^\epsilon \\ &\leq \frac{D_4}{\epsilon} \exp\left\{\frac{\sup_{\mathcal{A}_\delta} V}{2\epsilon^2} - \frac{\inf_{K_2} V}{2\epsilon^2} - \lambda_{\epsilon,2}(t-T)\right\} \|f\|_\infty \\ &\quad + D_1^{\frac{1}{p}} \exp\left\{-\frac{D_7}{\epsilon^2} - \frac{\lambda_{\epsilon,1}(t-T)}{p'}\right\} \|f\|_\infty, \quad \forall t > T + 2. \end{aligned}$$

It remains to estimate  $\hat{\alpha}_\epsilon(\mu, \delta)$ . Note that


$$\begin{aligned} \hat{\alpha}_\epsilon(\mu, \delta) &\leq \|\phi_{\epsilon,1}\|_{L^1(u_\epsilon^G)} \int_0^\infty \phi_{\epsilon,1} d\mu_T^\epsilon e^{\lambda_{\epsilon,1}T} \\ &= \|\phi_{\epsilon,1}\|_{L^1(u_\epsilon^G)} \int_0^\infty P_T^\epsilon \phi_{\epsilon,1} d\mu e^{\lambda_{\epsilon,1}T} = \|\phi_{\epsilon,1}\|_{L^1(u_\epsilon^G)} \int_0^\infty \phi_{\epsilon,1} d\mu = \alpha_\epsilon(\mu). \end{aligned} \quad (6.12)$$

The upper bound of  $\hat{\alpha}_\epsilon(\mu, \delta)$  follows from that of  $\alpha_\epsilon(\mu)$  given in (1). This completes the proof.  $\square$

## 7. Applications

In this section, we discuss some applications of Corollary A and Theorem C. Two typical situations where SDEs of the form (1.1) arise are described as follows.

**Chemical reactions.** Consider the chemical reactions:



where the positive numbers  $k_1$ ,  $k_{-1}$  and  $k_2$  are reaction rates. The concentration of the substance  $A$ , denoted by  $x_A$ , is assumed to remain constant. We assume  $k_1 x_A > k_2$ .

Denote by  $V \gg 1$  the volume of the system and  $X_t^V$  be the continuous-time Markov jump process counting the number of molecules of the substance  $X$ . The law of large numbers [19, 1] ensures that

as the volume  $V$  grows to infinity, the re-scaled process  $\frac{X_t^V}{V}$  converges to solutions of the following classical mean-field ODE for the concentration of  $X$ :

$$\dot{x} = -k_{-1}x^2 + k_1x_Ax - k_2x, \quad x \in [0, \infty). \quad (7.1)$$

Obviously, (7.1) is a standard logistic equation, and its unique positive equilibrium is given by  $x_e := \frac{k_1x_A - k_2}{k_{-1}}$ . The fluctuation of  $\frac{X_t^V}{V}$  around solutions of (7.1) is captured by the central limit theorem [19, 1], leading to the following *diffusion approximation* of  $\frac{X_t^V}{V}$ :

$$dx = (-k_{-1}x^2 + k_1x_Ax - k_2x)dt + \epsilon\sqrt{k_{-1}x^2 + k_1x_Ax + k_2x}dW_t, \quad x \in [0, \infty), \quad (7.2)$$

where  $\epsilon = \frac{1}{\sqrt{V}}$  and  $W_t$  is a standard one-dimensional Wiener process.

It is a routine task to check (see e.g. [31]) that solutions of (7.2) hit the extinction state 0 in finite time almost surely, while positive solutions of (7.1) are attracted by the equilibrium  $x_e$ . Such a dynamical disagreement between a mean-field model and its stochastic counterpart is often referred to as *Keizer's paradox* [33]. This is originally formulated in terms of the master equation for distributions of  $\frac{X_t^V}{V}$  (see e.g. [34, 55, 7]). In the work [7], the authors use a slow manifold reduction method to estimate the first eigenvalue and the gap between the first and second eigenvalues of the generator of  $\frac{X_t^V}{V}$  that respectively quantify the first passage time to the extinction state and QSD.

**Logistic BDPs.** Let  $\mathbb{N}_0 = \mathbb{N} \cup \{0\}$  and  $\lambda > \mu > 0$ . Consider a continuous-time birth-and-death process (BDP)  $Z_t^K$  on the state space  $\mathbb{N}_0$  with birth rates  $\lambda_n^K = \lambda n$ ,  $n \in \mathbb{N}_0$  and death rates  $\mu_n^K = n(\mu + \frac{n}{K})$ ,  $n \in \mathbb{N}$ , where  $K \gg 1$  is the carrying capacity. By the central limit theorem (see e.g. [38, 19]), for sufficiently large  $K$ , the process  $\frac{Z_t^K}{K}$  stays close to solutions of the following SDE:

$$dx = (\lambda x - \mu x - x^2)dt + \epsilon\sqrt{\lambda x + \mu x + x^2}dW_t, \quad x \in [0, \infty), \quad (7.3)$$

where  $\epsilon = \frac{1}{\sqrt{K}}$ . The SDE (7.3) is the diffusion approximation of  $\frac{Z_t^K}{K}$ .

Both situations give rise to SDEs of the form :

$$dx = (b_1x - b_2x^2)dt + \epsilon\sqrt{a_1x + a_2x^2}dW_t, \quad x \in [0, \infty), \quad (7.4)$$

where  $0 < \epsilon \ll 1$ ,  $b_1, b_2$  and  $a_1$  are positive constants, and  $a_2 \geq 0$ . It is straightforward to check that assumptions in Corollary A and Proposition A.1 are satisfied. Applying Corollary A, Theorem C and Proposition A.1 to (7.4), we find the following result.

**Theorem 7.1.** *The following statements hold.*

(1) *There hold*

$$-\lim_{\epsilon \rightarrow 0} \epsilon^2 \ln \lambda_{\epsilon,1} = r_1 := \begin{cases} 2 \left[ \frac{a_1b_2 + a_2b_1}{a_2^2} \ln \left( 1 + \frac{a_2b_1}{a_1b_2} \right) - \frac{b_1}{a_2} \right], & \text{if } a_2 > 0, \\ \frac{b_1^2}{a_1b_2}, & \text{if } a_2 = 0 \end{cases}$$

and  $\lim_{\epsilon \rightarrow 0} \lambda_{\epsilon,2} = b_1$ .

(2) *For each compact  $K \subset [0, \infty)$ , there are  $C = C(K) > 0$  and small  $\epsilon_* = \epsilon_*(K) > 0$  such that*

$$\begin{aligned} & \sup_{\substack{\mu \in \mathcal{P}([0, \infty)) \\ \text{supp}(\mu) \subset K}} \left\| \mathbb{P}_\mu^\epsilon [X_t^\epsilon \in \bullet] - [e^{-\lambda_{\epsilon,1}t} \mu_\epsilon + (1 - e^{-\lambda_{\epsilon,1}t}) \delta_0] \right\|_{TV} \\ & \leq \exp \left\{ \frac{C}{\epsilon^2} - \frac{b_1}{2} t \right\} + \gamma(\epsilon) e^{-\lambda_{\epsilon,1}t}, \quad \forall t > 0 \text{ and } \epsilon \in (0, \epsilon_*), \end{aligned}$$

where  $\gamma = \gamma_K : (0, 1) \rightarrow (0, 1)$  satisfies  $\gamma(\epsilon) \rightarrow 0$  as  $\epsilon \rightarrow 0$ .

- (3) For each  $0 < \delta \ll 1$  and compact sets  $K_1, K_2 \subset (0, \infty)$  with  $(\frac{b_1}{b_2} - \delta, \frac{b_1}{b_2} + \delta) \cap K_2 = \emptyset$ , there are  $T = T(K_1, \delta) > 0$ ,  $C = C(K_1, K_2, \delta) > 0$  and  $\epsilon_* = \epsilon_*(K_1, K_2, \delta) > 0$  such that for each continuous function  $f : (0, \infty) \rightarrow \mathbb{R}$  with  $\text{supp}(f) \subset K_2$ , there holds

$$\sup_{\substack{\mu \in \mathcal{P}([0, \infty)) \\ \text{supp}(\mu) \subset K_1}} \left| \mathbb{E}_\mu^\epsilon [f(X_t^\epsilon)] - e^{-\lambda_{\epsilon,1} t} \int_0^\infty f d\mu_\epsilon \right| \leq \Gamma(\epsilon) e^{-C\lambda_{\epsilon,1} t} \|f\|_\infty, \quad \forall t > T \text{ and } \epsilon \in (0, \epsilon_*),$$

where  $\Gamma = \Gamma_{K_1, K_2, \delta} : (0, 1) \rightarrow (0, 1)$  satisfies  $\Gamma(\epsilon) \rightarrow 0$  as  $\epsilon \rightarrow 0$ .

Note that  $\frac{a_1 b_2 + a_2 b_1}{a_2} \ln \left( 1 + \frac{a_2 b_1}{a_1 b_2} \right) = \frac{a_1 b_2}{a_2} \left( 1 + \frac{a_2 b_1}{a_1 b_2} \right) \ln \left( 1 + \frac{a_2 b_1}{a_1 b_2} \right) > \frac{a_1 b_2}{a_2} \frac{a_2 b_1}{a_1 b_2} = \frac{b_1}{a_2}$ .

We end this section by stating the following remark about the validity of diffusion approximations of  $\frac{X_t^V}{V}$  and  $\frac{Z_t^K}{K}$ .

**Remark 7.1.** *In general, diffusion approximations of re-scaled Markov jump processes can be justified only over finite time intervals of the form  $[0, T]$ . But, for the process  $\frac{X_t^V}{V}$ , its diffusion approximation holds at least over  $[0, T_\epsilon]$ , where  $T_\epsilon = e^{\frac{r}{\epsilon^2}}$  for  $r \in (0, r_1)$ , due to the special dynamical structure of the mean-field ODE, that is, it has a single globally asymptotically stable equilibrium. Indeed, this can be roughly seen from the classical result for diffusion approximations and the following facts:*

- *Theorem 7.4 says that a typical solution of (7.2) quickly approaches the QSD and then is trapped by it until  $T_\epsilon$ ; the process  $\frac{X_t^V}{V}$  exhibits similar dynamics (see [11]);*
- *combining Lemma A.1 (1) with a similar result for the QSD of  $\frac{X_t^V}{V}$  established in [11], we show in the forthcoming work [49] that the distance ( $p$ -Wasserstein distance for any  $p \in [1, \infty)$  and the Kolmogorov metric) between QSDs of  $\frac{X_t^V}{V}$  and (7.2) vanishes as  $\epsilon = \frac{1}{\sqrt{V}} \rightarrow 0$ .*

Similar results hold for  $\frac{Z_t^K}{K}$  and (7.3).

In the case that the mean-field ODE has multiple asymptotically stable equilibria, QSDs of the re-scaled Markov jump process and its diffusion approximation may stay far away from each other (see e.g. [51]), but they always stay close in some weak sense to the attractor of the mean-field ODE (see e.g. [32, 20, 53]), which is traditionally used to explain the chemical oscillations in a closed chemical reaction system or the persistence states of a species from a phenomenological point of view. Therefore, QSDs can be used to describe the long transient chemical oscillations in a closed chemical reaction system or the long persistence of a species in this case.

## Acknowledgements

The authors would like to express their sincere thanks to the anonymous referee for carefully reading the manuscript and providing invaluable suggestions.

## APPENDIX A. Asymptotic of the first eigenfunctions

We justify (1.6) when  $b$  changes its sign only once appealing to results established in the forthcoming work [49]. The formal statement is as follows. Recall  $\alpha_\epsilon(\mu)$  and  $\hat{\alpha}_\epsilon(\mu, \delta)$  from Theorem C.

**Proposition A.1.** *Assume (H). Suppose, in addition, that*

- $b \in C^2((0, \infty))$ ,  $b$  changes its sign only at  $x^* \in (0, \infty)$  and  $b'(x^*) < 0$ ;
- $a \in C^3((0, \infty))$  and  $\int_1^\infty \frac{1}{\sqrt{a(s)}} ds = \infty$ .

Then,  $\lim_{\epsilon \rightarrow 0} \|\phi_{\epsilon,1}\|_{L^1(u_\epsilon^G)} \phi_{\epsilon,1} = 1$  locally uniformly in  $(0, \infty)$ . In particular,

(1) for any compact  $K \subset [0, \infty)$ ,

$$\lim_{\epsilon \rightarrow 0} \alpha_\epsilon(\mu) = 1 \quad \text{uniformly in } \mu \in \mathcal{P}([0, \infty)) \text{ with } \text{supp}(\mu) \subset K;$$

(2) for any compact  $K \subset (0, \infty)$  and  $0 < \delta_1 < \delta_2 \ll 1$ ,

$$\lim_{\epsilon \rightarrow 0} \hat{\alpha}_\epsilon(\mu, \delta) = 1 \quad \text{uniformly in } \mu \in \mathcal{P}([0, \infty)) \text{ with } \text{supp}(\mu) \subset K \text{ and } \delta \in [\delta_1, \delta_2].$$

We remark that the condition  $\int_1^\infty \frac{1}{\sqrt{a(s)}} ds = \infty$  is only to make  $y_\infty = \lim_{x \rightarrow \infty} \xi(x) = \infty$ , where  $\xi$  is defined in Subsection 2.2, so that the Schrödinger operator  $\mathcal{L}_\epsilon^S$  is on  $(0, \infty)$ . Then, we can use decaying properties of its eigenfunctions near  $\infty$ .

Recall that  $u_\epsilon = \frac{\phi_{\epsilon,1} u_\epsilon^G}{\|\phi_{\epsilon,1}\|_{L^1(u_\epsilon^G)}}$  is the density of the unique QSD  $\mu_\epsilon$  of  $X_t^\epsilon$ . We need the following results from [49], whose proof indeed benefits from decaying properties of  $\tilde{U}_\epsilon U_\epsilon \phi_{\epsilon,1}$ , an eigenfunction of  $\mathcal{L}_\epsilon^S$  associated to  $\lambda_{\epsilon,1}$ , near  $\infty$ , where  $\tilde{U}_\epsilon$  and  $U_\epsilon$  are unitary transforms defined in Subsection 2.2.

**Lemma A.1** ([49]). *Suppose the conditions in Proposition A.1 hold.*

(1) There holds  $u_\epsilon = \frac{R_\epsilon}{\epsilon a} e^{-\frac{2}{\epsilon^2} \int_\bullet^{x_*} \frac{b(s)}{a(s)} ds}$ , where

$$\lim_{\epsilon \rightarrow 0} R_\epsilon(x) = R_0 := \sqrt{-\frac{a(x_*)b'(x_*)}{\pi}} \quad \text{locally uniformly in } x \in (0, \infty).$$

(2) For each  $0 < \delta \ll 1$ , there are  $0 < x_\delta \ll 1$  and  $0 < \epsilon_\delta \ll 1$  such that

$$e^{-\frac{2}{\epsilon^2} \left( \int_0^{x_*} \frac{b(s)}{a(s)} ds + \delta \right)} \leq u_\epsilon(x) \leq e^{-\frac{2}{\epsilon^2} \left( \int_0^{x_*} \frac{b(s)}{a(s)} ds - \delta \right)}, \quad \forall x \in (0, x_\delta), \quad \epsilon \in (0, \epsilon_\delta).$$

(3) There exist  $L \gg 1$ ,  $C > 0$ ,  $r_* > 0$  and  $0 < \epsilon_* \ll 1$  such that

$$u_\epsilon(x) \leq \frac{C}{[a(x)]^{\frac{3}{4}}} e^{-\frac{r_*}{\epsilon^2} (\xi(x) - \xi(L))} e^{\frac{1}{\epsilon^2} \int_L^x \frac{b(s)}{a(s)} ds}, \quad \forall x \in [L, \infty), \quad \epsilon \in (0, \epsilon_*).$$

Note that Lemma A.1 (1) justifies the WKB approximation of  $u_\epsilon$  up to the second order.

We prove Proposition A.1.

**Proof of Proposition A.1.** Lemma A.1 (1) gives  $\frac{\phi_{\epsilon,1} u_\epsilon^G}{\|\phi_{\epsilon,1}\|_{L^1(u_\epsilon^G)}} = \frac{R_\epsilon}{\epsilon a} e^{-\frac{2}{\epsilon^2} \int_\bullet^{x_*} \frac{b(s)}{a(s)} ds}$ , which together with the expression for  $u_\epsilon^G$  leads to

$$\phi_{\epsilon,1} = \frac{1}{\epsilon} e^{-\frac{2}{\epsilon^2} \int_0^{x_*} \frac{b(s)}{a(s)} ds} \|\phi_{\epsilon,1}\|_{L^1(u_\epsilon^G)} R_\epsilon. \quad (\text{A.1})$$

It follows that

$$1 = \int_0^\infty \phi_{\epsilon,1}^2 u_\epsilon^G dx = \|\phi_{\epsilon,1}\|_{L^1(u_\epsilon^G)} \int_0^\infty u_\epsilon \phi_{\epsilon,1} dx = \frac{1}{\epsilon} e^{-\frac{2}{\epsilon^2} \int_0^{x_*} \frac{b(s)}{a(s)} ds} \|\phi_{\epsilon,1}\|_{L^1(u_\epsilon^G)}^2 \int_0^\infty u_\epsilon R_\epsilon dx. \quad (\text{A.2})$$

We claim  $\lim_{\epsilon \rightarrow 0} \int_0^\infty u_\epsilon R_\epsilon dx = R_0$ . Indeed,  $u_\epsilon R_\epsilon = \epsilon a u_\epsilon^2 e^{\frac{2}{\epsilon^2} \int_\bullet^{x_*} \frac{b(s)}{a(s)} ds}$  thanks to Lemma A.1 (1). Let  $K \gg 1$ . It is easy to see from Lemma A.1 (2) that  $\lim_{\epsilon \rightarrow 0} \int_0^{\frac{1}{K}} u_\epsilon R_\epsilon dx = 0$ . By Lemma A.1 (3),

$$\begin{aligned} \int_K^\infty u_\epsilon R_\epsilon dx &\leq \epsilon C^2 e^{\frac{2\gamma_*}{\epsilon^2} \xi(L)} e^{\frac{2}{\epsilon^2} \int_L^{x_*} \frac{b(s)}{a(s)} ds} \int_K^\infty \frac{1}{\sqrt{a(x)}} e^{-\frac{2\gamma_*}{\epsilon^2} \xi(x)} dx \\ &= \epsilon C^2 e^{\frac{2\gamma_*}{\epsilon^2} \xi(L)} e^{\frac{2}{\epsilon^2} \int_L^{x_*} \frac{b(s)}{a(s)} ds} \int_{\xi(K)}^\infty e^{-\frac{2\gamma_*}{\epsilon^2} y} dy \rightarrow 0 \quad \text{as } \epsilon \rightarrow 0. \end{aligned}$$

Hence,  $\lim_{\epsilon \rightarrow 0} \int_0^\infty u_\epsilon R_\epsilon dx = \lim_{\epsilon \rightarrow 0} \int_{\frac{1}{K}}^K u_\epsilon R_\epsilon dx$ . Since  $\int_0^\infty u_\epsilon dx = 1$  and Lemma A.1 (2)(3) gives  $\lim_{\epsilon \rightarrow 0} \int_{\frac{1}{K}}^K u_\epsilon dx = 1$ , we conclude from Lemma A.1 (1) that  $\lim_{\epsilon \rightarrow 0} \int_{\frac{1}{K}}^K u_\epsilon R_\epsilon dx = R_0$ . This proves the claim.

Letting  $\epsilon \rightarrow 0$  in (A.2), the claim implies  $\lim_{\epsilon \rightarrow 0} \frac{1}{\epsilon} e^{-\frac{2}{\epsilon^2} \int_0^{x^*} \frac{b(s)}{a(s)} ds} \|\phi_{\epsilon,1}\|_{L^1(u_\epsilon^G)}^2 = \frac{1}{R_0}$ , which together with (A.1) and Lemma A.1 (1) yields the local uniform convergence of  $\|\phi_{\epsilon,1}\|_{L^1(u_\epsilon^G)} \phi_{\epsilon,1}$  to 1 as  $\epsilon \rightarrow 0$ .

(1) follows readily. To prove (2), we recall that  $\hat{\alpha}_\epsilon(\mu, \delta) := \|\phi_{\epsilon,1}\|_{L^1(u_\epsilon^G)} \int_{\mathcal{A}_\delta} \phi_{\epsilon,1} d\mu_T^\epsilon e^{\lambda_{\epsilon,1} T}$ , where  $T = T(K, \delta)$  is the smallest time such that  $\sup_{x_0 \in K} \text{dist}(x_t, \mathcal{A}) < \frac{\delta}{2}$  for all  $t > T$ , and  $\mu_T^\epsilon = \mathbb{P}_\mu^\epsilon[X_T^\epsilon \in \bullet]$ .

Let  $\eta_\delta : (0, \infty) \rightarrow [0, 1]$  be a smooth function satisfying  $\eta_\delta = 1$  on  $\mathcal{A}_{\frac{3\delta}{4}}$  and  $\eta_\delta = 0$  on  $\mathcal{A}_\delta^c$ . Then,

$$\begin{aligned} \hat{\alpha}_\epsilon(\mu, \delta) &\geq e^{\lambda_{\epsilon,1} T} \|\phi_{\epsilon,1}\|_{L^1(u_\epsilon^G)} \int_0^\infty \eta_\delta \phi_{\epsilon,1} d\mu_T^\epsilon \\ &= e^{\lambda_{\epsilon,1} T} \|\phi_{\epsilon,1}\|_{L^1(u_\epsilon^G)} \int_0^\infty P_T^\epsilon(\eta_\delta \phi_{\epsilon,1}) d\mu = e^{\lambda_{\epsilon,1} T} \int_0^\infty \mathbb{E}_\bullet^\epsilon [(\eta_\delta \|\phi_{\epsilon,1}\|_{L^1(u_\epsilon^G)} \phi_{\epsilon,1})(X_T^\epsilon) \mathbb{1}_{T < T_0^\epsilon}] d\mu, \end{aligned}$$

where we used Proposition 2.1 (4) in the second equality. Note that

$$\begin{aligned} &\mathbb{E}_\bullet^\epsilon [(\eta_\delta \|\phi_{\epsilon,1}\|_{L^1(u_\epsilon^G)} \phi_{\epsilon,1})(X_T^\epsilon) \mathbb{1}_{T < T_0^\epsilon}] \\ &\geq \int_{[X_T^\epsilon \in \mathcal{A}_{\frac{3\delta}{4}}]} (\eta_\delta \|\phi_{\epsilon,1}\|_{L^1(u_\epsilon^G)} \phi_{\epsilon,1})(X_T^\epsilon) \mathbb{1}_{T < T_0^\epsilon} d\mathbb{P}_\bullet^\epsilon \\ &\geq \left( \inf_{\mathcal{A}_{\frac{3\delta}{4}}} \|\phi_{\epsilon,1}\|_{L^1(u_\epsilon^G)} \phi_{\epsilon,1} \right) \mathbb{P}_\bullet^\epsilon [X_T^\epsilon \in \mathcal{A}_{\frac{3\delta}{4}}, T < T_0^\epsilon] \\ &\geq \left( \inf_{\mathcal{A}_{\frac{3\delta}{4}}} \|\phi_{\epsilon,1}\|_{L^1(u_\epsilon^G)} \phi_{\epsilon,1} \right) \left\{ \mathbb{P}_\bullet^\epsilon [X_T^\epsilon \in \mathcal{A}_{\frac{3\delta}{4}}] + \mathbb{P}_\bullet^\epsilon [T < T_0^\epsilon] - 1 \right\}. \end{aligned}$$

Since the limits  $\lim_{\epsilon \rightarrow 0} e^{\lambda_{\epsilon,1} T} = 1$ ,  $\lim_{\epsilon \rightarrow 0} \mathbb{P}_x^\epsilon [X_T^\epsilon \in \mathcal{A}_{\frac{3\delta}{4}}] = 1$  and  $\lim_{\epsilon \rightarrow 0} \mathbb{P}_x^\epsilon [T < T_0^\epsilon] = 1$  are uniform in  $x \in K$  and  $\delta \in [\delta_1, \delta_2]$ , we conclude the lower bound:

$$\liminf_{\epsilon \rightarrow 0} \hat{\alpha}_\epsilon(\mu, \delta) \geq 1 \quad \text{uniformly in } \mu \in \mathcal{P}([0, \infty)) \text{ with } \text{supp}(\mu) \subset K \text{ and } \delta \in [\delta_1, \delta_2].$$

The upper bound follows from  $\hat{\alpha}_\epsilon(\mu, \delta) \leq \alpha_\epsilon(\mu)$  due to (6.12), and the limit proven in (1) for  $\alpha_\epsilon(\mu)$ . This completes the proof.  $\square$

## REFERENCES

1. D. F. Anderson and T. G. Kurtz, *Stochastic analysis of biochemical systems*. Mathematical Biosciences Institute Lecture Series. Stochastics in Biological Systems, 1.2. Springer, Cham; MBI Mathematical Biosciences Institute, Ohio State University, Columbus, OH, 2015.
2. F. A. Berezin and M. A. Shubin, The Schrödinger equation. Translated from the 1983 Russian edition by Yu. Rajabov, D. A. Leites and N. A. Sakharova and revised by Shubin. With contributions by G. L. Litvinov and Leites. Mathematics and its Applications (Soviet Series), 66. Kluwer Academic Publishers Group, Dordrecht, 1991.
3. A. Bovier, M. Eckhoff, V. Gayrard and M. Klein, Metastability and low lying spectra in reversible Markov chains. *Comm. Math. Phys.* 228 (2002), no. 2, 219-255.
4. A. Bovier, V. Gayrard and M. Klein, Metastability in reversible diffusion processes. II. Precise asymptotics for small eigenvalues. *J. Eur. Math. Soc. (JEMS)* 7 (2005), no. 1, 69-99.
5. P. Cattiaux, P. Collet, A. Lambert, S. Martínez, S. Méléard and J. San Martín, Quasi-stationary distributions and diffusion models in population dynamics. *Ann. Probab.* 37 (2009), no. 5, 1926-1969.
6. P. Cattiaux and S. Méléard, Competitive or weak cooperative stochastic Lotka-Volterra systems conditioned on non-extinction. *J. Math. Biol.* 60 (2010), no. 6, 797-829.

7. P. Childs and J. P. Keener, Slow manifold reduction of a stochastic chemical reaction: exploring Keizer's paradox. *Discrete Contin. Dyn. Syst. Ser. B* 17 (2012), no. 6, 1775-1794.
8. N. Champagnat and D. Villemonais, Uniform convergence of conditional distributions for absorbed one-dimensional diffusions. *Adv. in Appl. Probab.* 50 (2018), no. 1, 178-203.
9. N. Champagnat and D. Villemonais, General criteria for the study of quasi-stationarity. <https://arxiv.org/pdf/1712.08092.pdf>
10. N. Champagnat and D. Villemonais, Lyapunov criteria for uniform convergence of conditional distributions of absorbed Markov processes. <https://arxiv.org/pdf/1704.01928.pdf>
11. J.-R. Chazottes, P. Collet and S. Méléard, Sharp asymptotics for the quasi-stationary distribution of birth-and-death processes. *Probab. Theory Related Fields* 164 (2016), no. 1-2, 285-332.
12. J.-R. Chazottes, P. Collet and S. Méléard, On time scales and quasi-stationary distributions for multitype birth-and-death processes. *Ann. Inst. Henri Poincaré Probab. Stat.* 55 (2019), no. 4, 2249-2294.
13. P. Collet, S. Martínez and J. San Martín, *Quasi-stationary distributions. Markov chains, diffusions and dynamical systems*. Probability and its Applications (New York). Springer, Heidelberg, 2013.
14. J.-M. Combes, P. Duclos and R. Seiler, Kreĭn's formula and one-dimensional multiple-well. *J. Funct. Anal.* 52 (1983), no. 2, 257-301.
15. A. Dembo and O. Zeitouni, *Large deviations techniques and applications*. Second edition. Applications of Mathematics (New York), 38. Springer-Verlag, New York, 1998.
16. A. Devinatz, R. Ellis and A. Friedman, The asymptotic behavior of the first real eigenvalue of second order elliptic operators with a small parameter in the highest derivatives. II. *Indiana Univ. Math. J.* 23 (1973/74), 991-1011.
17. A. Eizenberg and Y. Kifer, The asymptotic behavior of the principal eigenvalue in a singular perturbation problem with invariant boundaries. *Probab. Theory Related Fields* 76 (1987), no. 4, 439-476.
18. K.-J. Engel and R. Nagel, *One-parameter semigroups for linear evolution equations*. With contributions by S. Brendle, M. Campiti, T. Hahn, G. Metafune, G. Nickel, D. Pallara, C. Perazzoli, A. Rhandi, S. Romanelli and R. Schnaubelt. Graduate Texts in Mathematics, 194. Springer-Verlag, New York, 2000.
19. S. N. Ethier and T. G. Kurtz, *Markov processes. Characterization and convergence*. Wiley Series in Probability and Mathematical Statistics: Probability and Mathematical Statistics. John Wiley & Sons, Inc., New York, 1986.
20. M. Faure and S. J. Schreiber, Quasi-stationary distributions for randomly perturbed dynamical systems. *Ann. Appl. Probab.* 24 (2014), no. 2, 553-598.
21. M. I. Freidlin and A. D. Wentzell, *Random perturbations of dynamical systems*. Translated from the 1979 Russian original by Joseph Szűcs. Third edition. Grundlehren der Mathematischen Wissenschaften [Fundamental Principles of Mathematical Sciences], 260. Springer, Heidelberg, 2012.
22. A. Friedman, The asymptotic behavior of the first real eigenvalue of a second order elliptic operator with a small parameter in the highest derivatives. *Indiana Univ. Math. J.* 22 (1972/73), 1005-1015.
23. M. Fukushima, *Dirichlet forms and Markov processes*. North-Holland Mathematical Library, 23. North-Holland Publishing Co., Amsterdam-New York; Kodansha, Ltd., Tokyo, 1980.
24. A. Galves, E. Olivieri and M.E Vares, Metastability for a class of dynamical systems subject to small random perturbations. *Ann. Probab.* 15 (1987), no. 4, 1288-1305.
25. A. Hastings, Transients: The key to long-term ecological understanding? *Trends Ecol. Evol.* 19 (2004), no. 1, 39-45.
26. A. Hastings, K. C. Abbott, K. Cuddington, T. Francis, G. Gellner, Y. C. Lai, et al. Transient phenomena in ecology. *Science*. 361 (2018), No. 6406, eaat6412.
27. B. Helffer, M. Klein and F. Nier, Quantitative analysis of metastability in reversible diffusion processes via a Witten complex approach. *Mat. Contemp.* 26 (2004), 41-85.
28. B. Helffer and F. Nier, Quantitative analysis of metastability in reversible diffusion processes via a Witten complex approach: the case with boundary. *Mém. Soc. Math. Fr. (N.S.)* No. 105 (2006), vi+89 pp.
29. A. Hening and M. Kolb, Quasistationary distributions for one-dimensional diffusions with singular boundary points. *Stochastic Process. Appl.* 129 (2019), no. 5, 1659-1696.
30. R. A. Holley, S. Kusuoka and D. W. Stroock, Asymptotics of the spectral gap with applications to the theory of simulated annealing. *J. Funct. Anal.* 83 (1989), no. 2, 333-347.
31. N. Ikeda and S. Watanabe, *Stochastic differential equations and diffusion processes*. Second edition. North-Holland Mathematical Library, 24. North-Holland Publishing Co., Amsterdam; Kodansha, Ltd., Tokyo, 1989.
32. F. Jacobs and S. J. Schreiber, Random perturbations of dynamical systems with absorbing states. *SIAM J. Appl. Dyn. Syst.* 5 (2006), no. 2, 293-312.



33. J. Keizer, Master equations, Langevin equations, and the effect of diffusion on concentration fluctuations. *J. Chem. Phys.* 67 (1977), no. 4, 1473-1476.
34. J. Keizer, *Statistical Thermodynamics of Nonequilibrium Processes*. Springer-Verlag, New York, 1987.
35. Y. Kifer, On the principal eigenvalue in a singular perturbation problem with hyperbolic limit points and circles. *J. Differential Equations* 37 (1980), no. 1, 108-139.
36. Y. Kifer, The exit problem for small random perturbations of dynamical systems with a hyperbolic fixed point. *Israel J. Math.* 40 (1981), no. 1, 74-96.
37. V. N. Kolokoltsov, *Semiclassical analysis for diffusions and stochastic processes*. Lecture Notes in Mathematics, 1724. Springer-Verlag, Berlin, 2000.
38. T. G. Kurtz, Limit theorems for sequences of jump Markov processes approximating ordinary differential processes. *J. Appl. Probability* 8 (1971), 344-356.
39. Y.-C. Lai and T. Tél, *Transient chaos. Complex dynamics on finite-time scales*. Applied Mathematical Sciences, 173. Springer, New York, 2011.
40. D. Le Peutrec, Small eigenvalues of the Neumann realization of the semiclassical Witten Laplacian. *Ann. Fac. Sci. Toulouse Math. (6)* 19 (2010), no. 3-4, 735-809.
41. J. Littin C., Uniqueness of quasistationary distributions and discrete spectra when  $\infty$  is an entrance boundary and 0 is singular. *J. Appl. Probab.* 49 (2012), no. 3, 719-730.
42. P. Mathieu, Zero white noise limit through Dirichlet forms, with application to diffusions in a random medium. *Probab. Theory Related Fields* 99 (1994), no. 4, 549-580.
43. P. Mathieu, Spectra, exit times and long time asymptotics in the zero-white-noise limit. *Stochastics Stochastics Rep.* 55 (1995), no. 1-2, 1-20.
44. S. Méléard and D. Villemonais, Quasi-stationary distributions and population processes. *Probab. Surv.* 9 (2012), 340-410.
45. L. Miclo, Comportement de spectres d'opérateurs de Schrödinger à basse température. (French) [Low-temperature behavior of spectra of Schrödinger operators] *Bull. Sci. Math.* 119 (1995), no. 6, 529-553.
46. Y. Miura, Ultracontractivity for Markov semigroups and quasi-stationary distributions. *Stoch. Anal. Appl.* 32 (2014), no. 4, 591-601.
47. A. Morozov, et al., Long transients in ecology: Theory and applications. *Phys. Life Rev.* 32 (2020) 1-40.
48. E. Olivieri and M. E. Vares, *Large deviations and metastability*. Encyclopedia of Mathematics and its Applications, 100. Cambridge University Press, Cambridge, 2005.
49. W. Qi, Z. Shen and Y. Yi, Large deviation principle of quasi-stationary distributions for absorbed singular diffusions, in preparation.
50. A. Pazy, *Semigroups of linear operators and applications to partial differential equations*. Applied Mathematical Sciences, 44. Springer-Verlag, New York, 1983.
51. H. Qian, Nonlinear stochastic dynamics of mesoscopic homogeneous biochemical reaction systems - an analytical theory. *Nonlinearity* 24 (2011), no. 6, R19-R49.
52. S. K. Scott, B. Peng, A. S. Tomlin and K. Showalter, Transient chaos in a closed chemical system. *J. Chem. Phys.* 94, 1134 (1991).
53. Z. Shen, S. Wang and Y. Yi, Concentration of quasi-stationary distributions for one-dimensional diffusions with applications. <https://sites.ualberta.ca/~zhongwei/manuscript-Shen-Wang-Yi.pdf>.
54. B. Simon, Semiclassical analysis of low lying eigenvalues. I. Nondegenerate minima: asymptotic expansions. *Ann. Inst. H. Poincaré Sect. A (N.S.)* 38 (1983), no. 3, 295-308.
55. M. Vellela and H. Qian, A quasistationary analysis of a stochastic chemical reaction: Keizer's paradox. *Bull. Math. Biol.* 69 (2007), no. 5, 1727-1746.
56. A. D. Ventcel, The asymptotic behavior of the largest eigenvalue of a second order elliptic differential operator with a small parameter multiplying the highest derivatives. (Russian) *Dokl. Akad. Nauk SSSR* 202 (1972), 19-22.
57. J. Wang, P. G. Soerensen and F. Hynne, Transient period doublings, torus oscillations, and chaos in a closed chemical system. *J. Phys. Chem.* 98 (1994), no. 3, 725-727.

HUA LOO-KENG KEY LABORATORY OF MATHEMATICS, ACADEMY OF MATHEMATICS AND SYSTEMS SCIENCE, BEIJING 100190, PRC, AND SCHOOL OF MATHEMATICAL SCIENCES, UNIVERSITY OF CHINESE ACADEMY OF SCIENCES, BEIJING 100049, PRC

*Email address:* `jimin@math.ac.cn`

DEPARTMENT OF MATHEMATICAL AND STATISTICAL SCIENCES, UNIVERSITY OF ALBERTA, EDMONTON, AB T6G 2G1, CANADA

*Email address:* `wqi2@ualberta.ca`

DEPARTMENT OF MATHEMATICAL AND STATISTICAL SCIENCES, UNIVERSITY OF ALBERTA, EDMONTON, AB T6G 2G1, CANADA

*Email address:* `zhongwei@ualberta.ca`

DEPARTMENT OF MATHEMATICAL AND STATISTICAL SCIENCES, UNIVERSITY OF ALBERTA, EDMONTON, AB T6G 2G1, CANADA, AND SCHOOL OF MATHEMATICS, JILIN UNIVERSITY, CHANGCHUN 130012, PRC

*Email address:* `yingfei@ualberta.ca`